

# Contents

Chapter 1. Introduction	1
1.1. Outline of the proof	4
1.2. Outline of manuscript	8
1.3. Other approaches	8
1.4. Acknowledgements	10
 <b>Part 1. GEOMETRIC AND ANALYTIC RESULTS FOR RICCI FLOW WITH SURGERY</b>	11
Chapter 2. Ricci flow with surgery	13
2.1. The standard solution and surgery	14
2.2. Main existence theorem	15
2.3. Review of notation and definitions	17
2.4. Geometric limits of Ricci flows	21
Chapter 3. Limits as $t \rightarrow \infty$	25
3.1. Preliminary estimates on curvature and volume	25
3.2. Three propositions	29
3.3. The hyperbolic pieces	30
3.4. Locally volume collapsed part of the $(M_t, g(t))$	39
Chapter 4. Local results valid for large time	41
4.1. First local result	41
4.2. Second local result	66
4.3. A corollary	77
Chapter 5. Proofs of the three propositions	79
5.1. Proof of Proposition 3.2.4	79
5.2. Proof of Proposition 3.2.3	85
5.3. Proof of Proposition 3.2.1	86
 <b>Part 2. LOCALLY VOLUME COLLAPSED 3-MANIFOLDS</b>	93
Chapter 6. Introduction to Part II	95
6.1. Seifert fibered manifolds and graph manifolds	95
6.2. The statement	97
6.3. Stronger results	99

Chapter 7. The collapsing theorem	101
7.1. First remarks	101
7.2. The collapsing theorem	103
7.3. Proof that Theorem 7.2.1 implies Theorem 6.2.1	103
Chapter 8. Overview of the rest of the argument	107
Chapter 9. Basics of Gromov-Hausdorff convergence	111
9.1. Limits of compact metric spaces	111
9.2. Limits of complete metric spaces	113
9.3. Manifolds with curvature bounded below	113
Chapter 10. Basics of Alexandrov spaces	117
10.1. Properties of comparison angles	117
10.2. Alexandrov spaces of curvature $\geq 0$	119
10.3. Strainers	124
10.4. Alexandrov balls	125
10.5. The tangent cone	127
10.6. Consequences of the existence of tangent cones	128
10.7. Directional derivatives	131
10.8. Blow-up results	135
10.9. Gromov-Hausdorff limits of balls in the $M_n$	139
Chapter 11. 2-dimensional Alexandrov spaces	141
11.1. Basics	141
11.2. The interior	146
11.3. The boundary	149
11.4. The covering	153
11.5. Transition between the 2- and 1-dimensional parts	161
Chapter 12. 3-dimensional analogues	163
12.1. Regions of $M$ near generic 2-dimensional points	164
12.2. The global $S^1$ -fibration	168
12.3. Balls centered at points of $\partial M_n$	173
12.4. The interior cone points	175
12.5. Near almost flat boundary points	180
12.6. Boundary points of angle $\leq \pi - \delta$	190
12.7. Balls near open intervals	196
12.8. Determination of the constants	200
Chapter 13. The global result	205
13.1. Part of $M$ close to intervals	205
13.2. Part of $M$ near 2-dimensional Alexandrov spaces	212
13.3. Fixing the 3-balls and attaching solid cylinders	218
13.4. $\epsilon$ -chains	221

13.5.	The Seifert fibration containing $W_2 \setminus \cup_i U(\mathcal{C}_i')$	230
13.6.	Deforming the boundary of $W_2$	232
13.7.	Removing solid tori and solid cylinders from $W_2$	239
13.8.	Completion of the proof	241
<b>Part 3.</b>	<b>THE EQUIVARIANT CASE</b>	<b>243</b>
Chapter 14.	The equivariant case	245
14.1.	The statement	245
14.2.	Preliminary results on compact group actions	246
14.3.	Actions on canonical neighborhoods	251
14.4.	Equivariant Ricci flow with surgery	261
14.5.	Proof of Theorem 14.1.4.	271
	Bibliography	281
	Glossary of symbols	283
	Index	289

# Completion of the proof of the Geometrization Conjecture

John Morgan and Gang Tian \*

August 7, 2012

## Introduction

This paper builds upon and is an extension of [21]. Here, we complete a proof of the following:

**Geometrization Conjecture:** Any closed, orientable, prime<sup>1</sup> 3-manifold  $M$  contains a disjoint union of embedded incompressible<sup>2</sup> 2-tori and Klein bottles such that each connected component of the complement admits a complete, locally homogeneous Riemannian metric of finite volume.

**Geometric 3-manifolds.** Let us briefly review the nature of *geometric* 3-manifolds, that is to say complete, locally homogeneous Riemannian 3-manifolds of finite volume. Any such manifold is *modelled on* a complete, simply connected homogeneous manifold; that is to say, it is isometric to the quotient of a complete, simply connected homogeneous Riemannian manifold by a discrete group of symmetries acting freely. Here, *homogeneous* means that the isometry group of the manifold acts transitively on the manifold. Geometric 3-manifolds come in eight classes or types depending on the complete, simply connected homogeneous manifold they are modelled on. Here is the list, where, for simplicity we have restricted attention to the orientable case.

1. **Hyperbolic:** These are manifolds of constant negative sectional curvature. The complete, simply connected example of this geometry is hyperbolic 3-space. It can be presented as  $\mathbb{C} \times (0, \infty)$  with coordinates  $(z, y)$  with  $z \in \mathbb{C}$  and  $y \in \mathbb{R}^+$  and with the metric being  $(|dz|^2 + dy^2)/y^2$ . Complete hyperbolic manifolds are the quotients of hyperbolic 3-space by discrete, torsion-free, co-finite volume subgroups of its isometry group  $PSL(2, \mathbb{C})$ . These manifolds can be non-compact; a neighborhood of any end is diffeomorphic to  $T^2 \times [0, \infty)$ ,

---

\*Supported partially by NSF grants DMS 0706815 (Morgan), DMS 0703985 (Tian), and DMS 0735963 (Tian)

<sup>1</sup>Not diffeomorphic to  $S^3$  and with the property that every separating 2-sphere in the manifold bounds a 3-ball.

<sup>2</sup>Meaning the fundamental group of the surface injects into the fundamental group of the 3-manifold.

and the torus cross-sections are all conformally equivalent and have areas that are decaying exponentially fast as we go to infinity.

2. **Flat:** These are manifolds with 0 sectional curvature. They are quotients of  $\mathbb{R}^3$  by discrete torsion-free, co-finite volume subgroups of its isometry group. All such manifolds are compact and are finitely covered by a flat 3-torus.
3. **Round:** These are manifolds with constant positive sectional curvature. They are quotients of  $S^3$  with its natural round metric by finite groups of isometries acting freely. Examples are lens spaces and the Poincaré dodecahedral space.
4. **Modelled on hyperbolic 2-space times  $\mathbb{R}$ :** At every point two of the sectional curvatures are 0 and the third is negative. These manifolds are finitely covered by the product of a hyperbolic surface of finite area with  $S^1$ . There are non-compact examples but every neighborhood of an end of one of these manifolds is diffeomorphic to  $T^2 \times [0, \infty)$  and the torus cross sections have areas that decay exponentially as we go to infinity; one direction is of constant length and the other decays exponentially fast.
5. **Modelled on  $S^2 \times \mathbb{R}$ :** There are exactly two examples here:  $S^2 \times S^1$  and  $\mathbb{R}P^3 \# \mathbb{R}P^3$ .
6. **Modelled on Nil, the 3-dimensional nilpotent group:**

$$\begin{pmatrix} 1 & x & z \\ 0 & 1 & y \\ 0 & 0 & 1 \end{pmatrix}.$$

Any example here is compact and is finitely covered by a non-trivial circle bundle over  $T^2$ .

7. **Modelled on the universal covering of  $PSL_2(\mathbb{R})$ :** Any example is finitely covered by a circle bundle over a hyperbolic surface of finite area. Non-compact examples have ends that are diffeomorphic to  $T^2 \times [0, \infty)$ .
8. **Modelled on Solv, the 3-dimensional solvable group**

$$\mathbb{R}^2 \rtimes \mathbb{R}^*.$$

All examples here are compact and are finitely covered by non-trivial  $T^2$ -bundles over  $S^1$  with gluing diffeomorphism being an element of  $SL(2, \mathbb{Z})$  of whose trace has absolute value  $> 2$ .

Manifolds of the last seven types are easily classified and their classifications have been long known, see [30]. Finite volume hyperbolic 3-manifolds are not classified. It was only recently ([8]) that the hyperbolic 3-manifold of smallest volume was definitely established. It is known that the set of real numbers which are volumes of complete hyperbolic 3-manifolds is totally ordered by the usual order on  $\mathbb{R}$  and

also that the function that associates to a hyperbolic 3-manifold its volume is finite-to-one but not one-to-one. The Geometrization Conjecture reduces the problem of completely classifying 3-manifolds to the problem of classifying complete, finite volume hyperbolic 3-manifolds, or equivalently to classifying torsion-free, co-finite volume lattices in  $PSL(2, \mathbb{C})$ . These problems remain open.

There is another way to organize the list of eight types of geometric 3-manifolds that fits better with what Ricci flow with surgery produces:

1. **Semi-positive type:** compact and modelled on either  $S^3$  or  $S^2 \times \mathbb{R}$ .
2. **Flat:** compact with a flat metric.
3. **Essentially 1-dimensional:** geometric and modelled on Solv.
4. **Essentially 2-dimensional:** the interior of a compact Seifert fibered 3-manifold with incompressible boundary; the interior of the base 2-dimensional orbifold of this Seifert fibration admits a complete hyperbolic or Euclidean metric of finite area. The manifold is geometric and modelled on either the universal covering of  $PSL(2, \mathbb{R})$ , the product of the hyperbolic plane with  $\mathbb{R}$ , or Nil.
5. **Essentially 3-dimensional:** diffeomorphic to a complete hyperbolic 3-manifold of finite volume.

We shall use information about the structure of the *cusps* or neighborhoods of the ends of a finite volume hyperbolic 3-manifold. For any such orientable manifold  $H$  and any end  $\mathcal{E}$  of  $H$  there is a neighborhood of  $\mathcal{E}$  that is isometric to the quotient of subset of the upper half-space

$$\{(z, y) \in \mathbb{C} \times [y_0, \infty) \mid y_0 > 0\}$$

by a lattice subgroup of  $\mathbb{C}$  acting on the first factor by translations and acting trivially on the second factor. The quotients of the slices  $\{y = y_1\}$  are *horospherical tori* in the end. They foliate the neighborhood of the end. Each one of them cuts off a neighborhood of the end that is diffeomorphic to  $T^2 \times [0, \infty)$ . A *truncation* of a complete hyperbolic 3-manifold of finite volume is the compact submanifold obtained by cutting off a neighborhood of each end of the manifold along some horospherical torus in that end.

**Interpretation of the Geometrization Conjecture.** The Geometrization Conjecture can be viewed as saying that any closed, orientable, prime 3-manifold  $M$  maps to a graph  $\Gamma$  in such a way that:

- The map is transverse to the midpoints of the edges and the pre-image of the mid-point of each edge is an incompressible torus in  $M$ .
- Let  $\mathcal{T}$  be the union of the tori that are the pre-images of the midpoints of the edges of the graph, and let  $N$  be the result of cutting  $M$  open along  $\mathcal{T}$  (so that  $N$  is a compact manifold whose boundary consists of two copies of

$\mathcal{T}$ ). The manifold  $N$  naturally maps to the result  $\widehat{\Gamma}$  of cutting  $\Gamma$  open along the midpoints of its edges. This map induces a bijection from the connected components of  $N$  to those of  $\widehat{\Gamma}$ , the latter being naturally indexed by the vertices of  $\Gamma$ .

- Each connected component of  $N$  is either a twisted  $I$ -bundle over the Klein bottle or its interior admits a complete, locally homogeneous metric of finite volume (automatically of one of the eight types listed above).

**Statement for a general closed 3-manifold.** The statement for a general closed, orientable 3-manifold is that there is a two-step process. The first step is to cut the manifold open along a maximal family of essential 2-spheres (essential in the sense that none of the 2-spheres bounds a 3-ball in the manifold and no two of the 2-spheres are parallel in the manifold), and then attach a 3-ball to each boundary component to produce a new closed 3-manifold, each component of which is automatically prime. The second step is to remove a disjoint family of incompressible tori and Klein bottles so that each connected component of the result has a complete, locally homogeneous metric of finite volume. Notice that there is a fundamental difference in these two steps in that in the first one one has to add material (the 3-balls) by hand whereas in the second step nothing is added. By definition, a closed, orientable, connected 3-manifold  $M$  satisfies the Geometrization Conjecture if and only if each of its prime factors does.

**Uniqueness of the decomposition.** Every closed 3-manifold has a decomposition into prime factors and these factors are unique up to order (and diffeomorphism). Given an orientable, prime 3-manifold  $M$ , consider all families of disjointly embedded tori and Klein bottles in  $M$  for which the conclusion of the Geometrization Conjecture holds. We choose one such family  $\mathcal{T}$  with a minimal number of connected surfaces. Then for any other such family  $\mathcal{T}'$  with the same number of connected surfaces as  $\mathcal{T}$  there is isotopic of  $M$  carrying  $\mathcal{T}'$  to  $\mathcal{T}$ . Thus, families  $\mathcal{T}$  which satisfy the Geometrization Conjecture and have a minimal number of connected surfaces are unique up to isotopy. The geometric structures on the complementary components are not unique. For example, for those components that fiber over surfaces or Seifert fiber over two-dimensional orbifolds, there are the moduli of the geometric structure on those surfaces or orbifolds. In addition, there are non-compact examples of types (4) and (7) that are diffeomorphic,

## 0.1 Outline of the proof

The basic ingredient for the proof of the Geometrization Conjecture is the existence and properties of a Ricci flow with surgery. In [21], following Perelman's arguments, we showed that for any closed, oriented Riemannian 3-manifold  $(M_0, g(0))$  there is a Ricci flow with surgery defined for all time with  $(M_0, g(0))$  as the initial condition. This flow consists of a one-parameter family of compact, Riemannian 3-manifolds  $(M_t, g(t))$ , defined for  $0 \leq t < \infty$ . The underlying smooth manifolds are locally

constant and the Riemannian metrics are varying smoothly except for a discrete set  $\{t_i\}$  of surgery times. At these times the topological type of the  $M_t$  and Riemannian metrics  $g(t)$  undergo discontinuous (but highly controlled) changes. One consequence of the nature of these changes is that if  $M_{t_0}$  satisfies the Geometrization Conjecture for some  $t_0 < \infty$ , then  $M_t$  satisfies the Geometrization Conjecture for all  $0 \leq t < \infty$ , and in particular,  $M_0$  satisfies the Geometrization Conjecture.

The strategy for proving the Geometrization Conjecture should now be clear. Start with any closed, oriented 3-manifold  $M_0$ . Impose a Riemannian metric  $g(0)$  and construct the Ricci flow with surgery defined for all  $0 \leq t < \infty$  with  $(M_0, g(0))$  as initial condition. Then show, for all  $t$  sufficiently large, that  $M_t$  satisfies the Geometrization Conjecture. This manuscript concentrates on the topology and geometry of the manifolds  $(M_t, g(t))$  for all  $t$  sufficiently large.

The nicest statement one can imagine is that (after an appropriate rescaling) the Riemannian manifolds  $(M_t, g(t))$  converge smoothly as  $t \rightarrow \infty$  (meaning there are no surgery times for  $t$  sufficiently large and up to diffeomorphism as  $t \rightarrow \infty$  the metrics  $g(t)$  converge smoothly to a limiting metric  $g(\infty)$ ) to a locally homogeneous metric, which is automatically complete and of finite volume since the  $M_t$  are compact. As we shall see, this essentially happens under certain topological assumptions, namely infinite fundamental group which (i) is not a non-trivial free product and (ii) does not contain a non-cyclic abelian subgroup. In this case the limiting metric is hyperbolic. But in general this scenario is too optimistic, not all manifolds are geometric – somehow Ricci flow with surgery must allow for the cutting of the manifold into its prime factors and also allow for the torus decomposition.

A more accurate picture of what happens in general goes as follows. First of all the discontinuities (or surgeries) perform the connected sum decomposition including possibly redundant (i.e., trivial) such decompositions which simply split off new components diffeomorphic to the 3-sphere without changing the topology of the already existing components. For sufficiently large  $t$ , every connected component of  $M_t$  is either prime or diffeomorphic to  $S^3$ . Also, the surgeries remove all components with round metrics and with metrics modelled on  $S^2 \times \mathbb{R}$ . This is the full extent of the topological changes wrought by the surgeries. All of these statements follow from what was established in [21]. Thus, for all sufficiently large  $t$  we have the following: Each connected component of  $M_t$  either is prime or is diffeomorphic to  $S^3$ . Furthermore, if connected component of  $M_t$  has finite fundamental group or has a fundamental group with an infinite cyclic subgroup of finite index, then it is diffeomorphic to  $S^3$ . As we shall show in Part I here, it turns out that given  $(M_0, g(0))$ , there is a finite list of complete hyperbolic manifolds  $\mathcal{H} = H_1 \amalg \cdots \amalg H_k$  such that for any truncation  $\overline{\mathcal{H}}$  of  $\mathcal{H}$  along horospherical tori the following holds. For all  $t$  sufficiently large, there is an embedding  $\varphi_t: \overline{\mathcal{H}} \rightarrow M_t$  such that the rescaled pulled back metrics  $\frac{1}{t}\varphi_t^*g(t)$  converge to the restriction of the hyperbolic metric  $\overline{\mathcal{H}}$ . Furthermore, the image of the boundary tori  $\mathcal{T}$  of  $\overline{\mathcal{H}}$  under  $\varphi_t$  are incompressible tori in  $M_t$ . Lastly, the complement  $(M_t \setminus \varphi_t(\text{int}(\overline{\mathcal{H}})), g(t))$  is locally volume collapsed on the negative curvature scale (details on this notion below). Actually,  $\mathcal{H}$  depends only on the diffeomorphism type of  $M_0$ . The proof of the existence of  $\mathcal{H}$  and the embeddings as required are rescaled versions, valid near infinity, of the main finite-



time results that were used in [21] in the construction of a Ricci flow with surgery and an understanding of its singularity development. These deal with non-collapsing and bounded curvature at bounded distance for the rescaled metrics  $\frac{1}{t}g(t)$  as  $t \rightarrow \infty$ .

To complete the proof of the Geometrization Conjecture we must show that the locally volume collapsed pieces satisfy the appropriate relative version of the Geometrization Conjecture.

**The Relative Version of the Geometrization Conjecture:** Let  $M$  be a compact, orientable 3-manifold whose boundary components are incompressible tori. Suppose that  $M$  is prime in the sense that every 2-sphere in  $M$  bounds a 3-ball and no component of  $M$  is diffeomorphic to  $S^3$ . Then there is a finite disjoint union  $\mathcal{T}$  of incompressible tori and Klein bottles in  $\text{int } M$  such that every connected component of  $\text{int } M \setminus \mathcal{T}$  is either diffeomorphic to  $T^2 \times \mathbb{R}$  or admits a complete, locally homogeneous metric of finite volume.

It is a direct argument to see that the relative version of the conjecture implies the original version of the conjecture when the manifold in question is closed.

### Locally Volume Collapsed manifolds.

**Definition 0.1.** Suppose that  $M$  is a complete  $n$ -dimensional Riemannian manifold and  $w > 0$  and  $\psi: M \rightarrow [0, \infty)$  are given. Then we say that  $M$  is *w locally volume collapsed on scale  $\psi$*  if for every  $x \in M$  we have

$$\text{Vol } B(x, \psi(x)) \leq w\psi(x)^n.$$

**Definition 0.2.** Suppose that  $M$  is a complete, connected Riemannian manifold and that  $M$  does not have everywhere non-negative sectional curvature. Then we define

$$\rho: M \rightarrow [0, \infty)$$

such that for each  $x \in M$  the infimum of the sectional curvatures on  $B(x, \rho(x))$  is  $-\rho^{-2}(x)$ . Then  $\rho(x)$  is the *negative curvature scale* at  $x$ . We say that  $M$  is *w locally volume collapsed on the negative curvature scale* if it is *w locally volume collapsed on scale  $\rho$* .

The results on Ricci flow as  $t \rightarrow \infty$  indicated above produce truncated hyperbolic submanifolds of  $(M_t, g(t))$  whose complements are locally volume collapsed on the negative curvature scale. In fact, given  $w > 0$  for all  $t$  sufficiently large the complement of the hyperbolic pieces in  $(M_t, g(t))$  is *w locally volume collapsed on the negative curvature scale*. The idea for studying the complement is to first understand the balls  $B(x, \rho(x)) \subset M_t$ . Rescaling  $g(t)$  by  $\rho^{-2}(x)$  gives us a unit ball on which the sectional curvatures are bounded below by  $-1$ . This uniform lower curvature bound implies that any sequence of such balls with  $t \rightarrow \infty$  has a subsequence which converges in a weak sense (the Gromov-Hausdorff sense) to a metric space that is weaker than a Riemannian manifold but still has some curvature structure, a so-called Alexandrov space.

Let us briefly list the local models for the limit and the corresponding 3-dimensional models. By general results the Gromov-Hausdorff limit of a sequence of rescaled balls  $\rho^{-1}(x_n)B(x_n, \rho(x_n))$  is an Alexandrov ball of dimension  $\leq 3$  and curvature  $\geq -1$ . The fact that the volume of the  $\rho^{-1}(x_n)B(x_n, \rho(x_n))$  are tending to zero as  $n \rightarrow \infty$ , means that the limit has dimension  $\leq 2$ . Also, it turns out that we can assume that  $\rho(x_n) \leq \text{diameter}(M_n)/2$ . This implies that the limit is not a point and hence has dimension  $\geq 1$ . Thus, the Gromov-Hausdorff limit is either 1- or 2-dimensional.

Let us describe what happens when the limiting Alexandrov space is 1-dimensional. In this case the limit is either an interval (open, half-closed or closed) or a circle. The local structure of the 3-manifolds converging to such Alexandrov space near points converging to an interior point is a product of  $S^2 \times (0, 1)$  or  $T^2 \times (0, 1)$  where the surface fibers are of diameter converging to zero and the interval has length bounded away from zero. In fact we can view neighborhoods in the  $M_n$  as fibering over the limiting open interval with fibers of small diameter which are either  $S^2$ -fibers or  $T^2$ -fibers. Near an end point the structure is either a 3-ball or a punctured  $\mathbb{R}P^3$  (when the fibers over nearby interior points are  $S^2$ ) or a solid torus or a twisted  $I$ -bundle over the Klein bottle (when the fibers over the nearby interior points are 2-tori).

Now we consider the second possibility when the limiting Alexandrov space is 2-dimensional. As we shall see, we write a 2-dimensional Alexandrov space as a union four types of points for an appropriately chosen  $\delta_0 > 0$ :

- interior points that are the center of neighborhoods close to open balls in  $\mathbb{R}^2$ ,
- points at which the space is an almost circular cone of cone angle  $\leq 2\pi - \delta_0$ ,
- boundary points that are the center of neighborhoods close to open balls centered at boundary points of half-space, and
- boundary points at which is space is almost isometric to flat cone in  $\mathbb{R}^2$  of cone angle  $\leq \pi - \delta_0$ .

The local models for neighborhoods of  $x \in M_n$  in these four cases are:

- $S^1 \times \mathbb{R}^2$  with a Riemannian metric that is almost a product of a Riemannian metric on  $S^1$  with a flat Riemannian metric on  $\mathbb{R}^2$ ;
- a solid torus;
- $D^2 \times \mathbb{R}$ ;
- a 3-ball.

It turns out that these neighborhoods are glued together in a completely standard way. It then is an elementary problem in 3-dimensional topology to show that a 3-manifold covered by such neighborhoods intersecting in standard ways satisfies the relative version of the Geometrization Conjecture.

Thus, for all  $t$  sufficiently large, the  $(M_t \setminus \Phi_t(\text{int}\overline{\mathcal{H}}), t^{-1}g(t))$  satisfies the relative version of the Geometrization Conjecture. This then completes the proof of the

Geometrization Conjecture for  $M_t$  for  $t$  sufficiently large, and consequently also for  $M_0$ .

## 0.2 Outline of Manuscript

This manuscript has two parts. In Part I we cover the material in Sections 6 and 7 of [27], in particular the material from Lemma 6.3 through Section 7.3. This preliminary study of the limits as  $t \rightarrow \infty$  of the  $t$  time-slices  $(M_t, g(t))$  of a 3-dimensional Ricci flow with surgery produces a dichotomy. For any  $w > 0$  and for all  $t$  sufficiently large (given  $w$ ), the  $t$  time-slice is divided along incompressible tori into two parts. The first part is a disjoint union of components each of which is an almost complete hyperbolic manifold implying in particular that its interior is diffeomorphic to a complete hyperbolic manifold of finite volume. The second part,  $M_t(w, -)$ , is locally  $w$  volume collapsed on the negative curvature scale. Then we turn to the manifolds  $M_t(w, -)$  for  $w$  sufficiently small and  $t$  sufficiently large. The result we need to handle this case is stated by Perelman as Theorem 7.4 in [27], but no proof is provided in [27]. The second part of this work is devoted to giving a proof of Theorem 7.4 from [27] which is stated as Theorem 5.5 below. We review the background material from the theory of Gromov-Hausdorff convergence of metric spaces and the theory of Alexandrov needed to establish this result. In the final section we state and sketch the proof of the equivariant version of the Geometrization Conjecture for compact group actions on compact 3-manifolds.

## 0.3 Other Approaches

The Geometrization Conjecture was proposed by W. Thurston in early 1980s. It includes the Poincaré Conjecture as a special case. Thurston himself established this conjecture for a large class of 3-manifolds, namely those containing an incompressible surface; i.e., an embedded surface of genus  $\geq 1$  whose fundamental group injects into the fundamental group of the 3-manifold, see [25].

While Perelman's approach is the most direct, there are other approaches to the Geometrization Conjecture using Ricci flow with surgery and variations of Theorem 5.5. As was indicated above, if a 3-manifold  $M$  admits an incompressible torus, then it falls into the class of 3-manifolds for which the Geometrization Conjecture had been established by Thurston himself. A detailed proof of the Geometrization Conjecture for those 3-manifolds was given in [24] and [25]. In view of this, it suffices to prove Theorem 5.5 for closed manifolds (again appealing to the Ricci flow results from [26] and the material in [27] preceding Theorem 7.4). This is route followed in [15] and [4]. A version of Theorem 5.5 for closed 3-manifolds has been proved in a series of papers of Shioya-Yamaguchi ([32], [33]). They did not make use of Assumption 3 of Theorem 5.5 on bounds on derivatives of curvature<sup>3</sup>, so their result is more general and can be applied to 3-manifolds that do not necessarily arise from Ricci flow. However, because they are not relying on estimates on higher derivatives

---

<sup>3</sup>Their proof was mostly for manifolds with curvature bounded from below, but the extension to the case of curvature locally bounded from below is not difficult as they point out in an appendix.

of the curvature as stated in Assumption 3, to prove their result, Shioya-Yamaguchi need to use a stability theorem on Alexandrov spaces. This stability theorem is due to Perelman and its proof was given in an unpublished manuscript in 1993. Recently, V. Kapovitch posted a preprint, [14], which proposes a more readable proof for this stability theorem of Perelman. Putting all these together, one has a Perelman-Shioya-Yamaguchi-Kapovitch proof of Theorem 5.5 for closed manifolds without the assumption of higher curvature bounds. As we have indicated, this proof requires a more knowledge about Alexandrov spaces, in particular knowledge about 3-dimensional Alexandrov spaces than the proof we present. It also relies on Thurston's result for manifolds with incompressible tori to give a complete proof of Geometrization.

Our presentation of the collapsing space theory is motivated by, and to a large extent follows, the Shioya-Yamaguchi paper [33], however it differs from their's in two fundamental aspects. First of all, as indicated above, we follow Perelman and add the assumption concerning the control of the higher derivatives of the curvature, thus allowing us to simplify the argument and in particular avoid the use of the stability theorem for Alexandrov spaces. Also, again following Perelman, we directly treat the case of non-empty boundary so that we do not have to appeal to Thurston's proof of the Geometrization Conjecture for manifolds containing an incompressible surface.

There is another approach to the proof of the Geometrization Conjecture due to Bessi eres et al [2] which avoids using Theorem 5.5 below. This argument also relies on Thurston's theorem that 3-manifolds with incompressible surfaces satisfy the Geometrization Conjecture, so that one only needs to consider the case when the entire closed 3-manifold is collapsed. Rather than appealing to the theory of Alexandrov spaces, this approach relies on other deep works in geometry and topology, e.g., results on the Gromov norms of 3-manifolds.

**PART I: Geometric and Analytic Results for Ricci Flow with Surgery**

## 1 Ricci flow with surgery

Let us review briefly the way we will apply Ricci flow with surgery in order to establish the Geometrization Conjecture. Here we are briefly reprising the work in [21]. Thurston's Geometrization Conjecture suggests the existence of especially nice metrics on 3-manifolds and consequently, a more analytic approach to the problem of classifying 3-manifolds. Richard Hamilton formalized one such approach in [11], the approach that Perelman successfully adopted, by introducing the Ricci flow on the space of Riemannian metrics on a fixed smooth manifold:

$$\frac{\partial g(t)}{\partial t} = -2\text{Ric}(g(t)), \quad (1.1)$$

where  $\text{Ric}(g(t))$  is the Ricci curvature of the metric  $g(t)$ . In dimension 3, the fixed points (up to rescaling) of this equation are the Riemannian metrics of constant sectional curvature. Beginning with any Riemannian manifold  $(M, g_0)$ , in [11] Hamilton showed that there is a solution  $g(t)$  of this Ricci flow on  $M$  for  $t$  in some interval such that  $g(0) = g_0$ . The naive hope is that if  $M$  is a closed 3-manifold, then  $g(t)$  exists for all  $t > 0$ , after appropriate rescaling, and converges to a nice metric outside a part with well-understood topology. As an example of this, in [12], R. Hamilton showed that if the Ricci flow exists for all time and if there is an appropriate curvature bound together with another geometric bound, then as  $t \rightarrow \infty$ , after rescaling to have a fixed diameter, the metric converges to a metric of constant negative curvature.

However, the general situation is much more complicated to formulate and much more difficult to establish. There are many technical issues that must be handled: One knows that in general the Ricci flow will develop singularities in finite time, and thus a method for analyzing these singularities and continuing the flow past them must be found. Furthermore, as we shall see, even if the flow continues for all time, there remain complicated issues about the nature of the metrics as  $t$  tends to  $\infty$ .

Let us discuss the finite-time singularities. If the topology of  $M$  is sufficiently complicated, say it is a non-trivial connected sum, then, no matter what the initial metric is, one must encounter finite-time singularities, forced by the topology. More seriously, even if  $M$  has simple topology, beginning with an arbitrary metric, one expects to (and cannot rule out the possibility that one will) encounter finite-time singularities in the Ricci flow. These singularities may occur along proper subsets of the manifold, not the entire manifold. Thus, one is led to study a more general evolution process called *Ricci flow with surgery*, denoted  $(\mathcal{M}, G)$ , first introduced by Hamilton in the context of four-manifolds, [13]. This evolution process is parametrized by an interval in time, and for each  $t$  in the interval of definition the  $t$  time-slice  $(M_t, g(t))$  is a compact Riemannian 3-manifold. But there is a discrete set of times at which the manifolds and metrics undergo topological and metric discontinuities (surgeries). In each of the complementary intervals to the singular times, the evolution is the usual Ricci flow, though, because of the surgeries, the topological type of the manifold  $M_t$  changes as  $t$  moves from one complementary interval to the next. From an analytic point of view, the surgeries at the discontinuity times are

introduced in order to ‘cut away’ a neighborhood of the singularities as they develop and insert by hand, in place of the ‘cut away’ regions, geometrically nice regions. This allows one to continue the Ricci flow (or more precisely, restart the Ricci flow with the new metric constructed at the discontinuity time). Of course, the surgery process also changes the topology. To be able to say anything useful topologically about such a process, one needs results about Ricci flow, and one also needs to control both the topology and the geometry of the surgery process at the singular times. For example, it is crucial for the topological applications that we do surgery along 2-spheres rather than surfaces of higher genus. Surgery along 2-spheres produces the connected sum decomposition, which, as we indicated above, is well-understood topologically, while, for example, (Dehn) surgeries along tori can completely destroy the topology, changing any 3-manifold into any other.

The change in topology turns out to be completely understandable and amazingly, the surgery processes produce exactly the topological operations needed to cut the manifold into pieces that are either prime or are copies on  $S^3$ , and furthermore, on each of these pieces the Ricci flow produces metrics sufficiently controlled so that the topology can be recognized, and the Geometrization Conjecture can be established.

### 1.1 Main Existence Theorem

Following Perelman ([27]), in [21] we gave a detailed proof of the long-time existence result for Ricci flow with surgery. First, an elementary definition.

**Definition 1.1.** We say that a Riemannian metric  $g$  on an  $n$  manifold  $M$  is *normalized* if for all  $x \in M$  we have  $|Rm(x)| \leq 1$  and  $\text{Vol}(B(x, 1)) \geq \omega_n/2$ , where  $\omega_n$  is the volume of the unit ball in Euclidean  $n$ -space. Clearly, if the Riemannian manifold  $(M, g)$  is compact, or more generally of bounded geometry, then there is a positive constant  $\lambda$  so that  $(M, \lambda g)$  is normalized.

**Theorem 1.2.** Fix  $\epsilon > 0$  sufficiently small. Let  $(M, g_0)$  be a closed Riemannian 3-manifold, with  $g_0$  normalized. Suppose that there is no embedded, locally separating  $\mathbb{R}P^2$  contained<sup>4</sup> in  $M$ . Then there is a Ricci flow with surgery, say  $(M, G)$ , defined for all  $t \in [0, \infty)$  with initial metric  $(M, g_0)$ . The set of discontinuity times for this Ricci flow with surgery is a discrete subset of  $[0, \infty)$ . The topological change in the time-slice  $M_t$  as  $t$  crosses a surgery time is a connected sum decomposition together with removal of connected components, each of which is diffeomorphic to one of  $S^2 \times S^1$ ,  $\mathbb{R}P^3 \# \mathbb{R}P^3$ , the non-orientable 2-sphere bundle over  $S^1$ , or a manifold admitting a metric of constant positive curvature. Furthermore, there are four non-increasing functions  $r(t) > 0$ ,  $\kappa(t) > 0$ ,  $\bar{\delta}(t) > 0$ , and  $h(t) > 0$  (independent of  $(M, g_0)$ ) such that: (1) surgery at time  $t$  is done with  $\bar{\delta}(t)$  control along 2-spheres with curvature  $\geq h^{-2}(t)$  (see the discussion immediately after Definition 15.5 in [21]); (2)  $(M_t, g(t))$  is  $\kappa(t)$ -non-collapsed (see [21] Definition 9.1); and (3) any point  $x \in M_t$  with  $R(g(t)) \geq r^{-2}(t)$  satisfies the so called strong  $(C, \epsilon)$ -canonical

<sup>4</sup>That is, no embedded  $\mathbb{R}P^2$  in  $M$  with trivial normal bundle. Clearly, all orientable manifolds satisfy this condition.

*neighborhood assumption for appropriate choices of  $C$  and  $\epsilon$  (see [21] Definition 9.78 and Theorem 15.9).*

Theorem 1.2 is central for all applications of Ricci flow to the topology of three-dimensional manifolds. The book [21] dealt with the case that  $M_t = \emptyset$  for  $t$  sufficiently large, that is, the case when the Ricci flow with surgery becomes extinct at finite time. Under this assumption, it follows from the above theorem that the initial manifold  $M$  is diffeomorphic to a connected sum of copies of  $S^2 \times S^1$ , the non-orientable 2-sphere bundle over  $S^1$ , and manifolds of the form  $S^3/\Gamma$ , where  $\Gamma \subset O(4)$  is a finite group acting freely on  $S^3$ . It was shown in [21] that if  $M$  is a simply-connected 3-manifold, then for any initial metric  $g_0$  the corresponding Ricci flow with surgery becomes extinct at finite time, see also ([28] and [5]). Consequently,  $M$  is diffeomorphic to  $S^3$ , thus proving the Poincaré Conjecture. More generally, in [21] we showed that if the fundamental group of  $M^3$  is a free product of finite groups and infinite cyclic groups, then  $M_t = \emptyset$  for all  $t$  sufficiently large. Hence, these manifolds are diffeomorphic to connected sums of prime manifolds admitting locally homogeneous metrics modelled either on the round metric on  $S^3$  (i.e., metrics of constant positive curvature) or on  $S^2 \times \mathbb{R}$  (the only prime examples of the latter being  $S^2$ -sphere bundles over  $S^1$ ).

In the case when  $M_t \neq \emptyset$  for every  $t$ , we showed (Corollary 15.4 of [21]) that if  $M_t$  satisfies the geometrization conjecture for some  $t > 0$  then so does the initial manifold  $M_0$ . Thus, it suffices to show that for any Ricci flow with surgery  $(\mathcal{M}, G)$ , for all  $t$  sufficiently large the  $t$  time-slice  $(M_t, g(t))$  satisfies the Geometrization Conjecture in order to conclude that it holds in general for all closed orientable 3-manifolds. This motivates a more detailed study of the time-slices  $(M_t, g(t))$  as  $t \rightarrow \infty$  for Ricci flows with surgery.

## 1.2 Review of notation and definitions

Here we recall the technical definitions for Ricci flows with surgery from [21] that will be used in the arguments we present here.

A *generalized Ricci flow of dimension  $n$*  is a smooth  $(n+1)$ -manifold  $U$  together with a time function  $t: U \rightarrow \mathbb{R}$  which is a submersion, a vector field  $\chi$ , and a smooth section  $g_{\text{hor}}$  of  $\text{Sym}^2((\text{Ker } dt)^*)$  subject to the following conditions:

1.  $\chi(t) = 1$ .
2.  $g_{\text{hor}}$  is a positive definite metric on  $\text{Ker } dt$ , and we denote by  $\text{Ric}(g_{\text{hor}})$  the symmetric 2-tensor on this bundle that is the Ricci curvature of the metric  $g_{\text{hor}}$ .
3. Denoting the Lie derivative with respect to  $\chi$  by  $\mathcal{L}_\chi$ , we have

$$\mathcal{L}_\chi(g_{\text{hor}}) = -2\text{Ric}(g_{\text{hor}}).$$

Said another way, for each point  $x \in U$ , setting  $t_0 = t(x)$ , there is a neighborhood  $V^n \subset t^{-1}(t_0)$  and a  $\xi > 0$  such that integrating flow lines of  $\chi$  through points of  $V$



determines a diffeomorphism from  $V \times (t_0 - \xi, t_0 + \xi)$  to a neighborhood of  $x$  in  $U$ . Furthermore, pulling back  $g_{\text{hor}}$  gives a smooth 1-parameter family of metrics  $g(t)$ ,  $t_0 - \xi < t < t_0 + \xi$ , on  $V$  that satisfies Equation (1.1), the Ricci flow equation. The special case of a generalized Ricci flow when the level sets of  $t$  are compact manifolds, or more generally when the flow lines of  $\chi$  determine a global product structure, is an ordinary Ricci flow. In a 3-dimensional Ricci flow with surgery  $(\mathcal{M}, G)$  the complement of the union of the surgery caps is a generalized Ricci flow, but of course it is not an ordinary Ricci flow since the topology of the time-slices changes.

Given a Ricci flow with surgery,  $(\mathcal{M}, G)$ , we denote by  $(M_t, g(t))$  the  $t$  time-slice. This is a compact Riemannian 3-manifold. For any  $(x, t) \in M_t$  and any  $r > 0$  we denote by  $B(x, t, r)$  the ball of radius  $r$  centered at  $x$  in  $(M_t, g(t))$ . Suppose that for some  $\Delta t > 0$  every  $y \in B(x, t, r)$  has the property that the flow-line of the Ricci flow with surgery through  $(y, t)$  extends backwards to at least  $t - \Delta t$ . Then we define the (backward) parabolic neighborhood  $P(x, t, r, -\Delta t)$  to be the union of these flow lines on the interval  $[t - \Delta t, t]$ . We then have an embedding  $B(x, t, r) \times [t - \Delta t, t] \subset \mathcal{M}$  and the pull-back of the Ricci flow with surgery by this embedding gives an ordinary Ricci flow on the product. In this case, we say *the Ricci flow with surgery contains the entire parabolic neighborhood  $P(x, t, r, -\Delta t)$*  or alternatively *the entire parabolic neighborhood exists in  $(\mathcal{M}, G)$* . There are analogous definitions and notation for forward parabolic neighborhoods  $P(x, t, r, \Delta t)$ .

We shall use other notation and definitions from [21]. Recall that, as we indicated above, Theorem 15.9 and Corollary 15.10 of [21], describing Ricci flows with surgery, make reference to two universal constants  $0 < \epsilon < 1/100$  and  $10 < C < \infty$  and four non-increasing, positive functions  $\kappa(t)$ ,  $r(t)$ ,  $\bar{\delta}(t)$  and  $h(t)$ . The function  $\kappa(t)$  is called the *non-collapsing function*: for every point  $(x, t)$  and radius  $r$  with  $0 < r \leq \epsilon$  with the property that the Ricci flow is defined on all of  $P(x, t, r, -r^2)$  and has all sectional curvatures on this set bounded in absolute value by  $r^{-2}$  also has the property that  $\text{Vol } B(x, t, r) \geq \kappa(t)r^3$ . This function is a step function on  $[0, \epsilon)$ ,  $[\epsilon, 2\epsilon)$ ,  $[2\epsilon, 4\epsilon)$ , etc. The function  $r(t)$  is the *canonical neighborhood function*. Every point  $(x, t) \in \mathcal{M}$  with  $R(x, t) \geq r^{-2}(t)$  has a  $(C, \epsilon)$ -canonical neighborhood (see below for the definition of the latter). It is also a step function on the same intervals as  $\kappa(t)$ . The function  $\bar{\delta}(t)$  is the *surgery control function*: Surgeries at time  $t$  along 2-spheres are performed along central 2-spheres of strong  $\bar{\delta}(t)$ -necks. The condition on  $\bar{\delta}(t)$  is that it be less than a universal non-increasing function  $\Delta(t)$  which is always less than  $\epsilon$  and limits to 0 as  $t \rightarrow \infty$ . The function  $\Delta(t)$  is also a step function on the same intervals as  $\kappa(t)$  and  $r(t)$ . The three step functions  $\kappa, r, \Delta$  are defined by interlocking induction one step at a time. Finally,  $h(t)$  is the *surgery scale function* in the sense that the 2-sphere surgeries at time  $t$  are done on the central 2-spheres of  $\bar{\delta}(t)$ -necks, 2-spheres through a point with scalar curvature  $h^{-2}(t)$ . The conditions on  $h(t)$  are two-fold. First, we require  $h(t) < \bar{\delta}^2(t)r(t)$ . Secondly,  $h(t)$  must be small enough so that any point  $(p, t)$  in an  $\epsilon$ -horn whose ‘big end’ has scalar curvature at least  $\bar{\delta}(t)r(t)$  and which satisfies  $R(p, t) \geq h^{-2}(t)$  is at the center of a strong  $\bar{\delta}(t)$ -neck. The function  $h(t)$  can be chosen arbitrarily subject to these two conditions after the other three functions have been defined for all  $t$ .

One of the conditions that our Ricci flows with surgery satisfy is the *curvature pinching hypothesis*. Setting  $X(x, t)$  equal to the maximum of the negative of the smallest eigenvalue of  $Rm(x, t)$  and zero, and assuming, as we always shall implicitly, that the initial conditions are normalized we have

$$R(x, t) \geq 2X(x, t) (\log(X(x, t)(t + 1)) - 3), \quad (1.2)$$

see Section 15 of [21].

In doing surgery at time  $t$  (see Section 14 of [21]) we remove connected components on which the scalar curvature is everywhere at least  $r^{-2}(t)$ . These are covered by  $(C, \epsilon)$ -canonical neighborhoods and thus by the results in the Appendix of [21] each such component either admits a round metric or admits a metric modelled on  $S^2 \times \mathbb{R}$  and hence these components satisfy the Geometrization Conjecture. We also cut open the  $\epsilon$ -horns of the limiting incomplete metric along the 2-spheres and remove the non-compact ends of these horns. We then add surgery caps at time  $t$ ; these are 3-disks added to the boundary 2-spheres created by the cutting process. The union of the surgery caps is exactly the set of points in  $M_t$  at which no flow line extends backwards, and hence exactly the set of points where the Ricci flow with surgery fails to satisfy the conditions to be a generalized Ricci flow. Every surgery cap at time  $t$  has diameter  $\leq 5h(t)$  and the scalar curvature on the surgery cap is bounded between  $3h^{-2}(t)/4$  and  $3h^{-2}(t)$ .

Recall that there are three types of  $(C, \epsilon)$ -canonical neighborhoods:

1. A strong  $\epsilon$ -neck centered at  $(x, t)$ . This is an evolving region in the Ricci flow with surgery on which the flow, after rescaling the metric and time by  $R(x, t)$  and shifting time so that the central point is at time 0, is within  $\epsilon$  in the  $C^{[1/\epsilon]}$ -topology of the standard product flow on  $S^2 \times (-\epsilon^{-1}, \epsilon^{-1}) \times [-1, 0]$  where the scalar curvature of the 2-spheres at time  $t$  is  $(1 - t)^{-1}$ .
2. A  $(C, \epsilon)$ -cap is an open submanifold  $\mathcal{C}$  of a time-slice, diffeomorphic to either an open 3-ball or the complement in  $\mathbb{R}P^3$  of a closed 3-ball, with a neighborhood  $N$  of the non-compact end of  $\mathcal{C}$  being the final time-slice of a strong  $\epsilon$ -neck. The complement  $\mathcal{C} \setminus N$  is called the *core* of the cap. Furthermore, the diameter of  $\mathcal{C}$  is at most  $CR(y)^{-1/2}$  for any  $y \in \mathcal{C}$ . There are also other bounds on curvature that are not relevant for us here.
3. The other type of  $(C, \epsilon)$ -canonical neighborhood consists of closed components of positive curvature. They will not play a role in this paper.

Furthermore, we require that in a  $(C, \epsilon)$ -canonical neighborhood we have

$$\left| \frac{dR(x, t)}{dt} \right| \leq CR^2(x, t) \quad (1.3)$$

$$|\nabla R(x, t)| \leq CR^{3/2}(x, t). \quad (1.4)$$

One of the main results of [21] is that in a Ricci flow with surgery is given  $\epsilon > 0$  there is a function  $r(t) > 0$  and  $C < \infty$  so that any point  $x$  in the  $t$ -time-slice  $M_t$  of

a Ricci flow with surgery with  $R(x) \geq r^{-2}(t)$  is either the center of a strong  $\epsilon$ -neck, is contained in the core of a  $(C, \epsilon)$ -cap or is contained in a  $(C, \epsilon)$ -component.

We shall use one further property of  $(C, \epsilon)$ -caps that was not required in [21] but which can be easily seen to be arrange from the construction: every point of  $N$  is itself at the center of an  $\epsilon$ -neck in  $M$ . To see how to arrange this, given  $\epsilon$  then fix  $C$  so that the result holds for  $(C, \epsilon/5)$ . For any  $(C, \epsilon/5)$ -cap with  $\epsilon/3$ -neck  $N$  as the complement of the core. Let  $N'$  be the middle  $1/5$  of  $N$ . Then  $N'$  is an  $\epsilon$ -neck and the union of the compact complementary component of  $N'$  with  $N'$  is a  $(C, \epsilon)$ -cap with the extra property that every  $x \in N'$  is the center of an  $\epsilon$ -neck in  $M$ . From now on we take this condition as part of the definition of a  $(C, \epsilon)$ -cap.

## 2 Limits as $t \rightarrow \infty$

We have finished our recap of the results, definitions, and notation from [21] that are necessary background. We now turn to the geometry of the of the volume non-collapsed part of the manifolds  $(M_t, g(t))$  as  $t \rightarrow \infty$ .

Recall (Equation 3.7 on page 41 of [21]) that for the 3-dimensional Ricci flow  $g(t)$ , one has the evolution equation on its scalar curvature  $R$

$$\frac{dR}{dt} = \Delta R + 2|\text{Ric}^0|^2 + \frac{2}{3}R^2, \quad (2.1)$$

where  $\text{Ric}^0$  is the trace-free part of  $\text{Ric}$ . Let  $R_{\min}(t)$  be the minimum of the scalar curvature  $R(g(t))$  of  $g(t)$ . Then by the usual (scalar) maximum principle we have

$$\frac{dR_{\min}}{dt} \geq \frac{2}{3}R_{\min}^2. \quad (2.2)$$

This inequality remains valid for Ricci flows with surgery, at least as long as  $R_{\min} < 0$ , since the surgery is done at a point with large positive scalar curvature. (In fact, the surgery is done at points where the scalar curvature is much larger than the threshold  $r^{-2}(t)$  for the existence of canonical neighborhood, so this equation remains true unless  $R_{\min}$  is greater than this threshold. If  $R_{\min}$  is greater than this threshold, then the manifold is covered by  $(C, \epsilon)$ -canonical neighborhoods and hence has a standard topology as described in the appendix of [21].) Because of the normalized initial conditions (see Assumption 1 in Chapter 15 of [21]),  $R_{\min}(0) \geq -6$ , it follows that

$$R_{\min}(t) \geq -\frac{3}{2(t + 1/4)}. \quad (2.3)$$

Furthermore, it follows from Equation (2.1) that if  $R_{\min}(0) > 0$  then the Ricci flow with surgery becomes extinct after a finite time, and according to the main theorem of [21], in this case the manifold is a connected sum of 3-dimensional spherical space-forms (quotients of  $S^3$  by finite groups of isometries acting freely) and copies of  $S^2$ -bundles over the circle. If  $R_{\min}(0) = 0$ , then by the strong maximum principle either it  $R_{\min}(t)$  is positive for all  $t > 0$  and the previous case applies, or  $R(x, 0) = 0$  for all  $x \in M_0$ . In the latter case, it follows from Equation (2.1) that either the

Ricci curvature and thus the sectional curvature vanishes identically and  $(M_0, g(0))$  is a flat manifold, or  $R_{\min}(t)$  is positive for all  $t > 0$  and the previous case applies. The conclusion is that if  $R_{\min}(0) \geq 0$ , then the initial manifold  $M_0$  satisfies the Geometrization Conjecture. From now on we assume that  $R_{\min}(t) < 0$  for all  $t$ , and hence that Inequality (2.3) holds for the Ricci flow with surgery. In fact as the Ricci flow with surgery proceeds and possibly breaks the manifold into several connected components, we remove from the Ricci flow any connected component of  $M_t$  on which the scalar curvature is everywhere non-negative. Hence, for all  $t$  and for every connected component  $C_t$  of  $M_t$  we have the  $\min_{x \in C_t} R(x, t) < 0$ .

**Definition 2.1.** Given a Ricci flow with surgery  $(\mathcal{M}, G)$  we set  $V(t)$  equal to the volume of  $(M_t, g(t))$  and we define  $\widehat{V}(t) = V(t)/(t + 1/4)^{3/2}$ . Define  $\widehat{R}(t) = R_{\min}(t)V^{2/3}(t)$ , where  $R_{\min}(t)$  is the minimum of the scalar curvature of  $(M_t, g(t))$ .

**Lemma 2.2.** *For any Ricci flow with surgery  $\widehat{V}(t)$  is a positive, non-increasing function of  $t$  and  $\widehat{R}(t)$  is a negative, non-decreasing function of  $t$*

*Proof.* Clearly,  $\widehat{V}(t) > 0$ . We have

$$\frac{d\widehat{V}(t)}{dt} = \frac{dV(t)/dt}{(t + 1/4)^{3/2}} - \frac{3}{2(t + 1/4)}\widehat{V}(t).$$

Since  $dV(t)/dt = -\int R(t)dV \leq -R_{\min}(t)V(t)$ , we have

$$\frac{d\widehat{V}(t)}{dt} \leq -\widehat{V}(t) \left( R_{\min} + \frac{3}{2(t + 1/4)} \right). \quad (2.4)$$

Inequality (2.3) implies that the right-hand side of the previous inequality is non-positive, so that  $\widehat{V}(t)$  is a non-increasing function of  $t$  in each interval between successive surgery times. Every surgery also reduces the volume in the sense that at every surgery time  $t_0$  we have  $\lim_{t \rightarrow t_0^-} V(t) \geq V(t_0)$ . The first statement follows.

Similarly,  $\widehat{R}(t) < 0$  since  $R_{\min} < 0$ . Using the inequality  $dR_{\min}(t)/dt \geq 2R_{\min}^2(t)/3$  and the equation  $dV(t)/dt = -\int R(t)dV$ , we have

$$\frac{d\widehat{R}(t)}{dt} \geq \frac{2}{3}\widehat{R}(t)V^{-1}(t) \int (R_{\min} - R)dV, \quad (2.5)$$

which is non-negative since  $\widehat{R}(t) < 0$ . As we have observed before since  $R_{\min}(t)$  is continuous at surgery times, it follows from the above that at any surgery time  $t_0$  we have

$$\lim_{t \rightarrow t_0^-} \widehat{R}(t) \geq \widehat{R}(t_0).$$

□

**Definition 2.3.** For a Ricci flow with surgery we define  $\widehat{V}(\infty) = \lim_{t \rightarrow \infty} \widehat{V}(t)$  and  $\widehat{R}(\infty) = \lim_{t \rightarrow \infty} \widehat{R}(t)$ .

**Lemma 2.4.** *Suppose that  $(\mathcal{M}, G)$  is a Ricci flow with surgery and that  $\widehat{V}(\infty) > 0$ . Then  $R_{\min}(t)$  is asymptotic to  $-3/2t$ , or equivalently  $\widehat{R}(\infty)\widehat{V}^{-2/3}(\infty) = -3/2$ .*

*Proof.* By Inequality (2.4) we have

$$\frac{d(\log \widehat{V})}{dt} \geq - \left( R_{\min} + \frac{3}{2(t+1/4)} \right).$$

Since we are assuming  $\widehat{V}(\infty) > 0$ , it follows that

$$\int_0^\infty R_{\min} + \frac{3}{2(t+1/4)} dt < \infty.$$

Now consider  $R_{\min}(t)(t+1/4) = \widehat{R}(t)/\widehat{V}(t)^{2/3}$ . Taking limits as  $t \rightarrow \infty$  gives

$$\lim_{t \rightarrow \infty} R_{\min}(t)(t+1/4) = \widehat{R}(\infty)/\widehat{V}(\infty)^{2/3}.$$

Thus,  $R_{\min}(t)(t+1/4)$  has a finite limit as  $t \rightarrow \infty$ . By the first inequality, that limit must be  $-3/2$ .  $\square$

The above results indicate that rescaling the metrics  $g(t)$  by  $1/t$  can lead to reasonable limits for  $R_{\min}(t)$ . As the next result shows that the same rescaling produces hyperbolic limits provided that we are working on regions on which these rescalings converge smoothly, see Section 7.1 of [27].

**Corollary 2.5.** *Let  $(\mathcal{M}, G)$  is a Ricci flow with surgery. Suppose that we have  $r > 0$ , a sequence  $t_n \rightarrow \infty$  as  $n \rightarrow \infty$ , and a sequence of parabolic neighborhoods  $P(x_n, t_n, r\sqrt{t_n}, -r^2 t_n)$  on which the Ricci flow with surgery is defined with the property that the rescaling of space and time in the  $n^{\text{th}}$ -parabolic neighborhood by  $t_n^{-1}$  converges smoothly as  $n \rightarrow \infty$  to a limit Ricci flow defined on an abstract parabolic neighborhood  $P(x_\infty, 1, r, -r^2)$ . Then, for every  $s \in (1-r^2, 1]$ , the sectional curvatures of the limit flow are constant on the  $s$  time-slice and equal to  $-1/4s$ .*

*Proof.* For each  $n \geq 1$  and each  $s \in [1-r^2, 1]$ , set  $g_n(s) = \frac{1}{t_n} g(st_n)$  on  $M_{st_n}$ . Denote by  $V_n(s)$ ,  $R_n(x, s)$  and  $\widehat{R}_n(s)$  the volume, the scalar curvature and the function  $\widehat{R}$  as defined above for  $(M_{st_n}, g_n(s))$ . By Inequality 2.5 we have

$$\widehat{R}(t_n) - \widehat{R}((1-r^2)t_n) \geq \int_{(1-r^2)t_n}^{t_n} \left[ \widehat{R}(s) V^{-1}(s) \int_{M_s} (R_{\min}(s) - R(x, s)) dV(s) \right] ds.$$

Changing variables, replacing  $s$  by  $t_n s$ , the right-hand side can be written as

$$\int_{1-r^2}^1 \left[ \widehat{R}(st_n) V^{-1}(st_n) \int_{M_{st_n}} (R_{\min}(st_n) - R(x, st_n)) dV(st_n) \right] t_n ds.$$

Since  $\widehat{R}$  is scale invariant, we can rewrite the right-hand side as

$$\begin{aligned} & \int_{1-r^2}^1 \left[ \widehat{R}_n(s) V_n^{-1}(s) t_n^{-3/2} \int_{M_{st_n}} t_n^{-1} (R_{n,\min}(s) - R_n(x, s)) t_n^{3/2} dV_n(s) \right] t_n ds \\ &= \int_{1-r^2}^1 \left[ \widehat{R}_n(s) V_n^{-1}(s) \int_{M_{st_n}} (R_{n,\min}(s) - R_n(x, s)) dV_n(s) \right] ds. \end{aligned}$$

Of course

$$V_n(s) = \widehat{V}(st_n) \cdot \left( \frac{t_n + 1/4}{t_n} \right)^{3/2},$$

and  $R_{n,\min}(s) - R_n(x, s) \leq 0$ . Thus, we have

$$\begin{aligned} \widehat{R}(t_n) - \widehat{R}((1-r^2)t_n) &\geq \\ \int_{1-r^2}^1 &\left[ \widehat{R}(st_n) \widehat{V}^{-1}(st_n) \left( \frac{t_n}{t_n + 1/4} \right)^{3/2} \int_{B(x,1,r)} (R_{n,\min}(s) - R_n(x, s)) dV_n(s) \right] ds. \end{aligned}$$

Taking limits as  $n \rightarrow \infty$  and denoting the limit metric on  $P(x, 1, r, -r^2)$  by  $g_\infty(s)$ , its scalar curvature by  $R_\infty(x, s)$  and its volume by  $V_\infty(s)$ , we have

$$0 \geq \widehat{R}(\infty) \widehat{V}^{-1}(\infty) \int_{1-r^2}^1 \int_{B(x,1,r)} (R_{\infty,\min}(s) - R_\infty(x, s)) dV_\infty(s) ds.$$

Since  $\widehat{R}(\infty) \widehat{V}^{-1}(\infty) < 0$  and the integrand is  $\leq 0$ , it follows that the integrand is identically zero, i.e.

$$R_{\infty,\min}(s) = R_\infty(x, s).$$

But we have already seen that  $R_{\infty,\min}(s) = -3/4s$ . This proves that  $g_\infty(s)$  has constant scalar curvature equal to  $-3/4s$ . It then follows from Equation 2.1 that  $\text{Ric}^0(g_\infty(s)) = 0$ , and hence  $g_\infty(s)$  is of constant sectional curvature  $-1/4s$  for every  $s \in [1-r^2, 1]$ .  $\square$

This analysis gives us control on the nature of the  $(M_t, \frac{1}{t}g(t))$  as  $t \rightarrow \infty$ , at sequences points whose times are going to infinity and for which there is a smooth limit on a parabolic neighborhood, and in fact is the source of the hyperbolic limits at infinity. But in order to apply this corollary we need to understand when these rescalings have limits. For this we need three local results that are more delicate. In the next section we will establish technical results that are used in proving the following three propositions. In all three propositions we are considering sequences of time-slices where  $t \rightarrow \infty$  and implicitly we are rescaling the metric by  $1/t$ . It turns out that these technical results require further conditions, another upper bound, on the surgery control parameter  $\bar{\delta}(t)$  (see Assumption 3.9) and on the surgery scale function  $h(t)$  (see Assumption 3.10) beyond those stated in Corollary 15.10 in [21]. Since Corollary 15.10 of [21] is valid as long as  $\bar{\delta}(t)$  was non-increasing and less than some fixed function  $\Delta(t) > 0$ , we can simply take  $\bar{\delta}(t)$  less than  $\Delta(t)$  and also less the new upper bound required here. Similarly, since the choices of the  $\kappa(t), r(t)$ , and  $\Delta(t)$  are independent of the choice of  $h(t)$  satisfying the given conditions, we are also free to add this extra condition as an upper bound for  $h(t)$ . **Throughout this section we assume that the surgery control parameter  $\bar{\delta}(t)$  and the surgery scale function  $h(t)$  satisfy the conditions given in Assumptions 3.9 and 3.10.**

## 2.1 Three propositions

Next, we state three geometric and analytic propositions that allow us to control the nature of the Ricci flow with surgery at large times. The first proposition shows that one can take limits and hence the curvature is close to  $-1/4t$  on regions that are volume non-collapsed with lower curvature bounds. Furthermore, there is a stability in that the limits exist forward for a certain amount of time. This is Lemma 7.2 of [27].

**Proposition 2.6.** (a) *Given  $w > 0, r > 0, \xi > 0$  there is  $T = T(w, r, \xi) < \infty$  such that the following holds for any Ricci flow with surgery  $(\mathcal{M}, G)$  satisfying Assumptions 3.9 and 3.10. If, for some  $t_0 \geq T$  and some  $x_0 \in M_{t_0}$ , the ball  $B(x_0, t_0, r\sqrt{t_0})$  has volume at least  $w r^3 t_0^{3/2}$  and sectional curvatures bounded below by  $-r^{-2} t_0^{-1}$ , then*

$$|2t_0 \text{Ric}(x_0, t_0) + g(x_0, t_0)|_{g(t_0)} < \xi. \quad (2.6)$$

(b) *In addition, given  $A < \infty$ , there is  $T' = T'(w, r, \xi, A) \geq T(w, r, \xi)$ , and provided that  $t_0 \geq T'$ , the Ricci flow with surgery contains the entire forward parabolic neighborhood  $P(x_0, t_0, Ar\sqrt{t_0}, Ar^2 t_0)$  and Equation (2.6) holds with  $(x_0, t_0)$  replaced by any  $(x, t)$  in this forward parabolic neighborhood.*

Before stating the second proposition we need a definition.

**Definition 2.7.** Given a Ricci flow with surgery  $(\mathcal{M}, G)$ , we define a function from  $\rho: \mathcal{M} \rightarrow (0, \infty)$  by setting  $\rho(x, t)$  equal to the largest real number with the property that  $Rm|_{B(x, t, \rho(x, t))} \geq -\rho^{-2}(x, t)$ .

The fact that no component of  $(M_t, g(t))$  has non-negative curvature implies that the function  $\rho$  exists and takes finite values.

The second proposition is a volume collapsing result at points where  $\rho$  is sufficiently small (see Section 7.3 of [27]).

**Proposition 2.8.** *For any  $w > 0$  there is  $\bar{\rho} = \bar{\rho}(w) > 0$  such that for all  $t$  sufficiently large (how large depending on  $w$ ) for any Ricci flow with surgery  $(\mathcal{M}, G)$  satisfying Assumptions 3.9 and 3.10, and for any  $x \in M_t$ , if  $\rho(x, t) < \bar{\rho}\sqrt{t}$  we have*

$$\text{Vol } B(x, t, \rho(x, t)) < w \rho^3(x, t).$$

The third result shows that, under the hypotheses of volume non-collapsing with a lower curvature bound, we have bounds on the norm of the Riemannian curvature and all its covariant derivatives. (This is the last hypothesis in Theorem 7.4 of [27].)

**Proposition 2.9.** *For every  $w' > 0$  there exist  $\bar{r} = \bar{r}(w') > 0$  and constants  $K_m = K_m(w') < \infty$ ,  $m = 0, 1, \dots$ , such that the following holds for any Ricci flow with surgery  $(\mathcal{M}, G)$  satisfying Assumptions 3.9 and 3.10 and for all  $t$  sufficiently large, how large depending only on  $w'$ . For any  $0 < r \leq \bar{r}\sqrt{t}$ , for any  $x \in M_t$ , and for any  $m > 0$ . Suppose that the ball  $B(x, t, r)$  has volume at least  $w' r^3$  and sectional curvatures bounded below by  $-r^{-2}$ . Then the norms of the curvature and its  $m^{\text{th}}$ -order covariant derivatives at  $(x, t)$  are bounded by  $K_0 r^{-2}$  and  $K_m r^{-(2+m)}$ , respectively.*

For the rest of this section we assume these three results; they will be proved in the next section.

## 2.2 The hyperbolic pieces

Let us begin with some basic definitions from 3-dimensional hyperbolic geometry. Because of the curvature of the limits arising in Corollary 2.5 we take the following slightly non-standard definition of a hyperbolic manifold. For us a *hyperbolic metric* on a 3-manifold is a Riemannian metric of constant sectional curvature  $-1/4$ . By a *hyperbolic manifold* we mean a Riemannian 3-manifold with a hyperbolic metric.

### 2.2.1 3-dimensional hyperbolic manifolds of finite volume

**Definition 2.10.** Let  $H$  be a non-compact, complete, orientable hyperbolic 3-manifold of finite volume. Then the fundamental group of each end of  $H$  is a free abelian group of rank 2 acting on the  $S^2$  at infinity of hyperbolic 3-space by parabolic elements (i.e., elements with a single fixed point on the 2-sphere). Choosing upper half space coordinates on hyperbolic three space,  $\mathbb{C} \times (0, \infty)$ , so that the fixed point of these commuting elements is  $\infty$ , the group they generate leaves invariant each plane  $\mathbb{C} \times \{t\}$ , called the *horospheres at infinity* and the quotient of each of these planes by the resulting action of  $\mathbb{Z} \times \mathbb{Z}$  is a torus, called the *horospherical tori* of the end. The induced metric on the horospherical tori changes by a conformal factor  $t/t'$  as we move from the one at height  $t$  to the one at height  $t'$ . (The distance between these plans is  $\ln(t'/t)$ .) For all  $t$  sufficiently large, the horospherical torus at height  $t$  embeds into  $H$ . The region of this end cut off by such an embedded horospherical torus is called a *cuspidal end*. Each cusp is foliated by horospherical tori, and every end of  $H$  is a cusp. A *truncation*  $\bar{H}$  of  $H$  is a compact submanifold whose boundary is a disjoint union of horospherical tori and whose complement,  $H \setminus \bar{H}$ , is foliated by horospherical tori and hence contained in the union of the cusps. The complement is diffeomorphic to  $\partial H \times [0, \infty)$ .

According to a result of Margulis's there is a constant  $w_0 > 0$  such that the following holds for any  $0 < w \leq w_0$  and any complete hyperbolic 3-manifold,  $H$ , of finite volume.

1. For each end  $\mathcal{E}$  of  $H$  let  $\tilde{H}_{\mathcal{E}}$  be covering space corresponding to the fundamental group of the end. Let  $\tilde{T}_w$  denote the horospherical torus in  $\tilde{H}_{\mathcal{E}}$  with the property that for each  $\tilde{x} \in \tilde{T}_w$  we have  $\text{Vol}_{H_{\mathcal{E}}} B(\tilde{x}, 2) = 8w$ . Denote by  $\tilde{U}(w) \subset \tilde{H}_{\mathcal{E}}$  the open set of all points within distance 2 of  $\tilde{T}_w$ . The projection  $\tilde{H}_{\mathcal{E}} \rightarrow H$  embeds  $\tilde{U}(w)$  into  $H$ . The image,  $U(w)$ , is the neighborhood of size 2 about the horospherical torus  $T_w \subset H$  that is the image of  $\tilde{T}_w$ , and  $\text{Vol}_H B(x, 2) = 8w$  for all  $x \in T_w$ .
2. The open subset of  $H$  that is  $w$ -volume collapsed on scale 2 consists of the cuspidal ends cut off by the  $T_w$  and a finite number of solid torus neighborhoods of short geodesics.



For any  $0 < w \leq w_0$  and for any complete hyperbolic 3-manifold of finite volume, we define the  $w$ -truncation of  $H$ , denoted  $\overline{H}(w)$ , by taking as boundary the horospherical torus  $T_w$  of each end of  $H$ .

Notice that it is not necessarily true that for every point  $x \in \overline{H}(w)$  the ball  $B(x, 2)$  has volume at least  $8w$ . The reason is that  $H$  can have short geodesics and around each there is a solid torus of points that are  $w$ -volume collapsed on the scale 2. Any given hyperbolic manifold has only finitely many such short geodesics. Thus, given a complete hyperbolic manifold  $H$  of finite volume there is a positive constant  $w' = w'(H) \leq w_0$  such that no point of  $\overline{H}(w')$  is  $w'$ -volume collapsed on the scale 2.

The next result is a consequence of Mostow rigidity [22] for hyperbolic 3-manifolds as well as Margulis's description of the sufficiently volume collapsed regions of a hyperbolic manifold.

**Lemma 2.11.** *There are constants  $\nu_0 > 0$  and  $0 < w_1 \leq w_0/2$  such that the following holds. Suppose that  $H$  and  $H'$  are complete hyperbolic 3-manifolds of finite volume with  $g'$  being the hyperbolic metric on  $H'$ . Further, suppose that  $\varphi: \overline{H}(w_1) \rightarrow H'$  is a smooth embedding with  $\varphi^*g'$  within distance  $\nu_0$  in the  $C^\infty$ -topology to the restriction to  $\overline{H}(w_1)$  of the metric on  $H$ . Then  $H' \setminus \varphi(\text{int } \overline{H}(w_1))$  is contained in the part of  $H'$  that is  $2w_1$  volume collapsed on the scale of its negative curvature. Furthermore, this difference is a disjoint union of solid torus neighborhoods of short geodesics and components diffeomorphic to  $T^2 \times [0, \infty)$  and contained in the cusps of  $H$ . In particular, if  $H'$  has at least as many cusps as  $H$ , then  $H$  and  $H'$  are isometric.*

*Proof.* Fix  $0 < w_1 < w_0/2$ . It follows easily that the given embedding  $\varphi: \overline{H}(w_1) \rightarrow H'$  has image whose boundary is contained in the cusps of  $H'$  and solid tori about short geodesics of  $H'$ . Furthermore, each boundary torus is parallel in  $H'$  to either a horospherical torus in the cusp that contains it or to the boundary of the solid torus neighborhood of a short geodesic that contains it. Thus, topologically  $H'$  is obtained from  $H$  by Dehn filling some of its boundary components (i.e., by truncating some of the cusps and gluing solid tori to the resulting boundaries). Clearly, then  $H'$  has at most as many cusps as  $H$ , and if it has as many, then none of the boundary tori of  $H$  are filled in creating  $H'$ . In this case  $\varphi$  is a homotopy equivalence between  $\overline{H}(w_1)$  and  $H'$ . By Mostow rigidity [22], it follows that in this latter case  $H$  and  $H'$  are isometric.  $\square$

### 2.2.2 Hyperbolic limits at infinity

For this subsection we fix a Ricci flow with surgery  $(\mathcal{M}, G)$ . Now we shall use the three propositions stated in Section 2.1 to establish that the limits required by Corollary 2.5 exist and consequently that there exist complete, finite volume hyperbolic limits for the non-collapsing part of the  $(M_t, g(t))$  as  $t \rightarrow \infty$ . All estimates on how large  $t$  has to be for various conclusions to hold depend on the Ricci flow with surgery.

**Definition 2.12.** A *geometric limit at infinity* of a Ricci flow with surgery,  $(\mathcal{M}, G)$ , is a based complete Riemannian manifold  $(H, x_\infty)$  for which there is a sequence

$(x_n, t_n) \in \mathcal{M}$  with  $t_n \rightarrow \infty$  such that, setting  $X_n = (M_{t_n}, (1/t_n)g(t_n))$ , the sequence of based Riemannian manifolds  $(X_n, x_n)$  converges geometrically to  $(H, x_\infty)$ . This means that for every  $0 < R < \infty$ , for all  $n$  sufficiently large, there are embeddings  $f_{n,R}: B(x_\infty, R) \rightarrow X_n$ , sending  $x_\infty$  to  $x_n$  so that the Riemannian metrics  $f_{n,R}^*(1/t_n)g(t_n)$  converge smoothly to the restriction to  $B(x_\infty, R)$  of the hyperbolic metric on  $H$ .

Given a Ricci flow with surgery  $(\mathcal{M}, G)$  satisfying Assumptions 3.9 and 3.10, for any  $w > 0$  and any  $t < \infty$  we define

$$\widetilde{M}_t(w, -) = \{x \in M_t \mid \text{Vol } B(x, t, \rho(x, t)) < w\rho^3(x, t)\}.$$

We define

$$M_t(w, +) = M_t \setminus \widetilde{M}_t(w, -),$$

and we set  $\bar{\rho} = \bar{\rho}(w)$  from Proposition 2.8.

According to Proposition 2.8, for all  $t$  sufficiently large, we have  $\rho(x, t) \geq \bar{\rho}\sqrt{t}$  for all  $x \in M_t(w, +)$ . It then follows from Proposition 2.6 that, given any  $A < \infty$  and any  $\xi > 0$  sufficiently small, for  $t$  sufficiently large (given  $w, \xi$ , and  $A$ ) for every point  $x \in M_t(w, +)$  Equation (2.6) holds at every point of  $P(x, t, A\bar{\rho}\sqrt{t}, A\bar{\rho}^2t)$ . After rescaling the metrics and time by  $t^{-1}$ , this gives us a Ricci flow on a parabolic neighborhood  $P = P(x, 1, A\bar{\rho}, A\bar{\rho}^2)$  of (incomplete)  $\xi$ -almost hyperbolic manifolds, in the sense that Inequality (2.6) holds at all points of  $P$ . Now suppose that  $M_{t_n}(w, +) \neq \emptyset$  for a sequence  $t_n$  going to  $\infty$ , and for each  $n$  choose a point  $x_n \in M_{t_n}(w, +)$ . Consider the based Ricci flows  $(M_{t_n}, \frac{1}{t_n}g(t_n), (x_n, 1))$ . It follows from the above discussion that given any  $A$ , for all  $n$  sufficiently large, all sectional curvatures of the metrics  $(1/t_n)g(t_n)$  on  $B_{(1/t_n)g(t_n)}(x_n, A)$  for  $1 \leq t \leq A^2$  are close to constant  $-1/4t$ . Furthermore, the volume of  $B_{(1/t_n)g(t_n)}(x_n, 2)$  is bounded away from zero as  $n \rightarrow \infty$ . Thus, by Proposition 2.9, all the higher derivatives of the metrics in this sequence are controlled in these parabolic neighborhoods. This means that these parabolic neighborhoods converge smoothly to a flow on a parabolic neighborhood  $P(x_\infty, 1, A, A^2)$  which is a flow of incomplete hyperbolic manifolds with the curvature at time  $t$  being  $-1/4t$ . This is true for every  $A < \infty$ , and after passing to a subsequence, these limit flows can be embedded one in the next to produce a limiting Ricci flow of complete manifolds

$$(H, g_{\text{hyp}}(t), (x_\infty, 1)), 1 \leq t < \infty,$$

where  $(H, g_{\text{hyp}}(t))$  has constant sectional curvature  $-1/4t$ . The volume of  $(H, g_{\text{hyp}}(1))$  is at most  $\lim_{t \rightarrow \infty} V(t)/t^{3/2} = \widehat{V}(\infty)$  and hence is finite. The fact that the Ricci flows on the parabolic neighborhoods converge smoothly to the restriction of the flow of complete hyperbolic metrics implies that the  $(M_{t_n}, (1/t_n)g(t_n), x_n)$  converge geometrically to  $H$  and the that generalized Ricci flows starting with these manifolds (and rescaled by  $t_n^{-1}$ ) converge geometrically to the Ricci flow of complete hyperbolic manifolds. This establishes the following limiting result.

**Proposition 2.13.** *For any  $w > 0$ , for any Ricci flow with surgery  $(\mathcal{M}, G)$  satisfying Assumptions 3.9 and 3.10, for any sequence of  $t_n \rightarrow \infty$ , and for any sequence*

$x_n \in M_{t_n}(w, +)$ , after passing to a subsequence, the  $(M_{t_n}, (1/t_n)g(t_n), x_n)$  converge geometrically to a complete hyperbolic manifold of finite volume. Furthermore, for any sequence  $t'_n \rightarrow \infty$  and for any sequence  $y_n \in M_{t'_n}$ , with the property that the sequence of based Riemannian manifolds  $(M_{t'_n}, (1/t'_n)g(t'_n), y_n)$  has a geometric limit, that limit is a complete hyperbolic manifold of finite volume.

*Proof.* The first statement was established in the previous discussion. For the second, if there is a geometric limit of associated to the sequence  $(y_n, t'_n) \in \mathcal{M}$ , then for some  $0 < w \leq w_0$  we have  $y_n \in M_{t'_n}(w, +)$  for all  $n$  sufficiently large. Hence, by the first statement after passing to a subsequence of the  $y_n$  there is a geometric limit of the  $(M_{t'_n}, (1/t'_n)g(t'_n), y_n)$ , a limit that is a complete hyperbolic manifold of finite volume. The geometric limit of the entire sequence agrees with the geometric limit of this subsequence.  $\square$

**Definition 2.14.** Fix  $0 < w \leq w_0$  and  $\nu > 0$ . Let  $(\mathcal{M}, G)$  be a Ricci flow with surgery. A  $w$ -truncated,  $\nu$ -almost hyperbolic manifold at time  $t$  in  $(\mathcal{M}, G)$  is a complete hyperbolic 3-manifold  $H$  and an embedding  $\varphi: \overline{H}(w) \rightarrow M_t$  with the property that  $(1/t)\varphi^*g(t)$  is within  $\nu$  in the  $C^\infty$ -topology of the restriction of the hyperbolic metric of  $H$  to  $\overline{H}(w)$ .

The following strengthening of Proposition 2.13 follows from the discussion immediately preceding that proposition:

**Corollary 2.15.** Fix a Ricci flow with surgery  $(\mathcal{M}, G)$  satisfying Assumptions 3.9 and 3.10. Given  $0 < \nu \leq \nu_0$  and  $0 < w \leq w_0$  there is  $T(w, \nu) < \infty$  such that the following holds and for all  $t \geq T(w, \nu)$ . For any  $x \in M_t(w, +)$  there is a  $w$ -truncated  $\nu$ -almost hyperbolic manifold in  $(\mathcal{M}, G)$  at time  $t$ ,  $\varphi: \overline{H}(w) \subset M_t$ , containing  $x$  and with the property that every flow line in  $\mathcal{M}$  beginning at a point of  $\varphi(\overline{H}(w))$  exists for all  $t' \in [t, 2t]$ . In particular, flowing along these flow lines determines an embedding of  $\widehat{\varphi}: \overline{H}(w) \times [t, 2t] \hookrightarrow \mathcal{M}$ . For every  $t \leq t' \leq 2t$  the metric  $(1/t')\varphi^*g(t')$  is within  $\nu$  in the  $C^\infty$ -topology of the restriction of the hyperbolic metric on  $H$ .

**Definition 2.16.** A  $w$ -truncated,  $\nu$ -almost hyperbolic manifold at time  $t$  in  $(\mathcal{M}, G)$  that satisfies the conclusion of the previous corollary is said to last until time  $2t$ . In this case the embedding  $\widehat{\varphi}: \overline{H}(w) \times [t_0, 2t_0] \rightarrow \mathcal{M}$  has the property that for every  $t_0 \leq t' \leq 2t$ , the restriction of  $\widehat{\varphi}$  to  $\overline{H}(w) \times \{t'\}$  embeds  $\overline{H}(w)$  as a  $w$ -truncated  $\nu$ -almost hyperbolic manifold at time  $t'$ .

By a  $w$ -truncated  $\nu$ -almost hyperbolic tower starting at time  $t$  for  $(\mathcal{M}, G)$  we mean a complete hyperbolic manifold  $H$  and a sequence of embeddings of  $\varphi_k: \overline{H}(w) \rightarrow M_{2^k t}$ ,  $k = 0, 1, \dots$ , such that for each  $k \geq 0$  the image of  $\varphi_k$  is a  $w$ -truncated  $\nu$ -almost hyperbolic manifold at time  $2^k t$  that lasts until time  $2^{k+1}t$ . Furthermore, the image of flowing  $\varphi_k(\overline{H}(w))$  from time  $2^k t$  to time  $2^{k+1}t$ , which is  $\widehat{\varphi}_k(\overline{H}(w) \times \{2^{k+1}t\})$ , contains  $\varphi_{k+1}(\overline{H}(2w))$ . The tower is said to be constructed from the hyperbolic manifold  $H$ . We denote by  $\mathcal{T} \subset \mathcal{M}$  the union of the images  $\widehat{\varphi}_k(\overline{H}(w) \times [2^k t, 2^{k+1}t])$ , and by abuse of terminology we call this subset a  $w$ -truncated,  $\nu$ -almost hyperbolic tower. The  $k^{\text{th}}$  stage of the tower is the image of  $\widehat{\varphi}_{k-1}$ .

**Lemma 2.17.** *For any  $0 < w \leq w_1$ , for  $\nu > 0$  sufficiently small, for any complete hyperbolic manifold  $H$  of finite volume, and for any  $w$ -truncated,  $\nu$ -almost hyperbolic tower  $\mathcal{T} \subset \mathcal{M}$  constructed from  $H$ , any sequence  $(x_n, t_n) \in \mathcal{T}$  with  $t_n \rightarrow \infty$  has a subsequence converging geometrically to  $H$ .*

*Proof.* We suppose that  $0 < \nu < \nu_0/3$ . There is a  $w' = w'(H)$  such that every point of  $\overline{H}(w)$  is  $w'$ -volume non-collapsed on scale 2. It follows that, provided that  $\nu$  is sufficiently small, every point of  $\mathcal{T}$  is  $w'/2$ -volume non-collapsed on scale 2. Thus, by Proposition 2.13 given any sequence  $(x_n, t_n) \in \mathcal{T}$  with  $t_n \rightarrow \infty$ , after passing to a subsequence, there is a geometric limit which is a complete hyperbolic manifold  $H_\infty$  of finite volume. By the definition of geometric limits, for all  $n$  sufficiently large there is an embedding  $\psi: \overline{H}(w) \subset H_\infty$  with  $\psi^*$  of the hyperbolic metric on  $H_\infty$  within  $\nu$  of the restriction to  $\overline{H}(w)$  of the metric  $(1/t_n)(g(t_n))$ . On the other hand, by the definition of a hyperbolic tower, the restriction of the metric  $(1/t_n)g(t_n)$  to  $\overline{H}(w)$  is within  $\nu$  of the restriction of the hyperbolic metric of  $H$  to  $\overline{H}(w)$ . It follows that  $\psi^*$  of the hyperbolic metric on  $H_\infty$  is within  $\nu_0$  of the restriction to  $\overline{H}(w)$  of the hyperbolic metric on  $H$ . Thus, by Lemma 2.11 either  $H_\infty$  and  $H$  are isometric or  $H_\infty$  has fewer cusps than  $H$ . But  $H$  was chosen to have the minimal number of cusps of all geometric limits at infinity of  $(\mathcal{M}, G)$ . Consequently,  $H_\infty = H$ .  $\square$

**Definition 2.18.** We say that the tower  $\mathcal{T}$  given in Lemma 2.17 *converges* to  $H$ .

Suppose that we have any sequence  $(x_n, t_n) \in \mathcal{M}$  with  $t_n \rightarrow \infty$  with a geometric limit  $H_\infty$ . Then there is  $z \in H_\infty$  that is  $w_0$ -non-collapsed on scale 2. Hence, for a sequence  $(x'_n, t_n)$  converging to  $z$ , and for all  $n$  sufficiently large, we have  $(M_{t_n}, (1/t_n)g(t_n))$  is  $w_0/2$  volume non-collapsed at  $x_n$ . This means that for all  $n$  sufficiently large  $(x_n, t_n) \in M_{t_n}(w_0/2, +)$ . This shows that for any  $0 < w \leq w_0/2$  any geometric limit at infinity of  $(\mathcal{M}, G)$  is in fact the limit of a sequence  $(x_n, t_n)$  with  $x_n \in M_{t_n}(w, +)$ . In particular,  $(\mathcal{M}, G)$  has geometric limits at infinity if and only if  $M_{t_n}(w, +) \neq \emptyset$  for a sequence of  $t_n$  tending to  $\infty$ . It also follows from this that there is a sequence  $x_n \in M_{t_n}(w, +)$  with  $t_n \rightarrow \infty$  such that the sequence  $(M_{t_n}, (1/t_n)g(t_n), x_n)$  has a geometric limit  $H$  with the property that  $H$  has the minimal number of cusps among limits among all geometric limits at infinity of  $(\mathcal{M}, G)$ .

**Proposition 2.19.** *Fix a geometric limit at infinity  $H$  for  $(\mathcal{M}, G)$  with a minimal number of cusps among all such geometric limits. For any  $w > 0$  and  $\nu > 0$  sufficiently small, there is a  $w$ -truncated,  $\nu$ -almost hyperbolic tower  $\mathcal{T}$  converging to  $H$ .*

*Proof.* We take  $w \leq \min(w_0/2, w'(H))$  and  $\nu > 0$  small. Fix a sequence  $(x_n, t_n)$  with  $x_n \in M_{t_n}(w, +)$  with geometric limit  $H$ . By the definition of the limiting process, after passing to a subsequence, for all  $n$  there is an embedding  $\varphi_n: \overline{H}(w/2) \subset M_{t_n}$  containing the component of  $M_{t_n}(w, +)$  containing  $x_n$  such that  $\frac{1}{t_n}\varphi_n^*g(t_n)$ ,  $1 \leq t \leq 2$ , converges as  $n \rightarrow \infty$  to the restriction of the hyperbolic flow  $(H, g_{\text{hyp}}(t))$ ,  $1 \leq t < \infty$  to  $\overline{H}(w/2) \times [1, 2]$ . For all  $n$  sufficiently large this constructs a  $w/2$ -truncated,

$\nu$ -almost hyperbolic manifold  $\varphi_n: \overline{H}(w/2) \rightarrow M_{t_n}$  at time  $t_n$  that lasts to time  $2t_n$ . This is the first stage of the tower.

Now we fix  $0 < \nu < \nu_0/3$ .

**Claim 2.20.** *There is  $T_1 \geq T'(w/4, \nu)$  such that the following hold for any  $t \geq T_1$ . Suppose that  $\varphi: \overline{H}'(w/2) \rightarrow M_t$  is a  $w/2$ -truncated  $\nu$ -almost hyperbolic manifold at time  $t$ . Then  $H'$  has at least as many cusps as  $H$ .*

*Proof.* Suppose there is a sequence  $t_n \rightarrow \infty$  and  $w/2$ -truncated  $\nu$ -almost hyperbolic manifolds  $\overline{H}_n(w/2) \subset M_{t_n}$  at time  $t_n$  with each  $H_n$  having few cusps than  $H$ . Fix points  $(x_n, t_n) \in \overline{H}'_n(w/2) \subset M_{t_n}$ . Then according to Corollary 2.15 passing to a subsequence we can extract a limit of the  $(M_{t_n}, (1/t_n)g(t_n), x_n)$  and this limit is a complete hyperbolic manifold  $H_\infty$  of finite volume. By the definition of the limit, for all  $n$  sufficiently large we have an embedding  $\psi_n: \overline{H}'_n(w/2) \rightarrow H_\infty$  so that the pull-back of the hyperbolic metric on  $H_\infty$  is within  $\nu$  of the restriction of the hyperbolic metric on  $H'_n$ . It follows from Lemma 2.11 that  $H_\infty$  has at most as many cusps as  $H'_n$  for all sufficiently large  $n$  and hence it has fewer cusps than  $H$ . This contradicts the choice of  $H$  as having the fewest number of cusps among all geometric limits that are hyperbolic.  $\square$

Now we fix  $t$  equal to one of the  $t_n$  in the above subsequence with  $t_n > T_1$ . (Recall that  $T_1 \geq T'(w/4, \nu)$ .) We relabel the map  $\varphi_n$  above and call it  $\varphi_0$ . It is a map  $\varphi_0: \overline{H}(w/2) \rightarrow M_t$  giving a  $w/2$ -truncated,  $\nu$ -almost hyperbolic manifold at time  $t$  that lasts to time  $2t$ . The image  $\varphi_0(2t)(\overline{H}(w/2))$  is contained in  $M_{2t}(w/4, +)$  and contains a component  $V$  of  $M_{2t}(3w/4, +)$ . Invoking Proposition 2.6 again, we see that since  $t > T'(w/4, \nu)$ , there is a complete hyperbolic manifold  $H'$  of finite volume and an embedding  $\psi: \overline{H}'(w/2) \rightarrow M_{2t}$  containing  $V$  giving a  $w/2$ -truncated,  $\nu$ -almost hyperbolic manifold at time  $2t$  that lasts to time  $4t$ . We claim that  $H' = H$ . Denote by  $g'$  the hyperbolic metric on  $H'$ . Since  $\lambda = \psi^{-1} \circ \varphi_0(2t): \overline{H}(w) \rightarrow \overline{H}'(w/2)$  has the property that  $\lambda^*g'_{\text{hyp}}$  is within  $\nu_0$  in the  $C^\infty$ -topology of the restriction of  $g_{\text{hyp}}$  to  $\overline{H}(w)$ . Also, since  $t \geq T_1$  it follows from the previous claim that  $H'$  has at least as many cusps as  $H$ . Hence, by Lemma 2.11 we see that  $H'$  is isometric to  $H$ . This constructs the map  $\varphi_1: \overline{H}(w/2) \rightarrow M_{2t}$  which is a  $w/2$ -truncated,  $\nu$ -almost hyperbolic manifold whose image at time  $2t$  contains  $\widehat{\varphi}_0(\overline{H}(w) \times \{2t\})$ . This is as required for the second stage of the tower.

We simply repeat this construction *ad infinitum* to complete the proof of the proposition.  $\square$

**Addendum 2.21.** We could produce a more refined version of a hyperbolic tower  $\mathcal{T}$  as follows: given  $w_n \rightarrow \infty$  and  $\nu_n \rightarrow \infty$ , then there is a monotone increasing function  $k(n)$  such that for every  $n$ , the  $(k(n) + 1)^{\text{st}}$  stage of the tower  $\widehat{\varphi}_{k(n)}: \overline{H}(w/2) \times [2^{k(n)}t, 2^{k(n)+1}t]$  is the restriction of a  $w_n$ -truncated  $\nu_n$  almost hyperbolic manifold that lasts to time  $2^{k(n)+1}t$ .

Now we fix a hyperbolic tower  $\mathcal{T}$  converging to  $H$  as in Proposition 2.19. For any  $t'$  we denote by  $\mathcal{T}(t')$  the  $t'$  time-slice of  $\mathcal{T}$ . It is a  $w/2$ -truncated,  $\nu$ -almost

hyperbolic manifold at time  $t'$ . As we have seen above, given any  $x_\infty \in H$  with  $\text{Vol}_H B(x_\infty, 2) \geq 8w$  and any sequence  $t_n \rightarrow \infty$  then for all  $n$  sufficiently large there are points  $x_n \in \mathcal{T}(t_n)$  so that the  $(M_{t_n}, (1/t_n)g(t_n), x_n)$  converge geometrically to  $(H, x_\infty)$ .

**Claim 2.22.** *Given  $D < \infty$  for all  $t'$  sufficiently large if  $x \in M_{t'}(w_1, +)$  and if  $x \notin \mathcal{T}(t')$ , then  $d_{(1/t')g(t')}(x, \mathcal{T}(t')) > D$ .*

*Proof.* If the result does not hold for some  $D < \infty$  then there is a sequence  $t_n \rightarrow \infty$  and points  $(x_n, t_n) \in \mathcal{M} \setminus \mathcal{T}$  with  $d_{(1/t_n)g(t_n)}(\mathcal{T}(t_n), x_n) \leq D$  and with  $B_{(1/t_n)g(t_n)}(x_n, 2) \geq 8w_1$ . Fix a point  $y_\infty \in H$  and let  $y_n \in \mathcal{T}(t_n)$  be a sequence converging to  $y_\infty$ . Given any point in  $\overline{H}(w)$  there is a sequence  $y_n'' \in \mathcal{T}(t_n)$  converging to this point. Furthermore, there is  $r > 0$  such that if  $y_n'' \in \mathcal{T}(t_n)$  is a sequence converging to a point of  $\overline{H}(w)$ , then for every  $n$  sufficiently large the ball of radius  $r$  about  $y_n''$  in  $(M(t_n), (1/t_n)g(t_n))$  is contained in  $\mathcal{T}(t_n)$ .

Since the  $d_{(1/t_n)g(t_n)}(x_n, \mathcal{T}(t_n))$  are bounded above by  $D < \infty$ , there is an  $R < \infty$  such that for all  $n$  sufficiently large the almost isometric map  $\psi_n: B_H(x_\infty, R) \rightarrow M_{t_n}$  contains the point  $y_n$ . Let  $\tilde{y}_n \in H$  be a point with  $\psi_n(\tilde{y}_n) = y_n$ . The  $\tilde{y}_n$  are at a uniformly bounded distance from  $x_\infty$ , so that passing to a subsequence we can arrange that the  $\tilde{y}_n$  converge to a point  $\tilde{y}_\infty$  in  $H$ . Since  $\text{Vol} B(y_n, 2) \geq 8w_1$ , it follows that  $\tilde{y}_\infty \in \overline{H}(w_1)$  and hence there is a sequence  $y_n'' \in \mathcal{T}(t_n)$  also converging to  $\tilde{y}_\infty$ . This means that  $d_{(1/t_n)g(t_n)}(y_n, y_n'') \rightarrow 0$  as  $n \rightarrow \infty$ . and hence for all  $n$  sufficiently large  $y_n \in \mathcal{T}(t_n)$ . This is a contradiction and establishes the lemma.  $\square$

**Corollary 2.23.** *Given  $w > 0$ ,  $\nu > 0$  sufficiently small and the  $w$ -truncated  $\nu$ -almost hyperbolic tower  $\mathcal{T}$ , the following holds for all  $t$  sufficiently large. If a  $w$ -truncated  $\nu$ -almost hyperbolic manifold  $\overline{H}'(w/2) \subset M_t$  at time  $t$  contains a point  $(x, t)$  not in  $\mathcal{T}$ , and with  $\text{Vol} B_{(1/t)g(t)}(x, 2) \geq 8w_1$ , then  $\overline{H}'(w/2)$  is disjoint from  $\mathcal{T}$ .*

*Proof.* For each  $w' > 0$  there is a  $C(w') < \infty$  such that Removing from  $\overline{H}'(w/2)$  all points that are  $w'$ -volume collapsed on scale 2 yields a connected manifold of diameter  $< C$ . Fix  $w' \ll w$ . It follows from the above, that for all  $t$  sufficiently large, any set  $X \subset M_t$  of diameter  $< C(w')$  in the metric  $(1/t)g(t)$  that contains the point  $(x, t)$  as in the statement of the corollary is disjoint from  $\mathcal{T}$ . This means that if  $t$  is sufficiently large then  $\overline{H}'(w/2) \cap \mathcal{T}(t)$  is contained in the solid torus neighborhoods in  $H'$  around short geodesics, solid torus neighborhoods consisting of points at which  $H'$  is  $w'$ -volume collapsed on scale 2. The boundary tori of these neighborhoods are disjoint from  $\mathcal{T}$  and hence if one of these solid torus neighborhoods meets  $\mathcal{T}(t)$  it must contain a boundary point of  $\mathcal{T}(t)$ . But the metrics on the balls of radius 2 about these points in the metric  $(1/t)g(t)$  within  $\nu$  of hyperbolic metrics on ball of radius 2 in  $H'$  of volume  $8(w/2)^3$ . Since  $w' \ll w$ , this is impossible. Consequently,  $\overline{H}'(w/2)$  is disjoint from  $\mathcal{T}$ .  $\square$

Having fixed  $\mathcal{T}$  converging to  $H$ , we consider all sequences  $(x_n', t_n') \in M_{t_n'}(w_1, +)$  disjoint from  $\mathcal{T}$ . We choose such a sequence whose geometric limit has a minimal number of cusps among all geometric limits of sequences disjoint from  $\mathcal{T}$ .

Let  $H'$  be the geometric limit hyperbolic manifold. By the same argument as before we can also construct a hyperbolic tower  $\mathcal{T}'$ , disjoint from  $\mathcal{T}$ , converging to  $H'$ . to see this first note that by what we have established, for some constant  $T'_1 \geq T_1$ , depending on  $H, H', w, \nu$  the following holds.

1. For any  $t \geq T'_1$ , any point  $x \in M_t(w, +)$  is either contained in  $\mathcal{T}$  or is contained in a  $w/2$ -truncated,  $\nu$ -almost hyperbolic manifold  $\overline{H}''(w/2)$  that is disjoint from  $\mathcal{T}$ .
2. For any such  $H''$  has at least as many cusps as  $H'$ .

We fix  $t'$  equal to one of the  $t'_n > T'_1$ . Suppose inductively, that we have constructed stages of a  $w/2$ -truncated,  $\nu$ -almost hyperbolic tower  $\varphi'_k: \overline{H}'(w/2) \rightarrow M_{2^k t}, k = 0, \dots, k_0$ . We need to show that the  $w/2$ -truncated  $\nu$ -almost hyperbolic manifold that contains the image  $\varphi'_{k_0}(\overline{H}'(w'))$  under the flow from  $2^{k_0} t$  to  $2^{k_0+1} t$  is isometric to  $H'$ . But the hyperbolic manifold  $H_1$  whose truncation  $\overline{H}_1(w/2)$  contains this image is disjoint from the original tower  $\mathcal{T}$ , and thus by the fact that  $H'$  has a minimal of the number of cusps for  $w$ -truncated,  $\nu$ -almost hyperbolic manifolds that are disjoint from  $\mathcal{T}$ , the same argument applies to show that  $H_1 = H'$ . We repeat this inductively to construct a  $w/2$ -truncated,  $\nu$ -almost hyperbolic tower  $\mathcal{T}'$  converging to  $H'$  and disjoint from  $\mathcal{T}$ .

Now we repeat this argument for sequences of points in  $M_t(w, +)$  disjoint from the union of these two towers. Among all such we take one with a limit which has a minimal number of cusps among all such and repeat the argument. At each stage we construct a new hyperbolic tower disjoint from the previous (at least for sufficiently large time).

There is a uniform positive lower bound to the volume of any truncated version of a complete hyperbolic manifold of sectional curvature  $-1/4$ . Since the renormalized volume  $\hat{V}(t)$  limits to  $V(\infty) < \infty$ , it follows that for all  $t$  sufficiently large there is a uniformly bounded number disjoint truncated versions of complete hyperbolic manifolds of finite volume containing  $M_t(w, +)$ . Thus, there is a bound to the number of disjoint hyperbolic towers in the Ricci flow with surgery. This means the above iterative process of constructing towers must terminate after a finite number of steps. This proves:

**Theorem 2.24.** *For every  $w > 0$  and every  $\nu > 0$ , both sufficiently small, the following holds. Given a Ricci flow with surgery  $(\mathcal{M}, G)$  satisfying Assumptions 3.9 and 3.10 there is a finite set of  $w$ -truncated,  $\nu$ -almost hyperbolic towers  $\mathcal{T}_1, \dots, \mathcal{T}_N$  starting at times  $t_1 < \dots < t_N$  and converging to hyperbolic manifolds  $H_1, \dots, H_N$  with the following properties:*

1. The  $\mathcal{T}_i$  are pairwise disjoint.
2. For all  $t$  sufficiently large, the union  $\cup_{i=1}^N \mathcal{T}_i$  contains  $M_t(w, +)$  and is contained in  $M_t(w/4, +)$ .

Now let us consider what happens when we replace  $w$  by a smaller constant  $w' \leq w/2$ .

Consider limits of sequences of based at  $x_n \in M_{t_n}(w', +)$  for a sequence with  $t_n \rightarrow \infty$ . These limits also will be complete hyperbolic 3-manifolds of finite volume. Suppose  $H'$  is one such. Then for an appropriate truncation and for all sufficiently large  $t$  we have embeddings  $\varphi'_t: \overline{H}'(w') \rightarrow M_t$ . The image of this embedding cannot be contained  $M_t(2w', -)$  because of the assumption  $2w' < w_0$ . Thus,  $\varphi'_t(\overline{H}'(w'))$  must have non-empty intersection with one of the  $\mathcal{T}_i(t)$ . (We take  $t > \max(t_1, \dots, t_N)$ .) Since the boundary of  $\varphi'_t(\overline{H}'(w'))$  is disjoint from  $\mathcal{T}_i(t)$ , the  $w$ -truncated  $\nu$ -almost hyperbolic manifold  $\varphi'_t(\overline{H}'(w'))$  must completely contain  $\mathcal{T}_i(t)$ , and that remains true for all  $t' > t$ . This means that  $H'(w') = H_i(w)$  and in fact the only difference between  $\overline{H}'(w')$  and  $\overline{H}_i(w)$  is that in  $\overline{H}'(w')$  we have truncated the cusps further out. This proves that for all  $t$  sufficiently large the towers  $w'$ -truncated  $\nu$ -almost hyperbolic towers are contained in extended versions of the  $\cup_{i=1}^N \mathcal{T}_i$  obtained by extending the embeddings of the  $\overline{H}(w/2)$  to  $\overline{H}(w'/2)$  (which is possible for  $t$  sufficiently large).

**Proposition 2.25.** *For all  $w > 0$  sufficiently small, and, given  $w$ , for all  $t$  sufficiently large, the boundary tori of the intersection of the  $\mathcal{T}_i$  with the time-slice  $M_t$  are incompressible tori in  $M_t$ .*

*Proof.* For a proof of this result see Sections 11 and 12 (especially Theorem 11.1) in [12]. The basic point in the argument is to assume that one of the boundary tori of one of the towers is compressible for all  $t$  sufficiently large. We consider the first-order change in the area of a minimal compressing disk for that boundary torus under the flow. One shows that the area of this disk goes to zero in finite time, which contradicts the fact that the torus exists for all time and is not compressible in a neighborhood, which as  $t \rightarrow \infty$ , converges to a complete hyperbolic manifold with the torus converging to a horospherical torus. Thus, any compressing disk must exit from this region and this provides a positive lower bound to its area. This gives a contradiction.

There are alternative proofs. One is due to Perelman, see in Proposition 8.2 of [27]. A variant of this idea was used by John Lott (see 93.1 in [15]) to give a simpler proof, one that uses the volume of the metric normalized by the minimum of scalar curvature.  $\square$

### 2.3 Locally volume collapsed part of the $(M_t, g(t))$

At this point let us define

$$M_t(w, -) = M_t \setminus \prod_{i=1}^N \mathcal{T}_i.$$

Then for all  $t$  sufficiently large, the manifold  $M_t(w, -)$  is a compact 3-manifold with locally convex boundary consisting of incompressible tori. Using the metric  $(1/t)g(t)$  on this manifold, the boundary has a topological collar neighborhood that contains all the points within distance 1 of the boundary and on which the curvature is close to  $-1/4$  (how close depending on  $t$  with the difference going to zero as  $t \rightarrow \infty$ ).



Also, the diameter of each boundary component is at most  $Kw$  for a constant  $K$  that depends on the limiting hyperbolic manifolds  $H_1, \dots, H_N$  but not on  $t$ . For every  $t$  sufficiently large, and for every  $(x, t) \in M_t(w, -)$ , we have

$$\text{Vol } B(x, t, \rho(x, t)) < w\rho^3(x, t) \quad \text{and}$$

$$Rm|_{B(x, t, \rho(x, t))} \geq -\rho^{-2}(x, t).$$

We take up the study of the  $(M_t(w, -), (1/t)g(t))$  in Part II.

### 3 Local results valid for sufficiently large time

The proofs of Propositions 2.6, 2.8, and 2.9 are based on important technical results reminiscent of the results which go into the proof of the existence of a Ricci flow with surgery defined for all time. To establish the existence of limits for the rescaled flows we must show that the rescaled metrics have uniform non-collapsing at the base point and have bounded curvature at bounded distance from the base point. These are the content of the local results in this section. While the conclusions are the same as the results for bounded time, these results are different in that, unlike the former results where the constants decay as time goes to  $\infty$ , the results here apply uniformly for all time sufficiently large. But to compensate for this, they are local, requiring a curvature and volume hypotheses near the central point around which we are working.

**In this section Ricci flow with surgery means a Ricci flow with surgery as in the hypothesis of Theorem 1.2. Later in this section we will put an additional requirement on the surgery control function  $\delta(t)$ .**

#### 3.1 First local result

The first result presents local versions of the non-collapsing result, the canonical neighborhood result and the bounded curvature at bounded distance result. These results do not follow from the results in Chapters 15 – 17 of [21] since we are not assuming a finite upper bound on the time. Rather, here we assume that we are working near a parabolic neighborhood where the curvature and volume are controlled. This result is Proposition 6.3 of [27].

**Proposition 3.1.** *For every  $A < \infty$  there are constants  $\kappa > 0$ ,  $K_1 < \infty$ ,  $K_2 < \infty$  and  $\bar{r} > 0$  depending on  $A$  and for each  $t_0 < \infty$  there is a constant  $\bar{\delta}' = \bar{\delta}'_A(t_0) > 0$ , depending as the notation indicates on  $A$  and  $t_0$ , such that the following hold. Suppose that we have a Ricci flow with surgery satisfying Corollary 15.10 from [21]. Suppose that for some  $t_0 < \infty$  the surgery control function  $\bar{\delta}(t)$  satisfies  $\bar{\delta}(t_0/2) \leq \bar{\delta}'$ . Suppose also that:*

- (i) *For some  $r_0 \leq \sqrt{t_0/2}$  the Ricci flow with surgery contains the entire parabolic neighborhood  $P = P(x_0, t_0, r_0, -r_0^2)$ ,*
- (ii)  *$|Rm(x, t)| \leq r_0^{-2}$  for all  $(x, t) \in P$ , and*

(iii)  $\text{Vol } B(x_0, t_0, r_0) \geq A^{-1}r_0^3$ .

Then:

1. The Ricci flow with surgery is  $\kappa$ -non-collapsed on all scales  $\leq r_0$  at every  $(x, t_0) \in B(x_0, t_0, Ar_0)$ .
2. Any  $(x, t_0) \in B(x_0, t_0, Ar_0)$  with  $R(x, t_0) \geq K_1 r_0^{-2}$  has a  $(C, \epsilon)$ -canonical neighborhood.
3. If  $r_0 \leq \bar{r}\sqrt{t_0}$ , then  $R(x, t_0) \leq K_2 r_0^{-2}$  for all  $(x, t_0) \in B(x_0, t_0, Ar_0)$ .

**Proof of non-collapsing.** To prove the  $\kappa$ -non-collapsing result we need to consider a localized version of the arguments in Sections 6, 8, and 16 of [21]. The idea is, given  $(x, t_0) \in B(x_0, t_0, Ar_0)$ , to find a path from  $(x, t_0)$  to a point of  $B(x_0, t_0 - r_0^2/2, r_0/10)$  whose  $\ell$ -length (see the next paragraph for the definition of  $\ell$ -length) is bounded above by a constant depending only on  $A$ . From this, the curvature control on  $P(x_0, t_0, r_0, -r_0^2)$ , and the volume control on  $B(x_0, t_0, r_0)$ , we easily establish the non-collapsing result by the standard argument using monotonicity of reduced volume as in Theorem 8.1 of [21].

**Definition 3.2.** Recall call Perelman's  $\mathcal{L}$ -length for a Ricci flow. Given a Ricci flow  $(M, g(t))$ , we view the flow as a metric on the horizontal sub-bundle in the tangent bundle of  $M \times [a, b]$  whose value on the horizontal tangent space at  $(x, t)$  is  $g_t(x)$ . Let  $\gamma(\tau)$   $0 \leq \tau \leq \tau_0$  be a path starting at time  $t_0$  and defined by backwards time in a Ricci flow, in the sense that  $\gamma(\tau) \in M \times \{t_0 - \tau\}$ . Then

$$\mathcal{L}(\gamma) = \int_0^{\tau_0} \sqrt{\tau} (R(\gamma(\tau)) + |\dot{\gamma}(\tau)|^2) d\tau,$$

where  $\dot{\gamma}(\tau)$  is the horizontal component of the tangent vector to  $\gamma$ . We also define the reduced  $\mathcal{L}$ -length denoted by

$$\bar{\mathcal{L}} = \frac{1}{2\sqrt{\tau_0}} \mathcal{L}.$$

We shall denote by  $\ell$  the reduced length function based at  $(x, t_0)$ . Its value at a point  $(y, t)$  with  $t < t_0$  is the infimum over all paths  $\gamma$  starting at  $(x, t_0)$  and parametrized by backwards time and ending at  $(y, t)$  of  $\mathcal{L}(\gamma)$ .

We shall study  $\ell$  on  $B(x_0, t_0, r_0/10) \times \{t_0 - r_0^2/2\}$ . To do this we need to invoke a cut-off function with some control on its first and second derivatives. The following is elementary (see Equation 8.1 of [26]) and provides the control function.

**Claim 3.3.** For any  $A < \infty$  there exist a constant  $C(A) < \infty$  and a smooth function  $\phi: (-\infty, \infty) \rightarrow [1, \infty]$  such that  $\phi = 1$  on  $(-\infty, 1/20)$  and  $\phi$  increases monotonically to  $+\infty$  on  $[1/10, \infty)$  with

$$2(\phi')^2/\phi\phi'' \geq (2A + 300)\phi' - C(A)\phi.$$

Fix  $(x, t_0) \in B(x_0, t_0, Ar_0)$ , and let us consider the Ricci flow with surgery where we rescale the metric and time by  $r_0^{-2}$ . We set  $\tau = t_0/r_0^2 - t$ . For any  $\tau \leq 1/2$  and any path  $\gamma(\tau')$  defined for  $0 \leq \tau' \leq \tau$ , we define

$$h(\gamma) = \phi(d_t(x_0, \gamma(\tau)) - A(1 - 2\tau)) (\bar{L}(\gamma) + 2\sqrt{\tau}),$$

where  $\bar{L}(\gamma) = 2\sqrt{\tau}\mathcal{L}(\gamma)$ . Then for any  $\tau \in [t_0/r_0^2 - 1/2, t_0/r_0^2]$ , we define  $h(y, \tau)$  to be the infimum of  $h(\gamma)$  over all paths  $\gamma$  parametrized by backwards time starting at  $(x, t_0/r_0^2)$  and ending at  $(y, (t_0/r_0^2) - \tau)$ .

**Claim 3.4.** *Suppose that for every  $\tau \in (0, 1/2]$  and for every minimum  $y$  of the function  $h(\cdot, \tau)$  every minimizing  $\mathcal{L}$ -geodesic from  $(x, t_0/r_0^2)$  to  $(y, t_0/r_0^2 - \tau)$  is contained in the smooth part of the Ricci flow with surgery. Then, denoting the minimum of  $h(\cdot, \tau)$  by  $h_0(\tau)$ , we have*

$$h_0(\tau) \leq 2\sqrt{\tau} \exp(C(A)(\tau) + 100\sqrt{\tau}).$$

*Proof.* First notice that the maximum principle and the fact that the initial conditions of the original Ricci flow with surgery are normalized, imply that in the original flow  $R(z, t) \geq -\frac{6}{1+4t}$ . Since  $t_0 \geq 2r_0^2$ , on the interval  $[t_0 - r_0^2/2, t_0]$  the scalar curvature is at least  $-r_0^{-2}$ . This means that in the rescaled flow on the interval  $[t_0/r_0^2 - 1/2, t_0]$  (i.e., the interval  $0 \leq \tau \leq 1/2$ ), the scalar curvature is at least  $-1$ . Since  $\tau \leq 1/2$ , it follows easily that  $\bar{L}(y, \tau) + 2\sqrt{\tau} > 0$  on the interval  $0 \leq \tau \leq 1/2$ . Thus, the minimum of  $h(\cdot, \tau)$  occurs in the region  $B(x_0, t, 1/10)$ . Direct computation using the inequality for  $\phi$  from Claim 3.3 gives

$$\frac{d}{d\tau} \log \left( \frac{h_0(\tau)}{\sqrt{\tau}} \right) \leq C(A) + \frac{50}{\sqrt{\tau}}.$$

As  $\tau \rightarrow 0^+$  we see that  $\lim_{\tau \rightarrow 0^+} \mathcal{L}(x, \tau) = 0$ , so that  $\lim_{\tau \rightarrow 0^+} h(x, \tau)/\sqrt{\tau} = 2$  and hence  $\lim_{\tau \rightarrow 0^+} h_0(\tau)/\sqrt{\tau} \leq 2$ . From this and the above differential inequality, it follows that

$$h_0(\tau) \leq 2\sqrt{\tau} \exp(C(A)\tau + 100\sqrt{\tau}).$$

□

Since the minimum at time  $t = t_0/r_0^2 - \tau$  must occur in the  $B(x_0, t, A(1 - 2\tau) + 1/10)$ , under the assumption of Claim 3.4, it follows that for each  $\tau \leq 1/2$  there is a path  $\gamma(\tau')$  parametrized by backwards time and contained in the smooth part of the Ricci flow with surgery, a path beginning at  $(x, t_0/r_0^2)$  and ending at a point of  $B(x_0, t_0/r_0^2 - \tau, A(1 - 2\tau) + 1/10)$  with

$$2\sqrt{\tau}\mathcal{L}(\gamma) \leq 2\sqrt{\tau} \exp(C(A)\tau + 100\sqrt{\tau}).$$

Again assuming the hypotheses of Claim 3.4 hold, this means that in the original Ricci flow with surgery for each  $\tau \leq r_0^2/2$ , there is a path  $\gamma(\tau')$ ,  $0 \leq \tau' \leq \tau$ , contained in the smooth part of the Ricci flow with surgery starting at  $(x, t_0)$  and ending at a point of  $B(x_0, t_0 - \tau, r_0/10 + A(1 - 2\tau/r_0^2))$  with

$$2\sqrt{\tau} \int_0^\tau \sqrt{\tau'} (R(\gamma(\tau')) + |\dot{\gamma}(\tau')|^2) d\tau' \leq 2\sqrt{\tau} \exp(C(A)\tau + 100\sqrt{\tau}) r_0^2,$$

where  $\bar{\tau} = \tau/r_0^2$ . In particular, we have a path  $\gamma$  which starts at  $(x, t_0)$  and ends at a point of  $B(x_0, t_0 - r_0^2/2, 1/10)$  and which satisfies this inequality with  $\bar{\tau} = 1/2$ .

The next step is to show that the hypotheses of Claim 3.4 always hold. In order to see this we must show that paths that meet the surgery disks have larger value than this hoped-for minimum. This is the reason for the extra condition on  $\bar{\delta}(t)$ .

**Claim 3.5.** *There is a constant  $\bar{\delta}'_A(t_0) > 0$  depending only on  $A$  and  $t_0$  such that the following holds. We fix*

$$\tilde{C}_0 = \tilde{C}_0(A) = 2\exp\left(C(A)/2 + 100\sqrt{1/2}\right).$$

*If  $\bar{\delta}(t) \leq \bar{\delta}'_A(t_0)$  for all  $t \in [t_0/2, t_0]$ , then, for any  $\tau \leq r_0^2/2$  and any path  $\gamma(\tau')$  starting at  $(x, t_0)$  parametrized by backwards time and meeting the closure of a surgery cap, we have*

$$\mathcal{L}(\gamma) \geq \tilde{C}_0 r_0,$$

*and hence*

$$2\sqrt{\tau}\mathcal{L}(\gamma) \geq 2\sqrt{\bar{\tau}}\tilde{C}_0 r_0^2,$$

*where  $\bar{\tau} = \tau/r_0^2$ .*

*Proof.* Since  $r_0^2 \leq t_0/2$  if  $t_0 \in [\epsilon^{2i}, \epsilon^{2i+1})$ , then  $t_0 - r_0^2 \geq \epsilon^{2(i-1)}$ . Thus, it follows immediately from Section 16.5 and in particular Proposition 16.13 of [21] that given a positive constant  $\tilde{C}$  there is a constant  $\bar{\delta}'(\tilde{C}, t_0)$  such for any  $t_\gamma \in [t_0 - r_0^2, t_0]$  and any path  $\gamma$  parametrized by  $0 \leq \tau \leq t_0 - t_\gamma$  we have the following. If  $\bar{\delta}(t) \leq \bar{\delta}'(\tilde{C}, t_0)$  for all  $t \in [t_0/2, t_0]$  and if  $\gamma$  meets a surgery cap, then

$$\mathcal{L}(\gamma) > \tilde{C}\sqrt{t_0} \geq 2\tilde{C}r_0.$$

We then define  $\bar{\delta}'_A(t_0)$  to be the constant  $\bar{\delta}'(\tilde{C}_0(A), t_0)$ , and we suppose that the surgery scale constant  $\bar{\delta}(t)$  is  $\leq \bar{\delta}'_A(t_0)$  for all  $t \in [t_0/2, t_0]$ .  $\square$

Arguing as in Chapter 16 of [21] completes the proof that the hypotheses of Claim 3.4 are always satisfied provided that  $\bar{\delta}(t) \leq \bar{\delta}'_A(t_0)$  for all  $t \in [t_0/2, t_0]$ .

We conclude the following:

**Corollary 3.6.** *Suppose that  $\bar{\delta}(t) \leq \bar{\delta}'_A(t_0)$  for all  $t \in [t_0/2, t_0]$ . Then for every  $0 \leq \tau \leq r_0^2/2$ , there is a path  $\gamma$  parametrized by backwards time from  $(x, t_0)$  to a point  $(y, t_0 - \tau)$  contained in  $B(x_0, t_0 - \tau, r_0/10)$  with the property that*

$$2\sqrt{\tau}\mathcal{L}(\gamma) \leq 2\sqrt{\bar{\tau}}\exp\left(C(A)\bar{\tau} + 100\sqrt{\bar{\tau}}\right)r_0^2,$$

*where  $\bar{\tau} = \tau/r_0^2$ .*

This is the main step in the proof of  $\kappa$ -non-collapsing at all points of  $B(x_0, t_0, Ar_0)$ . Because of the bound on the Riemann curvature on  $P(x_0, t_0, r_0, -r_0^2)$ , for any  $y \in B(x_0, t_0 - r_0^2/2, r_0/10)$  the parabolic neighborhood  $P(y, t_0 - r_0^2/2, r_0/4, -r_0^2/2)$  is contained in  $P(x_0, t_0, r_0, -r_0^2)$ . Thus, connecting  $(y, t_0 - r_0^2/2)$  to any point  $z \in$

$B(y, t_0 - r_0^2/2, r_0/4) \times \{t_0 - r_0^2\}$  by a  $g(t_0 - r_0^2/2)$ -geodesic gives an upper bound on  $\ell$  on  $B(y, t_0 - r_0^2/2, r_0/4) \times \{t_0 - r_0^2\}$ , a finite upper bound that depends only on  $A$ . Also, the curvature bound on  $P(x_0, t_0, r_0, -r_0^2)$  and the volume lower bound of  $B(x_0, t_0, r_0)$  imply that the volume of  $B(y, t_0 - r_0^2/2, r_0/4) \times \{t_0 - r_0^2\}$  is bounded below by a constant only depending on  $A$ . This proves that there is an open subset in the  $(t_0 - r_0^2)$ -time slice whose reduced volume is bounded below by a constant depending only on  $A$ . Using monotonicity of reduced volume as in Corollary 6.80 of [21], this proves the existence of a  $\kappa$ , depending only on  $A$ , such that  $(x, t_0)$  is  $\kappa$ -on-collapsed on all scales  $\leq r_0$ .

**Proof of the canonical neighborhood result.** Now let us turn to the second statement, the existence of a canonical neighborhood threshold in  $B(x_0, t_0, Ar_0)$  depending only on  $A$ . Fix  $A$  and suppose that there is no such threshold  $K_1$ . Then there are a sequence of constants  $K_{1,n}$  tending to  $\infty$  as  $n \rightarrow \infty$ , a sequence of Ricci flow with surgery  $(\mathcal{M}_n, G_n)$ , a sequence of points  $(x_{0,n}, t_{0,n}) \in \mathcal{M}_n$ , and constants  $r_{0,n}$  such that the hypotheses of the proposition hold, and there are points  $(y_n, t_{0,n}) \in B(x_{0,n}, t_{0,n}, Ar_{0,n})$  at which the scalar curvature is at least  $K_{1,n}r_{0,n}^{-2}$  but  $(y_n, t_{0,n})$  does not have a  $(C, \epsilon)$ -canonical neighborhood. Since at and before any fixed finite time  $t$  there is uniform finite canonical neighborhood threshold  $r^{-2}(t)$ , and since  $r_{0,n} \leq \sqrt{t_{0,n}/2}$ , it follows that  $t_{0,n} \rightarrow \infty$  as  $n \rightarrow \infty$ .

We apply Lemma 9.37 of [21] to the function  $R(x, t)d_t^{-2}(x_0, x)$  and consider only with points that violate the  $(C, \epsilon)$ -canonical neighborhood assumption. This allows us to conclude from the existence of  $(y_n, t_{0,n}) \in B(x_{0,n}, t_{0,n}, Ar_{0,n})$  that for all  $n$  sufficiently large, there is a point  $(\bar{x}_n, \bar{t}_n) \in B(x_{0,n}, \bar{t}_n, 2Ar_{0,n})$  with  $\bar{t}_n \in [t_{0,n} - r_{0,n}^2/2, t_{0,n}]$  such that, setting  $\bar{Q}_n = R(\bar{x}_n, \bar{t}_n)$ , we have  $\bar{Q}_n \geq K_{1,n}r_{0,n}^{-2}$ , the point  $(\bar{x}_n, \bar{t}_n)$  does not have a  $(C, \epsilon)$ -canonical neighborhood and each point

$$(x, t) \in \tilde{P}_n = \left\{ (x, t) \mid \bar{t}_n - K_{1,n}\bar{Q}_n^{-1} \leq t \leq \bar{t}_n, \ d_t(x_0, x) \leq d_{\bar{t}_n}(x_0, \bar{x}_n) + K_{1,n}^{1/2}\bar{Q}_n^{-1/2} \right\},$$

with  $R(x, t) \geq 4\bar{Q}_n$  does have a  $(C, \epsilon)$ -canonical neighborhood.

Clearly, by the curvature bound on  $P(x_{0,n}, t_{0,n}, r_{0,n}, -r_{0,n}^2)$  there is a universal constant  $\alpha > 0$  such that for any  $t \in [t_{0,n} - 3r_{0,n}^2/4, t_0]$  we have

$$P(x_{0,n}, t, \alpha r_{0,n}, -(\alpha r_{0,n})^2) \subset P(x_{0,n}, t_{0,n}, r_{0,n}, -r_{0,n}^2).$$

Also, by volume comparison there is a constant  $A' < \infty$  depending only on  $A$  such that for every  $t \in [t_{0,n} - 3r_{0,n}^2/4, t_0]$  we have

$$\text{Vol } B(x_{0,n}, t, \alpha r_{0,n}) \geq (A')^{-1}(\alpha r_{0,n})^3.$$

Since

$$2Ar_{0,n} + K_{1,n}^{1/2}\bar{Q}_n^{-1/2} \leq \frac{2A+1}{\alpha}(\alpha r_{0,n}),$$

applying the conclusion in Part 1 of this proposition with  $A$  replaced by the maximum of  $A'$  and  $(2A+1)/\alpha$ , we conclude that for all  $n$  sufficiently large, the Ricci flow with surgery  $(\mathcal{M}, G)$  is  $\kappa'$ -non-collapsed at every point of  $\tilde{P}_n$  for a universal  $\kappa'$  that depends only on  $A$ .

Use  $(\bar{x}_n, \bar{t}_n)$  as the central point, shift  $\bar{t}_n$  to 0 and scale by  $\bar{Q}_n$ , and restrict attention to  $t \leq 0$ . This gives us a sequence of based Ricci flows with surgery  $(\mathcal{M}'_n, G'_n, (\bar{x}_n, 0))$  defined for  $-\bar{Q}_n \bar{t}_n \leq t \leq 0$ . The conditions established above translate to the following:

1.  $R(\bar{x}_n, 0) = 1$ .
2. Every point of  $\bar{Q}_n \tilde{P}_n$  with curvature  $\geq 4$  has a  $(C, \epsilon)$ -canonical neighborhood.
3. At every point of  $\bar{Q}_n \tilde{P}_n$  the Ricci flow with surgery is  $\kappa'$ -non-collapsed on scales  $\leq \bar{Q}_n^{1/2} r_{0,n}$ .

Also, we have  $\bar{Q}_n^{1/2} r_{0,n} \geq K_{1,n}^{1/2}$ , so that  $\bar{Q}_n^{1/2} r_{0,n} \rightarrow \infty$  as  $n \rightarrow \infty$ . It follows that for any  $D < \infty$  for all  $n$  sufficiently large,  $\mathcal{B}_n(D) = \cup_{-D \leq t \leq 0} B(\bar{x}_n, t, D) \subset \bar{Q}_n \tilde{P}_n$ .

Now the argument is identical to the one given in Section 17 of [21]. We sketch the main points of this argument.

**Claim 3.7.** *Given  $D < \infty$  there is  $Q(D) < \infty$  such that  $R < Q(D)$  on  $B(\bar{x}_n, 0, D)$  for all  $n$  sufficiently large.*

*Proof.* With one modification, this exactly Theorem 10.2 in [21]. The modification involves the hypothesis in the reference (reformulated in the notation here) that the  $\bar{Q}_n \rightarrow \infty$ . This hypothesis together with curvature pinching is used to show that after rescaling and passing to partial limits of subsequences the curvature of such limits is non-negative. This non-negativity is used at the very last step when Hamilton's strong maximum principle is invoked to rule out a non-negatively curved Ricci flow with the final slice being an open subset of a non-flat cone. Here we do not know that  $\bar{Q}_n$  tends to infinity, only that  $\bar{Q}_n t_{0,n} \geq \bar{Q}_n r_{0,n}^{-2} \geq K_{1,n} \rightarrow \infty$ . On the other hand, we have the stronger curvature pinching hypothesis as given in Inequality (1.2). This is enough to conclude that the curvature of the limits is non-negative, and hence the proof of Theorem 10.2 applies in this case as well.  $\square$

**Claim 3.8.** *For any  $D < \infty$  there is  $\eta(D) > 0$  such that for all  $n$  sufficiently large the Ricci flow  $(\mathcal{M}'_n, G'_n)$  contains the entire parabolic neighborhood  $P(\bar{x}_n, 0, D, -\eta(D))$  and has curvature bounded by  $2Q(D)$  there.*

*Proof.* It follows immediately from the previous claim, the fact that any point with  $R \geq 4$  in has a  $(C, \epsilon)$ -canonical neighborhood, and Inequality (1.3) that there is a constant  $\eta(D) > 0$  such that for all  $n$  sufficiently large on any backwards flow line beginning at a point of  $B(\bar{x}_n, 0, D)$  and moving backwards at most time  $\eta(D)$  we have  $R \leq 2Q(D)$ . To complete the proof we must show that for all  $n$  sufficiently large, no such flow line can end in a surgery cap, or equivalently we must show that the entire parabolic neighborhood  $P(\bar{x}_n, 0, D, -\eta(D))$  is contained in the Ricci flow with surgery  $(\mathcal{M}'_n, G'_n)$ . This is exactly the argument in Lemma 17.7 of [21]: were there such a flow line for some  $n$  sufficiently large ending in a surgery cap, then  $(\bar{x}_n, 0)$  would have a  $(C, \epsilon)$ -canonical neighborhood in  $(\mathcal{M}'_n, G'_n)$ , and hence in the original Ricci flow with surgery  $(\mathcal{M}_n, G_m)$ , the point  $(\bar{x}_n, \bar{t}_n)$  would have a  $(C, \epsilon)$ -canonical neighborhood, contrary to our assumption.  $\square$

Since the original Ricci flows with surgery,  $(\mathcal{M}_n, G_n)$ , are  $\kappa'$ -non-collapsed at  $(\bar{x}_n, \bar{t}_n)$ , these curvature bounds on parabolic neighborhoods allow us to extract a smooth limit  $(M_\infty, \bar{x}_\infty)$  of a subsequence of the  $t = 0$  time-slices of the based Ricci flows with surgery  $(\mathcal{M}_n, G_n, (\bar{x}_n, 0))$ . This limit is a complete Riemannian 3-manifold of non-negative curvature with the property that every point  $y$  with  $R(y) \geq 4$  has a  $(C, \epsilon)$ -canonical neighborhood, meaning that either the point is contained in the core of a  $(C, \epsilon)$ -cap or is the central point of the final time-slice of a strong  $\epsilon$ -neck. It then follows from Lemma 2.20 (or Corollary 2.21) in [21] that  $M_\infty$  has bounded curvature.

Now arguing as in Section 11.2 of [21] we extend the limit manifold  $(M_\infty, \bar{x}_\infty)$  to a limiting Ricci flow with bounded curvature defined backwards for a small amount of time. The length of time depends on the curvature bound on  $M_\infty$ . Again we invoke Lemma 17.7 of [21] to rule out the appearance of surgery caps as we move backward. Then repeating the argument as in Section 11.2 (see especially Theorem 11.8) of [21] (and invoking Lemma 17.7 from [21] to rule out the appearance of surgery caps as we move backwards each step), we produce a limiting Ricci flow defined for all time  $-\infty < t \leq 0$ . This limit is a  $\kappa$ -solution. But that contradicts the fact that none of the  $(\bar{x}_n, \bar{t}_n) \in \mathcal{M}_n$  have  $(C, \epsilon)$ -canonical neighborhoods, and completes the proof of the second conclusion in the proposition.

**Proof of bounded curvature.** Now let us consider the third conclusion, the curvature bound at bounded distance from  $(x_0, t_0)$  after rescaling. Suppose that there is no  $\bar{r}$  as required. Then there is a sequence  $\bar{r}_n \rightarrow 0$  as  $n \rightarrow \infty$ , and for each  $n$  a Ricci flow with surgery  $(\mathcal{M}_n, G_n)$ , points  $(x_{0,n}, t_{0,n}) \in \mathcal{M}_n$ , constants  $r_{0,n} \leq \bar{r}_n \sqrt{t_0}$ , and points  $(y_n, t_{0,n})$  with  $R(y_n, t_{0,n}) = \hat{K}_n r_{0,n}^{-2}$  where  $\hat{K}_n \rightarrow \infty$  as  $n \rightarrow \infty$ . Take a shortest geodesic  $\gamma_n$  from  $(x_{0,n}, t_{0,n})$  to  $(y_n, t_{0,n})$ . For all  $n$  sufficiently large let  $(z_n, t_{0,n})$  be the last point of  $\gamma_n$  with  $R(z_n, t_{0,n}) = K_1 r_{0,n}^{-2}$ . (There are such points since  $R(x_{0,n}, t_{0,n}) \leq 6r_{0,n}^{-2}$  and  $R(y_n, t_{0,n}) \geq \hat{K}_n r_{0,n}^{-2} > K_1 r_{0,n}^{-2}$  for all  $n$  sufficiently large.) Now we apply the version of Theorem 10.2 of [21] described above (using the stronger curvature pinching hypothesis) to  $(z_n, t_{0,n})$  and  $(y_n, t_{0,n})$ . The ratio  $R(y_n, t_{0,n})/R(z_n, t_{0,n}) = \hat{K}_n/K_1$  and hence tends to  $\infty$  as  $n \rightarrow \infty$ , and  $R(z_n, t_{0,n})t_{0,n} \geq K_1 r_{0,n}^{-2}t_{0,n} \geq K_1 \bar{r}_n^{-2}$  also tends to  $\infty$  as  $n \rightarrow \infty$ . The only other difference is that in Theorem 10.2 of [21] we assumed that the  $(C, \epsilon)$ -canonical neighborhood assumption holds for the entire flow whereas here, we only have it in  $B(x_0, t_0, Ar_0)$  for each  $A$ . But an examination of the proof given in Section 10 of [21] shows that is all that is necessary in order to take the requisite limits to produce the cone limit at the final time. Then the existence of the  $(C, \epsilon)$ -canonical neighborhoods around the points at final time close to the cone point gives the contradiction. This completes the proof of Proposition 3.1.

Now we impose the extra condition on the surgery control parameter  $\bar{\delta}(t)$ .

**Assumption 3.9.** From now on we assume that the surgery control function  $\bar{\delta}(t)$  is at most  $\Delta(t)$  as given in Corollary 15.10 of [21] and also at most the constant  $\bar{\delta}_{2t}^t(2t)$  from the previous lemma. Thus, for every  $t_0 > 0$ , the previous result holds at time  $t_0$  for any  $A \leq t_0$  since we also have required (in [21]) that  $\bar{\delta}(t)$  be a weakly monotone decreasing function of  $t$ .

### 3.2 Second Local Result.

Before stating the second proposition we introduce one new piece of notation. Let  $\hat{h}(t_0) = \sup_{t \in [t_0/2, t_0]} h(t)$ .

**Assumption 3.10.** The extra condition we require of  $h(t)$  is that for all  $t < \infty$  we have  $\hat{h}(t) \leq \bar{\delta}^2(t)r(t)$ . From now on implicitly a Ricci flow with surgery is assumed to be as in the hypothesis of Theorem 1.2 and in addition, its surgery control function satisfies Assumption 3.9 and its surgery curvature function  $h(t)$  is assumed to satisfy Assumption 3.10.

The next result is Lemma 6.4 of [27]. It provides parabolic neighborhoods on which the solution is defined and bounded. Recall that  $C$  is one of the canonical neighborhood constants and that  $r(t)$  is the canonical neighborhood threshold function.

**Proposition 3.11.** *There exist  $\tau > 0$ ,  $\bar{r}_1 > 0$ , and  $K < \infty$  such that the following holds for any Ricci flow with surgery  $(\mathcal{M}, G)$ . Suppose that  $r_0, t_0$  satisfy  $4C\hat{h}(t_0) \leq r_0 \leq \bar{r}_1\sqrt{t_0}$ . Assume that  $B(x_0, t_0, r_0)$  has sectional curvatures at least  $-r_0^{-2}$  and the volume of any sub-ball  $B(x, t_0, r) \subset B(x_0, t_0, r_0)$  is at least  $(1 - \epsilon)$  times the volume of the Euclidean ball of the same radius. Then the Ricci flow with surgery contains the entire parabolic neighborhood  $P(x_0, t_0, r_0/4, -\tau r_0^2)$  and satisfies  $R < Kr_0^{-2}$  on this parabolic neighborhood.*

The proof of this result is divided into two cases.

#### 3.2.1 Case 1: $4C\hat{h}(t_0) \leq r_0 \leq r(t_0)$

We shall show that in this case, as long as  $\tau \leq C^{-3}/8$  and  $K \geq 8C^2$ , the result holds. (Notice that since  $r(t) \rightarrow 0$  as  $t \rightarrow \infty$ , given any  $\bar{r}_1 > 0$  for all  $t_0$  sufficiently large, we have  $r(t_0) < \bar{r}_1\sqrt{t_0}$ .)

**Claim 3.12.**  $R < 4C^2r_0^{-2}$  on  $B(x_0, t_0, r_0/2)$ .

*Proof.* Suppose that for some  $(x, t_0) \in B(x_0, t_0, r_0/2)$  we have  $R(x, t_0) \geq 4C^2r_0^{-2}$ . Since we have no components of positive curvature, the only canonical neighborhoods we have are strong  $\epsilon$ -necks and  $(C, \epsilon)$ -caps. It then follows from the inequality for  $R(x, t_0)$  that  $(x, t_0)$  is the central point of a strong  $\epsilon$ -neck or is contained in a  $(C, \epsilon)$ -cap. If  $(x, t_0)$  is contained in a  $(C, \epsilon)$ -cap, then the diameter of this cap is at most  $CR(x_0, t_0)^{-1/2} \leq r_0/2$ . Hence, in this case  $B(x_0, t_0, r_0)$  contains the  $(C, \epsilon)$ -cap and consequently contains final time-slice of the strong  $\epsilon$ -neck that forms a neighborhood of the non-compact end of this cap. If  $(x_0, t_0)$  is the center of a strong  $\epsilon$ -neck, then the radius of the central 2-sphere in that neck is approximately  $R(x_0, t_0)^{-1/2} \leq C^{-1}r_0/2$ . Thus, in either case  $B(x_0, t_0, r_0)$  contains a ball centered at a central point of an  $\epsilon$ -neck whose radius is larger than the diameter of the central 2-sphere of the  $\epsilon$ -neck. This is impossible by the volume assumption on all sub-balls of  $B(x_0, t_0, r_0)$ . This completes the proof of the claim.  $\square$



By Inequality (1.3), if  $(x, t)$  has a canonical neighborhood then  $|\partial R(x, t)/\partial t| \leq CR^2(x, t)$ . In particular, this inequality holds at any point  $(x, t) \in P(x_0, t_0, r_0/2, -r_0^2)$  with  $R(x, t) \geq r_0^{-2} \geq r(t_0)^{-2} \geq r(t)^{-2}$ . Since  $R(x, t_0) < 4C^2r_0^{-2}$  for all  $(x, t) \in B(x_0, t_0, r_0/2)$  it follows that we have  $R < 8C^2r_0^{-2}$  on  $P(x_0, t_0, r_0/2, -C^{-3}r_0^2/8)$  and in particular, the flow contains the entire backward parabolic neighborhood. The reason is that if a backwards flow line starting in  $B(x_0, t_0, r_0/2)$  hits a surgery cap at time  $t \geq t_0 - C^{-3}r_0^2/8$ , then the scalar curvature at the point of the cap is at least  $3h^{-2}(t)/4 \geq 3\hat{h}^{-2}(t_0)/4$ . But the scalar curvature is bounded above by  $8C^2r_0^{-2}$  and  $r_0 \geq 4C\hat{h}(t_0)$  so that the scalar curvature along this flow line is bounded above by  $\hat{h}^{-2}(t_0)/2$  which is a contradiction.

Thus, it remains to consider:

### 3.2.2 Case 2: $r(t_0) < r_0 \leq \bar{r}_1\sqrt{t_0}$ .

Notice that since  $r(t_0) > 4C\hat{h}(t_0)$  for  $t_0$  sufficiently large, it is automatic that  $4C\hat{h}(t_0) < r_0$  in this case. To treat this case we need a couple of preliminary results. The next lemma is based on the fact that the asymptotic volume of any  $\kappa$ -solution is zero. It gives strong curvature and distance distortion control for sufficiently small backwards time.

**Lemma 3.13.** *Given  $w > 0$  there are  $\tau_0 > 0$ ,  $0 < B, C < \infty$  and  $w' > 0$  such that the following hold. Suppose that  $(\mathcal{U}, G)$  is a generalized 3-dimensional Ricci flow and suppose that for some  $0 < \tau \leq \tau_0$  the open set  $\mathcal{B} = \cup_{0 \leq t \leq \tau} B(x_0, -t, .95) \subset \mathcal{U}$  has compact closure in  $\mathcal{U}$ . Suppose also that the volume of  $B(x_0, 0, .95)$  is at least  $w$  and that  $Rm(x, t) \geq -1$  on  $\mathcal{B}$ . For any  $0 < r \leq .95$  we denote  $\mathcal{B}(r)$  the union  $\cup_{0 \leq t \leq \tau} B(x_0, -t, r)$ . Then:*

1. *For every  $t \in [0, \tau]$ , the entire forward parabolic neighborhood  $P(x_0, -t, .9, t)$  exists and is a subset of  $\mathcal{B}$ .*
2.  *$R(x, -t) < B + C/(\tau - t)$  for all  $(x, -t) \in \mathcal{B}(1/3)$ .*
3. *For any  $t' \in [0, \tau]$  and any  $1/100 \leq r \leq 1/3 - 1/100$  we have*

$$P(x_0, 0, r - 1/100, -t') \subset \mathcal{B}(r) \cap \{(x, -t) \mid 0 \leq t \leq t'\} \subset P(x_0, 0, r + 1/100, -t').$$

4.  *$\text{Vol } B(x_0, -\tau, 1/4) \geq w'$ .*

*Proof.* First, notice that the fact that  $\mathcal{B}$  has compact closure means that if  $\ell$  is maximal flow line in  $\mathcal{B}$  for the vector field  $\chi$  (which recall is part of the structure of the generalized Ricci flow) then each end of  $\ell$  either is compact and contained  $B(x_0, 0, .95)$  or  $B(x_0, -\tau, .95)$  or is non-compact and is compactified by adding a point of the form  $(y, -t) \in \overline{B(x_0, -t, .95)}$  with  $d_{-t}(x_0, y) = .95$ .

We take  $\tau \leq \ln(\sqrt{1.01})$ . Since  $Rm \geq -1$ , and hence  $Ric \geq -2$  on  $\mathcal{B}$ , if  $\gamma(s)$  is any smooth path in  $B(x_0, -t, .95)$ , then the derivative of the length of  $\gamma$  with respect to time is at most twice the length of  $\gamma$ . It follows easily that if  $(y, -t) \in B(x_0, -t, r)$  for some  $r \leq (.95)\exp(-2t)$  then the forward flow line through  $y$  is contained in

$y \in \cup_{0 \leq t' \leq t} B(x_0, -t', \text{rexp}(2t))$  and, because of the assumption that the closure of  $\mathcal{B}$  is compact, exists for all  $t \leq \tau - t'$ . Hence, for  $t' \leq \tau$  and for any  $r \leq .9$  we have

$$B(x_0, -t', r) \subset P(x_0, 0, (1.01)r, -t') \quad \text{and} \quad P(x_0, -t', r, t') \subset \mathcal{B}((1.01)r). \quad (3.1)$$

In particular we have  $P(x_0, -t, .9, t) \subset \mathcal{B}$  for all  $0 \leq t \leq \tau$ . This proves the first item.

Now we turn to the second and third items. We continue to assume that  $0 < \tau \leq \ln(\sqrt{1.01})$  and we first establish these items under the following:

**Stronger volume hypothesis:** Suppose that for all  $t \in [0, \tau]$  the volume of  $B(x_0, -t, .9)$  is bounded below by some  $w' > 0$ .

We shall show that there are positive constants  $B, C$  satisfying the inequality in the second item, where the constants  $B$  and  $C$  are allowed to depend on  $w'$ . Having fixed  $w' > 0$  and assuming the stronger volume hypothesis, suppose that the second item does not hold for any constants  $B, C$ . We take sequences  $0 < B_n, C_n$  with  $B_n + C_n$  tending to  $\infty$  as  $n \rightarrow \infty$  and examples  $\mathcal{B}_n$  centered at points  $x_n$ , defined for  $-\tau_n \leq t \leq 0$  with  $0 < \tau_n \leq \ln(\sqrt{1.01})$  satisfying the stronger volume hypothesis and with  $Rm \geq -1$  on  $\mathcal{B}_n$ . We set  $A_n = \alpha \sqrt{\min(B_n, C_n)}$ , for a constant  $\alpha > 0$  to be determined. For each  $n$  we have a point  $(y_n, -t_n) \in B(x_n, -t_n, 1/3)$  with  $Q_n = R(y_n, -t_n) \geq B_n + C_n/(\tau_n - t_n)$ , and hence  $Q_n \rightarrow \infty$  as  $n \rightarrow \infty$ .

**Claim 3.14.** *Provided that  $\alpha > 0$  is sufficiently small, for every  $n$  sufficiently large there is a point  $(y'_n, -t'_n) \in \mathcal{B}_n$  satisfying  $t_n \leq t'_n \leq t_n + A_n Q_n^{-1}$  and  $d_{-t'_n}(x_n, y'_n) < .4$  with the following properties:*

1. Setting  $Q'_n = R(y'_n, t'_n)$ , we have  $Q'_n \geq B_n + C_n/(\tau_n - t'_n)$ .
2.  $R(z, -t) < 2Q'_n$  for every  $(z, -t)$  with  $t'_n \leq t \leq t'_n + A_n(Q'_n)^{-1}$  and with

$$d_{-t}(x_n, z) \leq d_{-t'_n}(x_n, y'_n) + A_n(Q'_n)^{-1/2}.$$

*Proof.* We begin with  $(y_n^0, -t_n^0)$  with  $d_{-t_n^0}(y_n^0, x_n^0) < 1/3$  and  $Q_n^0 = R(y_n^0, -t_n^0) \geq B_n + C_n/(\tau_n - t_n^0)$ . If this point does not satisfy the second conclusion, then there is  $(y_n^1, -t_n^1)$  with  $t_n^0 \leq t_n^1 \leq t_n^0 + A_n(Q_n^0)^{-1}$  and  $d_{-t_n^1}(x_n, y_n^1) \leq d_{-t_n^0}(y_n^0, x_n) + A_n(Q_n^0)^{-1/2}$  with  $Q_n^1 = R(y_n^1, t_n^1) \geq 2Q_n^0$ . Direct computation shows that, provided that  $C_n$  is at least 4, we have  $R(y_n^1, -t_n^1) \geq B_n + C_n/(\tau_n - t_n^1)$ . We repeat this argument using  $(y_n^1, -t_n^1)$  instead of  $(y_n^0, -t_n^0)$  constructing counter-example points  $(y_n^k, -t_n^k)$  with  $Q_n^k \geq 2^k Q_n^0$ . Because of the geometric increase in  $Q_n^k$  we see that  $d_{-t_n^k}(y_n^k, x_n) < 1/3 + A_n(Q_n^0)^{-1/2}(1 + (1/2)^{1/2} + (1/2) + (1/2)^{3/2} + \dots) < 1/3 + A_n(Q_n^0)^{-1/2}/(1 - \sqrt{1/2})$ . Since  $A_n \leq \alpha \sqrt{Q_n^0}$ , provided that  $\alpha > 0$  is chosen sufficiently small, we see that  $d_{-t_n^k}(x_n, y_n^k) < .4$  for all  $k$  for which the construction can be performed. But now by compactness, the process must terminate in a finite number of steps. At the last step we have a point as required.  $\square$

Once we have  $(y'_n, -t'_n)$  as in the previous claim, we shift  $-t'_n$  to zero, rescale by  $Q'_n$  to get a sequence of Ricci flows centered at points  $(y'_n, 0)$ . We have curvature

bounds  $Rm \geq -(Q'_n)^{-1}$  and  $R < 2$ , and hence sectional curvature bounded independent of  $n$ , on the union  $U_n$  of balls  $B(y'_n, -t, A_n)$  for all  $-A_n \leq t \leq 0$ . Also  $R(y'_n, 0) = 1$ .

**Claim 3.15.** *Using the shifted and rescaled time and the rescaled metric,  $U_n$  contains the entire parabolic neighborhood  $P(y'_n, 0, A_n/2, -A_n/16)$ ; and in particular the sectional curvature on  $P(y'_n, 0, A_n/2, -A_n/16)$  is bounded independent of  $n$ .*

*Proof.* From the construction we have  $R < 2$  on  $U_n$ . Since  $Rm \geq -(Q'_n)^{-1}$ , it follows that  $Ric < 3$  on  $U_n$ . Invoking Proposition 3.21 of [21] with  $r_0 = 1$ , we see that the derivative of  $d_{-t}(y'_n, z)$  with respect to time is  $\geq -8$ . Thus, for any  $-A_n/16 \leq -t < -t' \leq 0$  we have  $d_{-t}(y'_n, z) \leq d_{-t'}(y'_n, z) + A_n/2$  as long as the forward flow line from  $(z, -t)$  to time  $-t'$  remains in  $U_n$ . The result follows easily.  $\square$

Now let us return for a moment to the  $\mathcal{B}_n$  with the original metric and time coordinates. By our stronger volume hypothesis and volume comparison (using the fact that  $Rm \geq -1$ ), there is a constant  $w'' > 0$  depending only on  $w'$  such that  $\text{Vol} B(x'_n, -t_n, .1) \geq w''$ . Since  $B(x'_n, -t'_n, .1) \subset B(y'_n, -t'_n, .5)$ , it follows that  $\text{Vol} B(y'_n, -t'_n, .5) \geq w''$ . Again by volume comparison, using the fact that  $B(y'_n, -t'_n, .5) \subset B(x_n, -t'_n, .9)$  and the curvature estimate  $Rm \geq -1$  on  $B(x_n, -t'_n, .9)$ , there is  $\nu > 0$ , depending only on  $w''$ , and hence only on  $w'$ , such that for any  $r \leq .5$  we have  $\text{Vol} B(y'_n, -t'_n, r) \geq \nu r^3$ .

It follows from these results for the  $\mathcal{B}_n$ , in the rescaled flow  $U_n$  we have  $\text{Vol} B(y'_n, 0, r) \geq \nu r^3$  for all  $r \leq A_n/2$ . The uniform curvature control on the entire parabolic neighborhood  $P(y'_n, 0, A_n/2, -A_n/16) \subset U_n$  and uniform volume lower bound on  $B(y'_n, 0, A_n/2) \subset U_n$  implies that, after passing to a subsequence, there is a geometric limit of the  $(U_n, (y'_n, 0))$  which is a complete, ancient solution of bounded, non-negative curvature with scalar curvature at the base point (the limit of the  $(y'_n, 0)$ ) equal to 1. We claim that this is a contradiction. The reason is that since  $\text{Vol} B(y'_n, 0, A_n/2) \geq \nu(A_n/2)^3$  for every  $n$  and since  $A_n \rightarrow \infty$  as  $n \rightarrow \infty$ , this implies that the asymptotic volume of the 0 time-slice of the limit is  $\geq \nu$ , and hence using the fact that the limit has non-negative curvature, the solution is a  $\kappa$ -solution for  $\kappa = \nu$  equal. But this contradicts the fact (see Theorem 9.59 of [21]) that the asymptotic volume of a  $\kappa$ -solution is zero. This contradiction establishes the inequality in the second item under the stronger volume hypothesis.

Now continuing to work with the stronger volume hypothesis, we show that there is a  $0 < \hat{\tau}_0 \leq \ln(\sqrt{1.01})$  such that the third item holds as long as  $0 < \tau \leq \hat{\tau}_0$ . First of all, because of the bound on  $R$  just established and the fact that  $Rm \geq -1$ , we see that  $Ric(x, -t) \leq (\hat{B} + C/(\tau - t))$  for all  $0 \leq t \leq \tau$  and all  $(x, -t) \in B(x_0, -t, r_0/3)$ , where  $\hat{B} = B + 1$ . By Proposition 3.21 in [21] for any  $(x, -t) \in B(x_0, -t, r_0/3)$  with  $0 \leq t \leq \tau$  we have the derivative of the function  $d_s(x_0, x)$  at  $s = -t$  is at least  $-8\sqrt{\hat{B} + C/(\tau - t)}/\sqrt{3}$ . It now follows by integrating that

$$d_0(x_0, x) \geq d_{-t}(x, x_0) - \left(8\sqrt{\hat{B}t} + (8\sqrt{C})\sqrt{t}\right) / \sqrt{3}.$$

Thus, assuming that  $\hat{\tau}_0 > 0$  is sufficiently small, how small depending only on  $B$  and  $C$ , and that  $\tau \leq \hat{\tau}_0$ , we see that  $d_0(x_0, x) \geq d_{-t}(x_0, x) - 1/100$  for all  $0 \leq t \leq \tau$  and all  $(x, -t) \in B(x_0, -t, 1/3 - 1/100)$ . It follows that for all  $r \leq 1/3 - 1/100$  and all  $0 \leq t \leq \tau$  we have

$$P(x_0, 0, r, -\tau) \subset \mathcal{B}(r + 1/100).$$

For  $1/100 \leq r \leq 1/3 - 1/100$ , the inclusion  $B(x_0, -t, r - 1/100) \subset P(x_0, 0, r, -t)$  follows from Equation (3.1). This completes the proof of the third item under the stronger volume hypothesis

To complete the proof of the second and third items we must show that there is a  $0 < \tau_0 \leq \hat{\tau}_0 \leq \ln(\sqrt{1.01})$  such that under the hypothesis of the lemma the stronger volume hypothesis holds provided  $\tau \leq \tau_0$ ; that is to say, that there is a  $w' > 0$  depending only on  $w$  such that  $\text{Vol } B(x_0, -t, .9) \geq w'$  for all  $t \in [0, \tau]$ . We denote by  $V_{\text{hyp}}(r)$  the volume of the ball of radius  $r$  in hyperbolic 3-space. First notice that if  $\text{Vol } B(x_0, 0, .95) \geq w$  then by the Bishop-Gromov volume comparison we have  $\text{Vol } B(x_0, 0, r) \geq w(r) = wV_{\text{hyp}}(r)/V_{\text{hyp}}(.95)$  for every  $r < .95$ . We set  $w' = w(1/16)/2$ , and we consider the maximal  $\tau'$  such that  $\text{Vol } B(x_0, -t, .9) \geq w'$  for all  $t \in [0, \tau']$ . We conclude the proof in this case by showing that there is  $0 < \tau_0 \leq \hat{\tau}_0$  such that if  $\tau \leq \tau_0$  then  $\tau' = \tau$ . By what we have just established there are constants  $B'$  and  $C'$  depending only on  $w'$ , and hence depending only on  $w$ , such that  $R(x, -t) \leq B' + C'/(t' - t)$  on  $\cup_{0 \leq t \leq \tau'} B(x_0, -t, 1/3)$ , and there is  $\hat{\tau}'_0 > 0$ , depending only on  $B'$  and  $C'$ , and hence depending only on  $w$ , such that if  $\tau' \leq \hat{\tau}'_0$  then image of the ball  $B(x_0, -t, 1/8)$  under the Ricci flow from time  $-t$ , with  $0 < t \leq \hat{\tau}'_0$  to time 0 includes the ball  $B(x_0, 0, 1/16)$ . This implies that the volume of the result  $\hat{B}$  of flowing  $B(x_0, -t, 1/8)$  to time 0 is at least  $w(1/16)$ . On the other hand,  $dV/dt = -\int R dV \leq -R_{\min} V$ . Since  $Rm \geq -1$ , we see that  $R_{\min} \geq -6$ . Thus,  $\text{Vol } \hat{B} \leq \text{Vol } B(x_0, -t, 1/8) \exp(6t)$ . Taking  $\tau_0 = \min(\hat{\tau}'_0, \ln 2/12)$ , we conclude that

$$\text{Vol } B(x_0, -t, 1/8) > \text{Vol } \hat{B}/2 \geq w(1/16)/2.$$

A fortiori, we have  $\text{Vol } B(x_0, -t, .9) > w(1/16)/2$  under the same assumption. This means that it must be the case that  $\tau' = \tau$ . This shows the volume estimate  $\text{Vol } B(x_0, -t, .9) \geq w(1/16)/2$  holds for all  $t \in [0, \tau]$  provided that  $\tau \leq \tau_0$ . This completes the proof that the hypotheses of the lemma imply the stronger volume hypothesis for all  $0 \leq t \leq \tau$  provided that  $\tau \leq \tau_0$ . This completes the proof of Parts 2 and 3 of the lemma.

In the course of proving the second and third parts of the proposition, we showed that  $\text{Vol } B(x_0, -t, 1/8) > w(1/16)/2$  for every  $0 \leq t \leq \tau$ . Hence,  $\text{Vol } B(x_0, -\tau, 1/4) > w(1/16)/2$ . This establishes the last item and completes the proof of Lemma 3.13.  $\square$

**Corollary 3.16.** (a) *Given  $w > 0$  there are  $\tau_0 > 0$ ,  $w' > 0$ , and  $K_0 < \infty$  such that the following holds for any  $0 < \tau \leq \tau_0$ . Suppose that we have a generalized Ricci flow on  $\mathcal{U}$  and suppose  $\mathcal{B} = \sup_{0 \leq t \leq \tau} B(x_0, -t, .95)$  has compact closure in  $\mathcal{U}$ . Suppose that the sectional curvatures are bounded below by  $-1$  on  $\mathcal{B}$  and suppose that the volume of the ball  $B(x_0, 0, .95)$  is at least  $w$ , then:*

1.  $R(x, t) \leq K_0 \tau^{-1}$  for all  $(x, t) \in P(x_0, 0, 1/4, -\tau/2)$ , and
2. the ball  $B(x_0, -\tau, 1/4)$  has volume at least  $w'$ .

(b) Suppose the entire parabolic neighborhood  $P(1) = P(x_0, 0, 1, -\tau)$  is contained in  $\mathcal{U}$  with compact closure. Suppose that  $Rm \geq -1$  on  $P$  and the volume of  $B(x_0, 0, 1) \geq w$ . There are constants  $\tilde{\tau}_0 > 0$ ,  $\tilde{K}_0 < \infty$  and  $\tilde{w}' > 0$  such that the above two conclusions hold with  $\tilde{\tau}_0$  replacing  $\tau_0$ , with  $\tilde{K}_0$  replacing  $K_0$  and with  $\tilde{w}' > 0$  replacing  $w'$ .

*Proof.* Given  $w > 0$  we take  $\tau_0$  and  $w' > 0$  as in Lemma 3.13, and we set  $K_0 = B\tau_0 + 2C$ , where  $B$  and  $C$  are the constants from this lemma. Set  $P(1/4) = P(x_0, 0, 1/4, -\tau)$ . By Lemma 3.13 we have  $P(1/4) \subset \cup_{0 \leq t \leq \tau} B(x_0, -t, 1/3)$  and  $R(x, -t) \leq B + C/(\tau - t)$  for all  $(x, -t) \in P(1/4)$ . Thus,  $R < K_0 \tau^{-1}$  on the parabolic neighborhood  $P(x_0, 0, 1/4, -\tau/2)$ . Also, this lemma also implies that  $\text{Vol } B(x_0, -\tau, 1/4) \geq w'$ . This proves all the statements in Part (a) on the corollary.

As Part (b), since  $Rm \geq -1$  on  $P(1)$  and since  $\tau \leq \ln(\sqrt{1.01})$ , we have

$$B(x_0, -t, .95) \subset P(1)$$

for every  $0 \leq t \leq \tau$ . Also, there is  $\tilde{w} > 0$  depending only on  $w$  such that  $\text{Vol } B(x_0, 0, .95) \geq \tilde{w}$ . We simply apply what we have already established with  $w$  replaced by  $\tilde{w}$ .  $\square$

The second lemma shows the existence of a sub-ball with a Euclidean volume estimate for all further sub-balls.

**Lemma 3.17.** *Fix  $n > 0$ . For any  $w > 0$  there is  $\theta_0 = \theta_0(w) > 0$  such that if  $B(x, 1)$  is a metric ball of volume at least  $w$  compactly contained in a  $n$ -manifold without boundary with sectional curvatures at least  $-1$ , then there exists a ball  $B(y, \theta_0) \subset B(x, 1)$  such that every sub-ball  $B(z, r) \subset B(y, \theta_0)$  of any radius  $r$  has volume at least  $(1 - \epsilon)$  times the volume of the  $n$ -dimensional Euclidean ball of the same radius.*

*Proof.* Recall that a  $(k, \delta)$  strainer centered at a point  $x$  in an Alexandrov space consists of  $a_1, \dots, a_k, b_1, \dots, b_k$  such that for all  $1 \leq i, j \leq k$  the comparison angles satisfy:

$$\begin{aligned} \tilde{\angle} a_i x a_j &> \pi/2 - \delta \quad \text{for all } i \neq j \\ \tilde{\angle} a_i x b_i &> \pi - \delta \\ \tilde{\angle} a_i x b_j &> \pi/2 - \delta; \quad \text{for all } i \neq j \\ \tilde{\angle} b_i x b_j &> \pi/2 - \delta \quad \text{for all } i \neq j. \end{aligned}$$

The size of the strainer is the minimum of the  $2k$  lengths  $|xa_i|, |xb_i|$ . In an  $n$ -dimensional Alexandrov space  $X$ , for any  $\delta > 0$  the set of points with an  $(n, \delta)$ -strainer is an open dense set. Furthermore, for  $\delta > 0$  sufficiently small, if  $y \in X$  has an  $(n, \delta)$  strainer of size  $s$ , then there exist a constant  $r = r(s) > 0$  and a  $(1 - \epsilon)$  bilipschitz homeomorphism from  $B(y, r)$  the ball of radius  $r$  in Euclidean  $n$ -space. For more details on all these facts see [3].

Now fix  $w > 0$  and suppose that the result does not hold. Take a sequence  $\theta_k \rightarrow 0$  as  $k \rightarrow \infty$  and balls  $B(x_k, 1)$  that do not satisfy the conclusion for  $\theta = \theta_k$ . Pass to a subsequence and take a limit  $X$  of the  $B(x_k, 1)$ . Because of the uniform positive lower bound for the volumes of the  $B(x_k, 1)$ , the limit  $X$  is an  $n$ -dimensional Alexandrov ball of radius 1. As such it contains a point  $y$  with a  $(n, \delta)$ -strainer (with  $\delta > 0$  as above) of size  $s > 0$ . This strainer defines a  $(1 - \epsilon)$  bilipschitz map from a smaller ball  $B(y, r(s))$  centered at  $y$  to the corresponding ball in  $\mathbb{R}^n$ . This then implies that the result holds for the  $B(x_k, 1)$  for a sequence of points  $y_k$  converging to  $y$  and radius  $r(s)$ . This is a contradiction.  $\square$

**Definition 3.18.** We say that a ball  $B(x, t, r)$  in a 3-manifold has a *Euclidean volume estimate* if for every sub-ball  $B' = B(y, t, s) \subset B(x, t, r)$ , the volume of  $B'$  is at least  $(1 - \epsilon)$  times the volume of a Euclidean 3-ball of the same radius.

With these results in place we return to the proof of Proposition 3.11 in the case  $r(t_0) < r_0$ . We fix  $\tau_0 > 0$ ,  $w' > 0$ , and  $K_0 < \infty$  as in Corollary 3.16 for  $w = (1 - \epsilon)$  times the volume of a Euclidean ball of radius 1. (Recall that  $\epsilon > 0$  is one of the canonical neighborhood parameters.) We set  $\theta_0 = \min(\theta_0(4^3 w')/4, 1/20)$  where  $\theta_0(w')$  is the constant from Lemma 3.17. We shall show that there is  $\bar{r}_1 > 0$  such that the conclusion of Proposition 3.11 holds provided that  $r(t_0) < r_0 \leq \bar{r}_1 \sqrt{t_0}$  for  $\tau = \tau_0/2$  and for  $K = 2K_0 \tau_0^{-1}$ . Suppose that there is no  $\bar{r}_1 > 0$  as required. Then we take a sequence of  $\bar{r}_{1,n} \rightarrow 0$  and counter-examples consisting of Ricci flows with surgery  $(\mathcal{M}_n, G_n)$  containing balls  $B(x_n, t_n, r_n)$  with  $r(t_n) < r_n \leq \bar{r}_{1,n} \sqrt{t_n}$  for the given values of  $\tau$  and  $K$ . Since  $\bar{r}_{1,n} \rightarrow 0$  as  $n \rightarrow \infty$  and since  $\bar{r}_{1,n} > r(t_n)$ , it follows that  $t_n \rightarrow \infty$  as  $n \rightarrow \infty$ . It follows from the curvature pinching assumption and the fact that  $r_n^2 t_n^{-1} \leq \bar{r}_{1,n}^2 \rightarrow 0$  as  $n \rightarrow \infty$ , that given any constant  $1 \leq D < \infty$ , for all  $n$  sufficiently large if  $R(x, t) < Dr_n^{-2}$  for some  $t \geq r_n^2/4\bar{r}_{1,n}^2$ , then  $Rm(x, t) \geq -r_n^{-2}$  and consequently  $|Rm(x, t)| \leq Dr_n^{-2}$ .

Clearly, the hypotheses of the lemma and the negation of the conclusion of the lemma are closed under taking limits of  $(x, t, r)$ . Thus, we can choose  $t_n$  to be the first time where the lemma fails to hold for the constant  $\bar{r}_{1,n}$  and we choose  $r_n$  to be the minimal radius of a counter-example ball at time  $t_n$ . Our goal is to show that

$$\mathcal{B}_n = \cup_{0 \leq t \leq \tau_0 r_n^2} B(x_n, t_n - t, r_n),$$

has closure in the smooth part of the Ricci flow with surgery (which is a generalized Ricci flow), and  $Rm \geq -r_n^{-2}$  on  $\mathcal{B}_n$ . If we can establish this, then  $\mathcal{B}_n$  has compact closure in the generalized Ricci flow which is the smooth part of the Ricci flow with surgery. Rescaling by  $r_n^{-2}$ , applying the first part of Corollary 3.16 to the rescaled flow and then rescaling the conclusion of this result by  $r_n^2$ , we have  $R < K_0 \tau_0^{-1} r_n^{-2}$  on  $P(x_n, t_n, r_n/4, -\tau_0 r_n^2/2)$  as required by Proposition 3.11.

We shall establish that  $\mathcal{B}_n$  has closure contained in the smooth part of the Ricci flow with surgery and that  $Rm \geq -r_n^{-2}$  on  $\mathcal{B}_n$  on segments of time, moving backward one segment at a time. We shall find  $\Delta t > 0$  depending only on  $\epsilon, B, C$ , and  $\tau_0$ , and we shall show by induction on  $N$  that, as long as  $(N - 1)\Delta t \leq \tau_0$ , the Ricci flow with surgery satisfies  $Rm \geq -r_n^{-2}$  on  $\mathcal{B}_{n,N}$ , which is the intersection of  $\mathcal{B}_n$  with the pre-image in the Ricci flow with surgery of the time interval  $[t_n - N(\Delta t)r_n^2, t_n]$ .

Let us consider the first step in the induction. The appropriate value of  $\Delta t$  will emerge in the course of this argument. By Part (2) of Corollary 3.16

$$\text{Vol } B(x_n, t_n, r_n/4) \geq w' r_n^3.$$

By Lemma 3.17 there is a ball  $B(y_n, t_n, \theta_0 r_n) \subset B(x_n, t_n, r_n/4)$  that has a Euclidean volume estimate. Since  $t_n \rightarrow \infty$  as  $n \rightarrow \infty$  and since  $r_n > r(t_n) \geq \delta^{-2}(t_n) \hat{h}(t_n)$ , for all  $n$  sufficiently large,  $\theta_0 r_n > 4C\hat{h}(t_n)$ . Since  $B(y_n, t_n, \theta_0 r_n) \subset B(x_n, t_n, r_n)$  we have  $Rm \geq -r_n^{-2}$  on this ball. Since  $\theta_0 r_n < r_n$ , the fact that we chose  $r_n$  to be the minimal radius of a counter-example of Proposition 3.11 at time  $t_n$  implies that the proposition holds for  $B(y_n, t_n, \theta_0 r_n)$ . We have just checked that all the hypotheses of this proposition hold for  $B(y_n, t_n, \theta_0 r_n)$ . Thus, the proposition implies that the Ricci flow with surgery  $(\mathcal{M}_n, G_n)$  contains the entire parabolic neighborhood  $P(y_n, t_n, \theta_0 r_n/4, -\tau_0 \theta_0^2 r_n^2)$  and satisfies  $R < K\theta_0^{-2} r_n^{-2}$  on this parabolic neighborhood. By the curvature pinching result, for all  $n$  sufficiently large we have  $|Rm| \leq K\theta_0^{-2} r_n^{-2}$  and  $Rm \geq -r_n^{-2}$  on this parabolic neighborhood. We set

$$\alpha = \min(\sqrt{\tau_0}, K^{-1/2}, 1/4)\theta_0.$$

Then  $|Rm| \leq \alpha^{-2} r_n^{-2}$  on  $P_n(\alpha) = P(y_n, t_n, \alpha r_n, -\alpha^2 r_n^2)$ . The bound on  $|Rm|$  on  $P_n(\alpha)$  implies that for any  $t' \in [t_n - \alpha^2 r_n^2/2, t_n]$  we have  $P(y_n, t', \alpha r_n/4, -\alpha^2 r_n^2/16) \subset P_n(\alpha)$  and thus  $|Rm| \leq \alpha^{-2} r_n^{-2}$  on  $P(y_n, t', \alpha r_n/4, -\alpha^2 r_n^2/16)$  for all  $t' \in [t_n - \alpha^2 r_n^2, t_n]$ . Recall that  $r_n \leq \bar{r}_{1,n} \sqrt{t_n}$  with  $\bar{r}_{1,n} \rightarrow 0$  as  $n \rightarrow \infty$ . Thus, by Part 3 of Proposition 3.1, there is a constant  $K' < \infty$  such that  $|Rm| \leq K' \alpha^{-2} r_n^{-2}$  on  $B(y_n, t', 4r_n)$  for all  $n$  sufficiently large. Since  $d_{t_n}(x_n, x) < r_n$ , it follows that  $d_{t'}(x_n, y_n) < 2r_n$  for all  $t' \in [t_n - \alpha^2 r_n^2/4K', t_n]$ . Hence, for all such  $t'$  we have  $B(x_n, t', r_n) \subset B(y_n, t', 4r_n)$  and consequently,  $|Rm|$  is bounded by  $K' \alpha^2 r_n^2$  on the union of these balls. Since  $r_n \geq r(t_n)$  which in turn is at least  $\bar{\delta}(t_n)^{-2} \hat{h}(t_n)$  and since  $t_n \rightarrow \infty$  as  $n \rightarrow \infty$ , for  $n$  sufficiently large, the closure of  $B(x_n, t', r_n)$  is disjoint from the surgery caps for all  $t' \in [t_n - \alpha^2 r_n^2/2, t_n]$ . Also, by curvature pinching we conclude that provided that  $n$  is sufficiently large, we have  $Rm \geq -r_n^{-2}$  for every  $t' \in [t_n - \alpha^2 r_n^2/2, t_n]$ . We set  $\Delta t = \alpha^2/4K'$ . We have  $Rm \geq -r_n^{-2}$  on  $\mathcal{B}_{n,1}$ . This is the initial step in the induction.

Suppose that inductively for some  $N \geq 1$  with  $N\Delta t < \tau_0$ , we have shown that  $Rm \geq -r_n^{-2}$  on  $\mathcal{B}_{n,N}$  which is the intersection of  $\mathcal{B}_n$  with the time interval  $[t_n - N(\Delta t)r_n^2, t_n]$ . Then, by Part (2) of Corollary 3.16 we see that the volume of the ball  $B(x_n, t_n - N(\Delta t)r_n^2, r_n/4)$  is at least  $w' r_n^3$ . Hence, by Lemma 3.17 there is a ball  $B(y'_n, t_n - N(\Delta t)r_n^2, \theta_0 r_n) \subset B(x_n, t_n - N(\Delta t)r_n^2, r_n/4)$  that has a Euclidean volume estimate. Once again the fact that  $r_n > r(t_n)$  and  $t_n \rightarrow \infty$  as  $n \rightarrow \infty$  implies that for all  $n$  sufficiently large  $\theta_0 r_n > 4C\hat{h}(t_n)$ . The inductive hypothesis imply that  $Rm \geq -r_n^{-2}$  on  $B(y'_n, t_n - N(\Delta t)r_n^2, \theta_0 r_n)$ . This shows that the hypotheses of the proposition hold for  $B(y'_n, t_n - N(\Delta t)r_n^2, \theta_0 r_n)$ . Since  $t_n - N(\Delta t)r_n^2 < t_n$  by our assumption that  $t_n$  was the first counter-example time for  $\bar{r}_{1,n}$ , the proposition applies to  $B(y'_n, t_n - N(\Delta t)r_n^2, \theta_0 r_n)$  to show that the Ricci flow contains the entire parabolic neighborhood  $P(y'_n, t_n - N(\Delta t)r_n^2, \theta_0 r_n/4, -\tau_0 \theta_0^{-2} r_n^{-2})$  and satisfies  $R < K\theta_0^{-2} r_n^{-2}$  on this parabolic neighborhood. Arguing exactly as in the first step of the

induction, we see that for all  $n$  sufficiently large we have  $Rm \geq -r_n^{-2}$  on

$$\cup_{t_n - (N+1)(\Delta t)r_n^2 \leq t \leq t_n - N(\Delta t)r_n^2} B(x_n, -t, r_n),$$

and that the closure of this union is disjoint from all the surgery caps. Putting this together with what we have already established by induction, gives us the fact that  $Rm \geq -r_n^{-2}$  on  $\mathcal{B}_{n,N+1}$  and the closure of this neighborhood is disjoint from all the surgery caps. We continue this way until we have the result on  $\mathcal{B}_{n,N_0}$  where  $(N_0 - 1)\Delta t \leq \tau_0 < N_0\Delta t$ . Truncating to the time interval  $[-\tau_0, 0]$  we conclude that  $Rm \geq -r_n^{-2}$  on  $\mathcal{B}_n$  and the closure of  $\mathcal{B}_n$  is disjoint from the surgery caps. Hence,  $\mathcal{B}_n$  is contained with compact closure in the generalized Ricci flow that is the smooth part of the Ricci flow with surgery. Invoking Corollary 3.16 we see that  $R(x, t) < K_0\tau^{-1}$  for all  $(x, t) \in P(x_n, t_n, r_n/4, -\tau_0 r_n^2/2)$ . Since we have taken  $K = 2K_0\tau_0^{-1}$ , and  $\tau = \tau_0/2$ , this completes the proof of Proposition 3.11 in the second case, and hence completes the proof of the proposition.

### 3.2.3 A corollary

Now we can derive an important corollary of Proposition 3.11, which is Corollary 6.8 of [27].

**Corollary 3.19.** *For any  $w > 0$  there exist  $\tau' = \tau'(w) > 0$ ,  $K' = K'(w) < \infty$ ,  $\bar{r}' = \bar{r}'(w) > 0$  and  $\theta = \theta(w) > 0$  such that the following holds for any Ricci flow with surgery  $(\mathcal{M}, G)$ . Let  $t_0, r_0$  satisfy  $\theta^{-1}(w)\hat{h}(t_0) \leq r_0 \leq \bar{r}'\sqrt{t_0}$  and assume that there is a ball  $B(x_0, t_0, r_0) \subset \mathcal{M}$  on which the sectional curvatures are bounded below by  $-r_0^{-2}$  and suppose that the volume of  $B(x_0, t_0, r_0)$  is at least  $w r_0^3$ . Then the Ricci flow with surgery is defined in  $P = P(x_0, t_0, r_0/8, -\tau' r_0^2)$  and satisfies  $R(x, t) < K' r_0^{-2}$  for all  $(x, t) \in P$ .*

*Proof.* Fix  $\theta_0 = \min(\theta(w), 1)$  from Lemma 3.17. According to Lemma 3.17 there is a ball  $B(y, t_0, \theta_0 r_0) \subset B(x_0, t_0, r_0)$  that has a Euclidean volume estimate, and of course  $Rm \geq -r_0^{-2}$  on this ball. We shall take  $\theta \leq \theta_0/4C$  so that the condition  $\theta^{-1}\hat{h}(t_0) \leq r_0$  implies that  $4C\hat{h}(t_0) \leq \theta_0 r_0$ . Thus, assuming that  $r_0 \leq \bar{r}'\sqrt{t_0}$  with  $\bar{r}' \leq \bar{r}_1$  from Proposition 3.11, we can apply Proposition 3.11 to  $B(y, t_0, \theta_0 r_0)$  and conclude that the Ricci flow with surgery contains the entire parabolic neighborhood  $P(y, t_0, \theta_0 r_0/4, -\tau\theta_0^2 r_0^2)$  and satisfies  $R < K\theta_0^{-2} r_0^{-2}$  on this parabolic neighborhood. Since  $Rm \geq -r_0^{-2}$ , we have  $|Rm| \leq K\theta_0^{-2} r_0^{-2}$  on this parabolic neighborhood. Thus,  $|Rm| \leq \alpha^{-2} r_0^{-2}$  on  $P = P(y, t_0, \alpha r_0, -\alpha^2 r_0^2)$  where

$$\alpha = \min(1/4, \sqrt{\tau}, K^{-1/2})\theta_0,$$

so that  $\alpha$  depends only on  $w$ . Of course, for each  $t \in [t_0 - (\alpha r_0/2)^2, t_0]$  we have  $P(y, t, \alpha r_0/2, -(\alpha r_0/2)^2) \subset P$ . Now we take  $\bar{r}' = \min(\bar{r}_1, 2\alpha^{-1}\bar{r}(8/\alpha))$  where  $\bar{r}(8/\alpha)$  is the constant in Proposition 3.1. We can apply Proposition 3.1 and conclude that, provided that  $(4C/\theta_0)\hat{h}(t_0) \leq r_0 \leq \bar{r}'\sqrt{t_0}$  so that  $\alpha r_0/2 \leq \bar{r}(8/\alpha)$ , we have a bound on the scalar curvature  $R < K_2(\alpha r_0/2)^{-2}$  on  $\mathcal{B} = \cup_{t_0 - (\alpha r_0/2)^2 \leq t \leq t_0} B(y, t, 4r_0)$  for some  $K_2$  that depends only on  $\alpha$ , and hence only on  $w$ . By curvature pinching,



and the fact that  $r_0^{-2}t_0^{-1} \geq (\bar{r}')^{-2}$ , assuming that  $\bar{r}'$  is sufficiently small, it follows that  $|Rm| < K_2(\alpha r_0/2)^{-2}$  on  $\mathcal{B}$ . Now we set  $\theta(w) = \min(K_2^{-1/2}\alpha/4, \theta_0/4C)$ . Thus, the scalar curvature on  $\mathcal{B}$  is at most  $\theta(w)^{-2}r_0^{-2}/4$ . Since  $\theta^{-1}\hat{h}(t_0) \leq r_0$ , we have  $(4C/\theta_0)\hat{h}(t_0) \leq r_0$ . Thus, we see that the scalar curvature on  $\mathcal{B}$  is at most  $\hat{h}(t_0)/4$ , and hence the closure of  $\mathcal{B}$  is disjoint from the surgery caps, i.e., the closure of  $\mathcal{B}$  is contained in the smooth part of the Ricci flow. Since  $|Rm| < K_2(\alpha r_0/2)^{-2}$  and since  $d_{t_0}(x_0, y) < r_0$ , the open set  $\mathcal{B}$  contains a parabolic neighborhood of the form  $P(x_0, t_0, r_0, -\tau' r_0^2)$  for some  $\tau'$  depending only on  $K_2(\alpha/2)^{-2}$ , and hence depending only on  $w$ . This establishes the corollary with this value of  $\tau'$  with  $K' = 4K_2\alpha^{-2}$ , and for  $\bar{r}' > 0$  sufficiently small.  $\square$

## 4 Proof of Propositions 2.6, 2.8, and 2.9

### 4.1 Proof of Proposition 2.6

Let us begin by recalling the statement that we shall prove:

**Proposition 2.6.** (a) *Given  $w > 0, r > 0, \xi > 0$  there is  $T = T(w, r, \xi) < \infty$  such that the following holds for any Ricci flow with surgery  $(\mathcal{M}, G)$  satisfying Assumptions 3.9 and 3.10. If, for some  $t_0 \geq T$  and some  $x_0 \in M_{t_0}$ , the ball  $B(x_0, t_0, r\sqrt{t_0})$  has volume at least  $w r^3 t_0^{3/2}$  and sectional curvatures bounded below by  $-r^{-2}t_0^{-1}$ , then*

$$|2t_0 \text{Ric}(x_0, t_0) + g(x_0, t_0)|_{g(t_0)} < \xi. \quad (2.6)$$

(b) *In addition, given  $A < \infty$ , there is  $T' = T'(w, r, \xi, A) \geq T(w, r, \xi)$ , and provided that  $t_0 \geq T'$ , the Ricci flow with surgery contains the entire forward parabolic neighborhood  $P(x_0, t_0, Ar\sqrt{t_0}, Ar^2 t_0)$  and Equation (2.6) holds with  $(x_0, t_0)$  replaced by any  $(x, t)$  in this forward parabolic neighborhood.*

*Proof.* Fix  $w > 0, r > 0$ , and  $\xi > 0$  and suppose that (a) does not hold for these constants. Take a sequence of Ricci flows with surgery  $(\mathcal{M}_n, G_n)$  and points  $(x_n, t_n) \in \mathcal{M}_n$  with  $t_n \rightarrow \infty$  so that the hypotheses of Part (a) hold for each  $B(x_n, t_n, r\sqrt{t_n})$  but the conclusion fails for  $(x_n, t_n)$ . Set  $s = \min(r, \bar{r}'(w))$  and  $r_n = s\sqrt{t_n}$  where  $\bar{r}'(w)$  is the constant given in Corollary 3.19. Since  $\hat{h}(t) \rightarrow 0$  as  $t \rightarrow \infty$ , provided that  $n$  sufficiently large  $r_n > \theta^{-1}(w)\hat{h}(t_n)$ . Passing to a subsequence allows us to assume this holds for all  $n$ . Then the conclusions of Corollary 3.19 hold for  $B(x_n, t_n, r_n)$  for all  $n$ , which means that  $(\mathcal{M}_n, G_n)$  contains the entire parabolic neighborhood  $P(x_n, t_n, r_n/8, -\tau' r_n^2)$  and has scalar curvature bounded by  $K' r_n^{-2}$  on this parabolic neighborhood, where  $\tau' > 0$  and  $K' < \infty$  are the constants depending on  $w$  given in Corollary 3.19. Thus, passing to a subsequence, rescaling space and time by  $t_n^{-1} = r_n^{-2} s^2$ , we have a sequence of parabolic neighborhoods  $\tilde{P}_n = P(x_n, 1, s/8, -\tau' s^2)$  with metrics  $h_n(t) = (1/t_n)g(t_n t)$  on which the scalar curvature is bounded by  $K' s^{-2}$ . Since, by the hypothesis of Part (a) we have  $\text{Vol } B(x_n, t_n, r\sqrt{t_n}) \geq w r^3 t_n^{3/2}$  and  $Rm \geq -r_n^{-2} t_n$ , and since  $s \leq r$ , by volume comparison, there is a  $w' > 0$  depending only on  $w$  such that the volume of  $B(x_n, t_n, r_n/8) \geq w' r_n^3$ . Thus, the final time-slice of  $\tilde{P}_n$  has volume at least

$w's^3$ . Consequently, we can extract a subsequence limiting smoothly to an abstract parabolic neighborhood  $\tilde{P}_\infty = P(x_\infty, 1, s/8, -\tau's^2)$ . By Corollary 2.5 the sectional curvature of the final time-slice of the limit is constant and equal to  $-1/4$ , which means that the Ricci curvature of the  $t = 1$  time-slice of the limit is constant and equal to  $-1/2$ . Since the limiting process is smooth, it follows that Inequality (2.6) holds for all  $n$  sufficiently large.

Now let us establish that given  $A$  in addition to  $w, r$ , and  $\xi$ , provided that  $t_0$  is sufficiently large, Inequality (2.6) holds on  $B(x_0, t_0, Ar_0)$ . By the argument in Part (a) given immediately above, there are constants  $\tau' = \tau'(w) > 0$  and  $s = s(w, r) > 0$  (which we take less than 1) and  $K' = K'(w) < \infty$  such that for all  $t_0$  sufficiently large, the scalar curvature on  $P(x_0, t_0, s\sqrt{t_0}/8, -\tau's^2t_0)$  is bounded by  $K's^{-2}$ . Now take  $s' = \min(s/8, \sqrt{\tau'}s, (K')^{-1/2}s/\sqrt{2})$  so that  $s'$  depends only on  $w$  and  $r$ . Then the scalar curvature on  $P(x_0, t_0, s'\sqrt{t_0}, -(s')^2t_0)$  is bounded by  $(s')^{-2}t_0^{-1}/2$ . Provided that  $t_0$  is sufficiently large (how large depending on  $w$  and  $r$ ), it follows from the curvature pinching hypothesis that  $|Rm| \leq (s')^{-2}t_0^{-1}$  on this parabolic neighborhood. Also, by volume comparison the volume of  $B(x_0, t_0, s'\sqrt{t_0})$  is at least  $(A')^{-1}(s')^3t_0^{3/2}$  for some constant  $A' < \infty$ , depending only on  $w$  and  $r$ . Since we have required  $\bar{\delta}(t) \leq \bar{\delta}'_{2t}(2t)$  from Lemma 3.1, for any  $A' \leq t_0/r$  the conclusion of Lemma 3.1 holds for  $B(x_0, t_0, A'r\sqrt{t_0})$  for some constant  $K'_1 = K'_1(A', w, r)$ . Thus, given any  $A < \infty$ , there is a constant  $K''_1 = K''_1(A, w, r)$  such that, provided that  $t_0$  is sufficiently large, any point of  $B(x_0, t_0, Ar\sqrt{t_0})$  with scalar curvature at least  $K''_1r^{-2}t_0^{-1}$  has a  $(C, \epsilon)$ -canonical neighborhood. Suppose that there is such a point. Then the ball contains a point with scalar curvature exactly  $Q = K''_1r^{-2}t_0^{-1}$ , which also has a canonical neighborhood. We set  $\tilde{r} = r/\sqrt{K''_1}$ . This canonical neighborhood contains the ball of radius  $\tilde{r}\sqrt{t_0} = (K''_1)^{-1/2}r\sqrt{t_0}$ , and the volume of this ball is at least  $\kappa\tilde{r}^3r_0^{3/2}$  since canonical neighborhoods are  $\kappa$  non-collapsed for a universal  $\kappa$ . Also, the sectional curvature on the  $(C, \epsilon)$ -canonical neighborhood is bounded below by  $-\epsilon\tilde{r}^{-2}t_0^{-1}$ . Hence, if  $t_0$  is sufficiently large, how large depending on  $r/\sqrt{K''_1}$  and hence depending only on  $w, r$  and  $A$ , we can apply Part (a) of this result to see that Inequality (2.6) holds. This is of course absurd, since there are 2-planes where the sectional curvature is positive and of the order  $\tilde{r}^{-2}t_0^{-1}$ . This contradiction shows that, provided that  $T$  is sufficiently large, the scalar curvature on  $B(x_0, t_0, Ar\sqrt{t_0})$  is bounded above by  $K''_1r^{-2}t_0^{-1}$  for a constant  $K''_1$  depending on  $A, w, r$ . If  $t_0$  is sufficiently large, then by the curvature pinching assumption  $Rm > -r^{-2}t_0^{-1}$  on this ball.

By Lemma 3.1, there is  $\kappa'$  depending on  $A$  and  $w$  such that every point of  $B(x_0, t_0, Ar\sqrt{t_0})$  is  $\kappa'$ -non-collapsed on scales  $r\sqrt{t_0}$ . In particular, the volume of  $B(x_0, t_0, Ar\sqrt{t_0}) \geq (\kappa'/A^3)r^3t_0^{3/2}$ . Now we can apply just established conclusion in Part (a) of this result once again to see that Inequality (2.6) holds at every point of  $B(x_0, t_0, Ar\sqrt{t_0})$  provided that  $T$  is sufficiently large, how large depending on  $A, w$  and  $r$ .

Now we establish that for  $t_0$  sufficiently large (given  $A, w, r$ , and  $\xi$ ) Inequality (2.6) holds at every point of  $P(x_0, t_0, Ar\sqrt{t_0}, Ar^2t_0)$ . We consider the forward evolution of  $B(x_0, t_0, Ar\sqrt{t_0})$  under Ricci flow on  $[t_0, t_0 + Ar^2t_0]$ . If Inequality (2.6)

does not hold for this entire forward parabolic neighborhood, then there is a first time  $t' > t_0$  where it fails. Of course, at  $t'$  the weak form of Inequality (2.6) holds. By the curvature bound, the slice of the forward parabolic neighborhood at time  $t'$  is contained in  $B(x_0, t', A'r\sqrt{t_0})$  for some  $A'$  that depends only on  $A$ . On the other hand, since the Ricci curvature is controlled on the entire evolution from time  $t_0$  to time  $t' \leq t_0 + Ar^2t_0$ , we see that the ball  $B(x_0, t', r\sqrt{t'})$  has volume bounded below by a constant times  $r^3(t')^{3/2}$  where the constant depends only on  $w$  and  $A$ . Also, the sectional curvatures are bounded below by  $-r'^{-2}(t')^{-1}$ . Hence, we can apply what we just established to see that in fact, provided that  $T$  is sufficiently large, depending on  $A, w, r$ , the result holds for the  $t'$  time-slice of the forward parabolic neighborhood, which is a contradiction.

This completes the proof of Proposition 2.6.  $\square$

## 4.2 Proof of Proposition 2.8

Recall the statement that we shall prove:

**Proposition 2.8.** *For any  $w > 0$  there is  $\bar{\rho} = \bar{\rho}(w) > 0$  such that for all  $t$  sufficiently large (how large depending on  $w$ ) for any Ricci flow with surgery  $(M, G)$  satisfying Assumptions 3.9 and 3.10, and for any  $x \in M_t$ , if  $\rho(x, t) < \bar{\rho}\sqrt{t}$  we have*

$$\text{Vol } B(x, t, \rho(x, t)) < w\rho^3(x, t).$$

*Proof.* Fix  $w > 0$ .

**Case 1: For all  $t$  sufficiently large we have  $\rho(x, t) \leq \theta^{-1}(w)\hat{h}(t)$ .** Since  $\rho(x, t)$  is defined so that the infimum of the sectional curvatures on  $B(x, t, \rho(x, t))$  is  $-\rho(x, t)^{-2}$ , there is a point  $(y, t) \in B(x, t, \rho(x, t))$  with a sectional curvature at  $(y, t)$  less than  $-\rho(x, t)^{-2}/2$ . Since  $\rho(x, t) \leq \theta^{-1}(w)\hat{h}(t)$  and  $\hat{h}(t) \leq \bar{\delta}^2(t)r(t)$  where  $\bar{\delta}(t)$  is a monotone decreasing function of  $t$  with limit 0 as  $t \rightarrow \infty$ , it follows that, given  $K < \infty$ , provided that  $t$  is sufficiently large, there is a sectional curvature at  $(y, t)$  which is less than  $-Kr^{-2}(t)$ . By curvature pinching, again assuming that  $t$  is sufficiently large, we also have  $R(y, t) > Kr^{-2}(t)$ , so that  $(y, t)$  has a  $(C, \epsilon)$ -canonical neighborhood.

**Claim 4.1.** *Let  $Q_0 = Q_0(x, t) = \epsilon^{-2}\rho(x, t)^{-2}/16$ . Then, provided that  $t$  is sufficiently large, every point of  $B(x, t, \rho(x, t))$  has scalar curvature  $> Q_0$  and has a  $(C, \epsilon)$ -canonical neighborhood.*

*Proof.* By the discussion at above, provided that  $t$  is sufficiently large, there is a point  $(y, t) \in B(x, t, \rho(x, t))$  with  $R(y, t) \geq 2Q_0$ . Suppose that the claim does not hold. Then, there is a point  $(z, t) \in B(x, t, \rho(x, t))$  with  $R(z, t) = Q_0$ . Since, for all  $t$  sufficiently large  $\rho(x, t) \leq \theta^{-1}(w)\hat{h}(t) < r(t)$ , it follows that if  $t$  is sufficiently large then  $(z, t)$  has a  $(C, \epsilon)$ -canonical neighborhood. This canonical neighborhood contains the ball of radius  $\epsilon^{-1}/2\sqrt{Q_0} > 2\rho(x, t)$  centered at  $(z, t)$ . Since  $(z, t) \in B(x, t, \rho(x, t))$ , it follows that this canonical neighborhood contains  $B(x, t, \rho(x, t))$ . Every point of the canonical neighborhood has scalar curvature  $\leq CQ_0$ . But by the curvature pinching

result, for  $t$  sufficiently large, the absolute values of negative eigenvalues of  $Rm$  at any point of this canonical neighborhood are bounded above by an arbitrarily small constant (depending on  $t$ ) times the scalar curvature. This means that no point of the canonical neighborhood, and hence no point of  $B(x, t, \rho(x, t))$ , has Riemannian curvature with an eigenvalue less than  $-\rho(x, t)^{-2}/2$ . This contradicts the definition of  $\rho(x, t)$ . The contradiction shows that every point of  $B(x, t, \rho(x, t))$  has curvature  $> Q_0$ . Since  $\rho(x, t) < \theta^{-1}(w)\hat{h}(t) < r(t)$ , it follows from this lower bound on the scalar curvature that every point of  $B(x, t, \rho(x, t))$  has a  $(C, \epsilon)$ -canonical neighborhood.  $\square$

We keep the notation that  $(y, t)$  is a point in  $B(x, t, \rho(x, t))$  with a sectional curvature which is  $\leq -\rho^{-2}(x, t)/2$ . For any  $K < \infty$ , for  $t$  sufficiently large, the curvature pinching result implies that  $R(y, t) \geq K\rho^{-2}(x, t)$ . Let  $Q_{\max}(x, t)$  be the supremum of  $R$  over  $B(x, t, \rho(x, t))$ . Then, what we have just shown is that we can write  $Q_{\max}(x, t) = C_{\max}(x, t)\rho^{-2}(x, t)$  where  $C_{\max}(x, t)$  goes to infinity as  $t$  goes to infinity. The infimum  $Q_{\min}(x, t)$  of the scalar curvature on  $B(x, t, \rho(x, t))$ . It is at least  $Q_0(t) \geq \epsilon^{-2}\theta^2/16\hat{h}^{-2}(t)$  and hence  $Q_{\min}(x, t)$  goes to infinity as  $t \rightarrow \infty$ . On the other hand, we claim that  $Q_{\min}(x, t)$  cannot be bounded above by any fixed constant  $Q$  independent of  $t$  times  $\rho^{-2}(x, t)$ . For suppose that it were so bounded. Then for all  $t$  sufficiently large we have a point of  $B(x, t, \rho(x, t))$  with scalar curvature  $\leq Q\rho^{-2}(x, t)$  and a point of scalar curvature  $C_{\max}(x, t)\rho^{-2}(x, t)$  where  $C_{\max}(x, t)$  tends to  $\infty$  as  $t$  does. Furthermore, the canonical neighborhood threshold  $r(t)$  is greater than  $\rho(x, t)$ . This contradicts Theorem 10.2 in [21] and shows that the minimum value  $Q_{\min}(x, t)$  of  $R$  over  $B(x, t, \rho(x, t))$  is at least  $C_{\min}(t)\rho^{-2}(x, t)$  where  $C_{\min}(t) \rightarrow \infty$  as  $t \rightarrow \infty$ .

Having established this lower bound for the scalar curvature on  $B(x, t, \rho(x, t))$ , it follows immediately from the nature of  $(C, \epsilon)$ -canonical neighborhoods that the volume estimate for  $B(x, t, \rho(x, t))$  is bounded above by  $w(C_{\min}(t))\rho^3(x, t)$  where  $w(s)$  is a function that tends to zero as its argument  $s$  tends to infinity. It now follows that for a given  $w$ , for  $t$  sufficiently large, if  $\rho(x, t) \leq \theta^{-1}(w)\hat{h}(t)$  then  $\text{Vol}(B(x, t, \rho(x, t))) < w\rho^3(x, t)$ .

It remains to consider:

**Case 2: For  $t$  sufficiently large  $\rho(x, t) > \theta^{-1}(w)\hat{h}(t)$ .** Suppose there is no  $\bar{\rho}$  as required. Then there is a sequence  $\bar{\rho}_n \rightarrow 0$  and a sequence of Ricci flows with surgery  $(\mathcal{M}_n, G_n)$  and points  $(x_n, t_n)$  with  $t_n \rightarrow \infty$  such that  $\rho(x_n, t_n) < \bar{\rho}_n\sqrt{t_n}$  and  $\text{Vol } B(x_n, t_n, \rho(x_n, t_n)) \geq w\rho^3(x_n, t_n)$ . Now we pass to a subsequence so that for all  $n$  we have  $\bar{\rho}_n \leq \bar{r}'(w)$  from Corollary 3.19. Then according to that corollary there are  $\tau' > 0$  and  $K' < \infty$ , depending only on  $w$ , such that  $(\mathcal{M}_n, G_n)$  contains the entire parabolic neighborhood  $P(x_n, t_n, \rho(x_n, t_n)/4, -\tau'\rho^2(x_n, t_n))$  and has scalar curvature bounded above by  $K'\rho^{-2}(x_n, t_n)$  on this parabolic neighborhood. As before, let  $X(x, t)$  denote the minimum of 0 and the negative of the smallest eigenvalue of the Riemannian curvature tensor  $Rm(x, t)$ . If  $X(x_n, t_n)(t_n + 1) \geq e^4$ , then it follows from curvature pinching (Inequality 1.2) that

$$X(x_n, t_n) \leq K'\rho^{-2}(x_n, t_n)/2.$$

On the other hand, if  $X(x_n, t_n)(t_n + 1) < e^4$ , then  $X(x_n, t_n) < e^4 t_n^{-1}$ . Since  $\rho(x_n, t_n) \leq \bar{\rho}_n \sqrt{t_n}$ , we see that

$$X(x_n, t_n) < (e^4 \bar{\rho}_n^2) \rho^{-2}(x_n, t_n).$$

The latter term is less than  $\rho^{-2}(x_n, t_n)$  for all  $n$  sufficiently large. Thus, we see that for all  $n$  sufficiently large there is a constant  $K''$ , depending only on  $w$ , such that  $X(x, t) < K'' \rho^{-2}(x_n, t_n)$  for all  $(x, t) \in P(x_n, t_n, \rho(x_n, t_n)/4, -\tau' \rho^2(x_n, t_n))$ . Having the upper bound on the scalar curvature and  $X$ , there is a constant  $K_1$  depending only on  $w$  such that for all  $n$  sufficiently large, all sectional curvatures on  $P(x_n, t_n, \rho(x_n, t_n)/4, -\tau' \rho^2(x_n, t_n))$  are bounded in absolute value by  $K_1 \rho^{-2}(x_n, t_n)$ . We pass to a subsequence so that this inequality holds for all  $n$ . Now we set

$$\alpha = \alpha(w) = \min \left( 1/4, \sqrt{\tau'}, (K_1)^{-1/2} \right).$$

Then  $(\mathcal{M}_n, G_n)$  contains the entire parabolic neighborhood  $P(x_n, t_n, \alpha \rho(x_n, t_n), -(\alpha \rho(x_n, t_n))^2)$  and has sectional curvatures bounded in absolute value by  $\alpha^{-2} \rho^{-2}(x_n, t_n)$  on this parabolic neighborhood. Since  $|Rm| \leq K_1 \rho^{-2}(x_n, t_n)$  on  $B(x_n, t_n, \rho(x_n, t_n))$  and by supposition  $\text{Vol } B(x_n, t_n, \rho(x_n, t_n)) \geq w \rho^3(x_n, t_n)$ , there is a constant  $w' > 0$  depending only on  $w$  and  $K_1$  such that  $\text{Vol } B(x_n, t_n, \alpha \rho(x_n, t_n)) \geq w' \alpha^3 \rho^3(x_n, t_n)$ .

We take  $A = \max(\alpha^{-1}, (w')^{-1})$ , so that  $A$  depends only on  $w$ . Passing to a subsequence we can suppose that  $\bar{\rho}_n \leq \bar{r}(A)$  for all  $n$  where  $\bar{r}(A)$  the constant  $\bar{r}$  of Lemma 3.1 for this value of  $A$ . Apply Part (b) of Lemma 3.1 to these neighborhoods and the constant  $A$ . We conclude that there is a constant  $K_2$  depending only on  $A$  and hence depending only on  $w$  so that

$$R(y, t_n) \leq K_2 \alpha^{-2} \rho^{-2}(x_n, t_n)$$

for all  $(y, t_n) \in B(x_n, t_n, \rho(x_n, t_n))$ . By the definition of  $\rho(x_n, t_n)$  there is a point  $(y_n, t_n) \in B(x_n, t_n, \rho(x_n, t_n))$  with the smallest negative eigenvalue of  $Rm(y_n, t_n) \leq -\rho^{-2}(x_n, t_n)/2$ , and hence the ratio of  $X(y_n, t_n)/R(y_n, t_n)$  is bounded above by  $2K_2 \alpha^{-2}$ , which depends only on  $w$ . But, for all  $n$  sufficiently large, this contradicts the pinching inequality since  $\rho(x_n, t_n)/\sqrt{t_n} \leq \bar{\rho}_n \rightarrow 0$  and hence  $\rho(x_n, t_n)^{-2} t_n \rightarrow \infty$  as  $n \rightarrow \infty$ .

This establishes Case 2 and completes the proof of the Proposition 2.8.  $\square$

### 4.3 Proof of Proposition 2.9

Recall the statement that we shall prove:

**Proposition 2.9.** *For every  $w' > 0$  there exist  $\bar{r} = \bar{r}(w') > 0$  and constants  $K_m = K_m(w') < \infty$ ,  $m = 0, 1, \dots$ , such that the following holds for any Ricci flow with surgery  $(\mathcal{M}, G)$  satisfying Assumptions 3.9 and 3.10 and for all  $t$  sufficiently large, how large depending only on  $w'$ . For any  $0 < r \leq \bar{r} \sqrt{t}$ , for any  $x \in M_t$ , and for any  $m > 0$ . Suppose that the ball  $B(x, t, r)$  has volume at least  $w' r^3$  and sectional curvatures bounded below by  $-r^{-2}$ . Then the norms of the curvature and its  $m^{\text{th}}$ -order covariant derivatives at  $(x, t)$  are bounded by  $K_0 r^{-2}$  and  $K_m r^{-(2+m)}$ , respectively.*

*Proof.* Fix  $w' > 0$  and suppose that the result doesn't hold. Then we have a sequence of Ricci flows with surgery  $(\mathcal{M}_n, G_n)$  and balls  $B_n = B(x_n, t_n, r_n) \subset \mathcal{M}_n$  with  $r_n/\sqrt{t_n} \rightarrow 0$  and  $t_n \rightarrow \infty$  as  $n \rightarrow \infty$  such that  $\text{Vol } B_n \geq w' r_n^3$  and  $Rm|_{B_n} \geq -r_n^{-2}$ , yet there are no constants  $K_0, K_1, \dots$ , as required for this sequence.

**Case 1:**  $r_n \leq \theta^{-1}(w')\hat{h}(t_n)$  for all  $n$  sufficiently large. We divide this case into two subcases: either  $R(x_n, t_n) \geq r^{-2}(t_n)$  or  $R(x_n, t_n) < r^{-2}(t_n)$ . If  $R(x_n, t_n) \geq r^{-2}(t_n)$ , then  $(x_n, t_n)$  has a  $(C, \epsilon)$ -canonical neighborhood and the existence of the constants  $K_i$  as required is immediate from the  $C^\infty$ -bounds, on these neighborhoods, which follows from the compactness, up to rescaling, of the space of  $\kappa$ -solutions..

Thus, we can suppose that  $R(x_n, t_n) < r^{-2}(t_n)$ . Next, suppose that there is some point in  $(y_n, t_n) \in B(x_n, t_n, r_n)$  with  $R(y_n, t_n) \geq r^{-2}(t_n)$ . Then there is a point  $(z_n, t_n) \in B(x_n, t_n, r_n)$  with  $R(z_n, t_n) = r^{-2}(t_n)$ . This point has a  $(C, \epsilon)$ -canonical neighborhood which contains the ball of radius  $\epsilon^{-1}r(t_n)$ . Since  $r_n \leq \theta^{-1}(w')\hat{h}(t_n)$ , we have  $r_n < \epsilon^{-1}r(t_n)/2$ , provided that  $n$  is sufficiently large. It follows that  $B(x_n, t_n, r_n)$  is contained in the  $(C, \epsilon)$ -canonical neighborhood of  $(z_n, t_n)$ . Again the result follow from the  $C^\infty$ -bounds, up to scale, of  $(C, \epsilon)$ -canonical neighborhoods.

This means that we can assume that  $R(y_n, t_n) < r^{-2}(t_n)$  for all points  $(y_n, t_n) \in B(x_n, t_n, r_n)$ . Now we rescale by  $r_n^{-2}$ , and shift  $t_n$  to zero. This gives us balls of radius 1 on which the scalar curvature is less than  $r^{-2}(t_n)r_n^2$ . Since  $r_n \leq \theta^{-1}(w')\hat{h}(t_n) < \bar{\delta}^2(t_n)r(t_n)$ , this product  $r^{-2}(t_n)r_n^2$  tends to zero as  $n$  goes to infinity. This implies that the scalar curvature on these rescaled unit balls is tending to zero as  $n \rightarrow \infty$ , and hence by curvature pinching (recall that the  $r_n/\sqrt{t_n} \rightarrow 0$  as  $n \rightarrow \infty$ ) all sectional curvatures on these balls are also tending to zero as  $n \rightarrow \infty$ . Furthermore, the canonical neighborhood threshold at time  $t \in [-t_n r_n^{-2}, 0]$  for the rescaled and shifted version of the Ricci flow with surgery is  $\leq r(r_n^2 t + t_n)r_n^{-1}$ . Since  $r(t)$  is a weakly monotone decreasing function, any point with in the rescaled parabolic neighborhood with scalar curvature  $\geq r_n^2 r(t_n)^{-2}$  has a  $(C, \epsilon)$ -canonical neighborhood. If, for every  $n$  sufficiently large, the Ricci flow with surgery  $(\mathcal{M}_n, G_n)$  contains the entire parabolic neighborhood  $P(x_n, t_n, r_n, -r_n^2)$ , then we can apply Shi's theorem (Theorem 3.28 in [21]) to show that, after passing to a subsequence and rescaling by  $r_n^{-2}$ , the limit exists on an entire abstract parabolic neighborhood  $P(x_\infty, 0, 1, -1)$  and is flat. This implies in particular, that all the higher derivatives of the Riemann curvature tensor converge to zero on the  $r_n^{-1}B(x_n, t_n, r_n)$  as  $n$  goes to infinity. Hence, the constants  $K_n$  as required exist if the Ricci flows with surgery are defined on the entire parabolic neighborhoods  $P(x_n, t_n, r_n, -r_n^2)$ .

Now suppose that, after passing to a subsequence, for each  $n$  the Ricci flow with surgery does not contain the entire parabolic neighborhood  $P(x_n, t_n, r_n, -r_n^2)$ . Then there is a backwards flow line from a point of  $B(x_n, t_n, r_n)$  that meets a surgery cap at some time in the interval  $[t_n - r_n^2, t_n]$ . (For example, the ball itself might contain a point of the surgery cap.) We take the first such cap we reach in flowing backwards from  $B(x_n, t_n, r_n)$  and let  $t'_n \geq t_n - r_n^2$  be the corresponding surgery time. Passing to a subsequence, we can suppose that  $\alpha = \lim_{n \rightarrow \infty} (t_n - t'_n)r_n^{-2}$  exists. If  $\alpha > 0$ , then we can apply the same argument as before to the smaller parabolic neighborhoods  $P(x_n, t_n, r_n, -\alpha r_n^2/2)$  to conclude there are constants  $K_n$  as required.

Thus, we can assume that  $(t_n - t'_n)r_n^{-2} \rightarrow 0$  as  $n \rightarrow \infty$ . The same curvature

argument as before shows that the rescaled scalar curvature at any point of the surgery cap at time  $t'$  that lies on a backwards flow line emanating from  $B(x_n, t_n, r_n)$  tends to zero as  $n \rightarrow \infty$ . Because the scalar curvature on the union of the entire surgery cap and the remaining half of the  $\epsilon$ -neck that the cap is glued onto varies by at most a fixed multiplicative constant, it follows that the rescaled scalar curvature on this entire union goes to zero as  $n \rightarrow \infty$ . This means that  $r_n/h(t'_n)$  tends to zero as  $n \rightarrow \infty$  where  $h(t'_n)$  is the scale of surgeries at time  $t'_n$ .

This implies that for all  $n$  sufficiently large, the result of flowing  $B(x_n, t_n, r_n)$  backwards to time  $t'_n$  is contained in the union of the surgery cap and continuing half of the  $\epsilon$ -neck at time  $t'_n$ . Now rescale by  $h^{-2}(t'_n)$ . The time interval between the surgery cap and the ball in the rescaled flow in  $h^{-2}(t_n - t'_n) < r_n^2/h^2(t'_n)$  approaches 0 as  $n \rightarrow \infty$ . After rescaling by  $h^{-2}(t'_n)$  here is a bound on the  $C^\infty$ -topology of the union of the surgery cap at time  $t'_n$  and the half of the  $\epsilon$ -neck it is glued to (see Part 5 of Theorem 12.5 in [21]). Applying the refined version of Shi's theorem (Theorem 3.29 in [21], see also Corollary 16.9 of [21]), this implies that rescaling by  $h^{-2}(t'_n)$  for all  $n$  sufficiently large there are uniform bounds  $K_0$  on the curvature and, for each  $i \geq 1$  a bound  $K_i$  on the  $i^{th}$  derivatives of curvature on the backward parabolic neighborhood on the time interval  $[t'_n, t_n]$ , whose  $t_n$  time-slice is  $B(x_n, t_n, r_n)$ . Rescaling by  $h^2(t'_n)$  to get back to the original scale in  $(\mathcal{M}_n, G_n)$  gives us the required bound  $K_0 h^{-2}(t'_n)$  on the curvature of  $B(x_n, t_n, r_n)$ , and bounds  $K_i h^{-(2+i)}(t'_n)$  on the  $i^{th}$  derivatives of the curvature on  $B(x_n, t_n, r_n)$ . Since  $r_n/h(t'_n) \rightarrow 0$  as  $n \rightarrow \infty$ , this is a contradiction.

This completes the proof in Case 1 and allows us to assume that we are in the complementary case. Passing to a subsequence allows us to assume that:

**Case 2:**  $r_n > \theta^{-1}(w')\hat{h}(t_n)$  for all  $n$ . In this case we can apply Corollary 3.19 and conclude that there are constants  $\tau' > 0$  and  $K' < \infty$  so that for all  $n$  sufficiently large, the Ricci flow with surgery contains the entire parabolic neighborhood  $P(x_n, t_n, r_n/4, -\tau' r_n^2)$  and has  $R < K' r_n^{-2}$ . Rescaling the metric and time by  $r_n^{-2}$  gives us parabolic neighborhoods  $P(x_n, t_n, 1/4, -\tau')$  on which the scalar curvature is bounded by  $K'$  and hence by the curvature pinching assumption, the Riemannian curvature tensor is uniformly bounded on these neighborhoods. Applying Shi's Theorem (3.28 of [21]) gives us the required constants  $K_i$ . This is a contradiction, concluding the proof of Proposition 2.9.  $\square$

## **PART II: Locally Volume Collapsed 3-manifolds**



## 5 Introduction to Part II

In Part I we showed that for any 3-dimensional Ricci flow with surgery with normalized initial conditions,  $(M_t, g(t))$ , for any  $w > 0$  for all sufficiently large  $t$  the manifold  $(M_t, g(t))$  contains a finite disjoint union of truncated hyperbolic manifolds of finite volume  $\mathcal{H}$  with incompressible boundary such that the complement  $(M_t(w, -), g(t))$  is  $w$  locally volume collapsed.

To complete the proof of the geometrization conjecture it suffices to show that provided that  $w$  is sufficiently small and, given  $w$ , that  $t$  is sufficiently large, the manifolds  $M_t(w, -)$  are *graph manifolds*, that is to say that the  $M_t(w, -)$  are connected sums of manifolds that are themselves unions along incompressible tori of Seifert fibrations. For this, it suffices to take a sequence  $w_n \rightarrow 0$  as  $n \rightarrow \infty$  and for each  $n$  choose  $t_n$  sufficiently large so that the above results hold for  $(M_n, g_n) = (M_{t_n}(w_n, -), (1/t_n)g(t_n))$  (and also  $t_n \rightarrow \infty$  as  $n \rightarrow \infty$ ) and show that, for every  $n$  sufficiently large,  $M_n$  is a graph manifold. That is to say, it suffices to show that the relative version of the geometrization conjecture holds for  $M_n$  for all  $n$  sufficiently large. Since we know that for all  $t$  sufficiently large, each component of  $M_t(w, -)$  is either diffeomorphic to a 3-sphere or is aspherical, we shall show that these manifolds are unions along incompressible tori of Seifert fibrations.

### 5.0.1 Seifert Fibered Manifolds and Graph Manifolds

From now on 3-manifolds are implicitly assumed to be orientable. Recall that a *Seifert fibration structure* on a compact 3-manifold is a locally-free circle action on a 2-sheeted covering  $\widetilde{M}$  of  $M$  such that, denoting the covering transformation on  $\widetilde{M}$  by  $\tau$ , we have  $\tau(\zeta \cdot x) = \bar{\zeta} \cdot x$  for all  $x \in \widetilde{M}$  and all  $\zeta \in S^1$ . Seifert fibration structures are classified in terms of their base orbifolds, local Seifert invariants, and, when the base is closed, an ‘Euler class,’ see [31] or [23]. A compact 3-manifold is said to be *Seifert fibered* if it admits a Seifert fibration structure.

**Lemma 5.1.** *A compact, connected Seifert fibered 3-manifold with compressible boundary is diffeomorphic to a solid torus. A compact, connected Seifert fibered manifold with incompressible boundary is diffeomorphic either to  $T^2 \times I$  or to a twisted  $I$ -bundle over the Klein bottle or is geometric in the sense that its interior admits a complete, locally homogeneous metric of finite volume..*

*Proof.* Let  $M$  be a compact, connected Seifert fibered 3-manifold and denote by  $\Sigma$  be the quotient 2-dimensional orbifold. If the boundary of  $M$  has a compressible torus, then the corresponding boundary component of  $\Sigma$  does not generate an infinite cyclic subgroup of  $\pi_1^{\text{orb}}(\Sigma)$ . This means that  $\Sigma$  is a topological disk with at most one singular point, and consequently that  $M$  is diffeomorphic to a solid torus.

Suppose that  $\partial M$  consists of incompressible tori. If the orbifold Euler characteristic of  $\Sigma$  is negative, then  $\Sigma$  is equivalent to a hyperbolic orbifold and  $M$  admits a geometric structure modelled on either the universal covering of  $PSL(2, \mathbb{R})$  or the product of  $\mathbb{R}$  with hyperbolic 2-space. If the orbifold Euler characteristic of  $\Sigma$  is positive, then either  $\Sigma$  is a spherical 2-dimensional orbifold, in which case  $M$  is

geometric and either admits a round metric or is modelled on  $S^2 \times \mathbb{R}$ , or  $\Sigma$  is homeomorphic to  $S^2$  with at most two singular points. In the later case,  $M$  is the union of 2 solid tori and is hence geometric. Lastly, consider the case when the orbifold Euler characteristic of  $\Sigma$  is zero. If  $\Sigma$  is without boundary, then  $M$  is geometric and either admits a flat metric or a metric modelled on the 3-dimensional nilpotent group. If  $\partial\Sigma \neq \emptyset$ , then  $\Sigma$  is either an annulus, a möbius band, or topologically the 2-disk with 2 orbifold singular points of order 2. In these cases,  $M$  is diffeomorphic to either  $T^2 \times I$  or the twisted  $I$ -bundle over the Klein bottle.  $\square$

**Definition 5.2.** A *graph manifold* is a compact 3-manifold with torus boundary each of whose prime factors can be decomposed along incompressible tori into pieces that are Seifert fibered.

**Lemma 5.3.** A *prime graph manifold* either has incompressible boundary or is a solid torus.

*Proof.* Suppose that  $X$  is prime and  $\partial X$  contains a compressible 2-torus. Then the union of a collar neighborhood of  $\partial X$  with a compressing 2-disk is compact submanifold  $Y$  of  $X$  diffeomorphic to the complement of a 3-ball in a solid torus. Since  $X$  is prime, the complement of  $Y$  in  $X$  is a 3-ball, and hence  $X$  is diffeomorphic to a solid torus.  $\square$

**Proposition 5.4.** Suppose that  $M$  is a closed 3-manifold and suppose that  $\overline{\mathcal{H}} \subset M$  is an embedding of a truncated version of a complete hyperbolic manifold of finite volume into  $M$ , suppose that the image of the boundary of  $\overline{\mathcal{H}}$  is a disjoint union of incompressible tori, and suppose that  $N = M \setminus \text{int}(\overline{\mathcal{H}})$  is a graph manifold. Then  $M$  satisfies the Geometrization Conjecture.

*Proof.* First, notice that since the boundary of  $\overline{\mathcal{H}}$  is incompressible in  $M$ , any 2-sphere in  $N$  that does not bound a 3-ball in  $N$  does not bound a 3-ball in  $M$ . Hence, we obtain the prime decomposition of  $M$  by gluing together the prime decomposition,  $N'$ , of  $N$  and  $\overline{\mathcal{H}}$  along their common boundary. Hence, it suffices to show that the union,  $M'$ , along incompressible tori of a (possibly disconnected) hyperbolic manifold and a (possibly disconnected) prime graph manifold  $N'$  satisfies the Geometrization Conjecture. By hypothesis, the boundary of  $N'$  is incompressible. Hence, according to the previous lemma, each component of  $N'$  is either  $T^2 \times I$  or a twisted  $I$ -bundle over the Klein bottle or further decomposes along incompressible tori into Seifert fibrations with incompressible boundary. We start with  $\mathcal{T}_0$  equal to the boundary components of  $\overline{\mathcal{H}}$  together with all the incompressible tori that are used to divide components of  $N'$  into Seifert fibrations with incompressible boundary. This collection of tori decomposes  $M'$  into hyperbolic pieces, Seifert fibrations with incompressible boundary, pieces diffeomorphic to  $T^2 \times I$  and pieces diffeomorphic to twisted  $I$ -bundles over a Klein bottle. Any closed manifold that is a union along the boundary of twisted  $I$ -bundles over the Klein bottle and copies of  $T^2 \times I$  is geometric (either flat or modelled on the solvable 3-dimensional Lie group). This allows us to assume that no component of  $M'$  is of this form. Now if distinct components  $T$  and  $T'$  of  $\mathcal{T}_0$  are parallel, then we remove one of them from the collection and call the new

collection  $\mathcal{T}_0$ . It divides  $M'$  into the same types of pieces as the original collection does. We repeat this operation, until no distinct components of  $\mathcal{T}_0$  are parallel tori. Now if a component  $T$  of  $\mathcal{T}_0$  bounds a twisted  $I$ -bundle over the Klein bottle in  $M'$ , then by assumption it does so on only one side. In this case we change  $\mathcal{T}_0$  by replacing  $T$  by the 0-section Klein bottle in this neighborhood. Again, the complementary pieces are of the same types as before. Continuing in this manner, allows us to assume that no 2-torus in  $\mathcal{T}_0$  bounds a twisted  $I$ -bundle over the Klein bottle and no distinct 2-tori in  $\mathcal{T}_0$  are parallel. Hence, all the complementary components of  $\mathcal{T}_0$  are geometric.  $\square$

### 5.0.2 The Statement

Reformulating what we have established in Section 2.2, we see that the  $(M_n, g_n) = (M_{t_n}(w_n, -), (1/t_n)g(t_n))$  satisfy the hypotheses of the following theorem. The theorem then tells us that for all  $n$  sufficiently large  $M_n$  is a graph manifold.

**Theorem 5.5.** *(Theorem 7.4 of [27]) Suppose that  $(M_n, g_n)$  is a sequence of compact, oriented Riemannian 3-manifolds, closed or with convex boundary, and that  $w_n$  is a sequence of positive numbers tending to zero as  $n$  tends to  $\infty$ . Assume that:*

1. *For each point  $x \in M_n$  there exists a radius  $\rho = \rho_n(x)$  such that the ball  $B_{g_n}(x, \rho)$  has volume at most  $w_n \rho^3$  and all the sectional curvatures of the restriction of  $g_n$  to this ball are all bounded below by  $-\rho^{-2}$ ;*
2. *There is a constant  $K < \infty$  such that the following holds. Each component of the boundary of  $M_n$  is locally convex and is an incompressible torus of diameter at most  $Kw_n$  and with a topologically trivial collar containing the all points within distance 1 of the boundary and on which the sectional curvatures are between  $-5/16$  and  $-3/16$ ;*
3. *For every  $w' > 0$  there exist  $\bar{r} = \bar{r}(w') > 0$  and constants  $K_m = K_m(w') < \infty$  for  $m = 0, 1, 2, \dots$ , such that for all  $n$  sufficiently large, and any  $0 < r \leq \bar{r}$ , if the ball  $B_{g_n}(x, r)$  has volume at least  $w' r^3$  and sectional curvatures bounded below by  $-r^{-2}$ , then the curvature and its  $m^{\text{th}}$ -order covariant derivatives,  $m = 1, 2, \dots$ , at  $x$  are bounded by  $K_0 r^{-2}$  and  $K_m r^{-m-2}$ , respectively.*

*Then for every  $n$  sufficiently large  $M_n$  is a graph manifold.*

Take a sequence  $w_n \rightarrow 0$ . Recall from Theorem 2.24 and Proposition 2.25 in Part 1 the following hold provided that we have a Ricci flow with surgery satisfying the hypotheses of Corollary 15.10 of [21] with the surgery control function  $\bar{\delta}(t)$  and the surgery scale function  $h(t)$  satisfy Assumptions 3.9 and 3.10. There is a sequence  $t_n \rightarrow \infty$  and a complete, finite volume hyperbolic manifold  $\mathcal{H}$  such that for each  $n$  there is an embedding of a truncated version  $\bar{\mathcal{H}}(w_n)$  of  $\mathcal{H}$  into  $M_{t_n}$  whose complement satisfies the first two conditions of the above theorem. Also, by Proposition 2.9 the third condition in the above theorem also holds of the complement  $M_{t_n} \setminus \text{int}(\bar{\mathcal{H}}(w_n))$ . Thus, as a consequence of this theorem we have:

**Corollary 5.6.** *The Geometrization Conjecture is true for all closed, orientable 3-manifolds.*

Sections 6 through 12 are devoted to establishing Theorem 5.5.

### 5.0.3 Stronger Results

Using the full strength of what was proved in [21] we can in fact make a much stronger statement about  $M_t$ . Recall from Proposition 18.9 of [21] and the Poincaré Conjecture, it follows that for all  $t$  sufficiently large, that every component of  $M_t$  is irreducible and hence either prime or diffeomorphic to  $S^3$ . Thus, for sufficiently large time, every  $S^2$ -surgery is along a separating 2-sphere bounding a 3-ball and produces the disjoint union of a 3-sphere and a manifold diffeomorphic to the manifold before surgery. From this we deduce:

**Corollary 5.7.** *Given a Ricci flow with surgery satisfying the hypotheses of Corollary 15.10 in [21] with the surgery control function  $\bar{\delta}(t)$  and the surgery scale function  $h(t)$  satisfying Assumptions 3.9 and 3.10, for all  $t$  sufficiently large there is a finite set of incompressible tori in  $M_t$  such that each component of the complement satisfies one of the following:*

1. *The component is diffeomorphic to  $S^3$ .*
2. *The component admits a complete hyperbolic metric of finite volume.*
3. *The component is the interior of a compact Seifert fibered 3-manifold with incompressible boundary.*
4. *The component is closed and admits a locally homogeneous metric of Solv, Nil, or Flat type.*

Since a component of the third and four types either admits a complete, locally homogeneous metric of finite volume or the component is diffeomorphic to either  $T^2 \times \mathbb{R}$  or to the twisted  $\mathbb{R}$ -bundle over the Klein bottle, we have:

**Corollary 5.8.** *Given a Ricci flow with surgery satisfying Corollary 15.10 of [21] and with the surgery control function  $\bar{\delta}(t)$  and the surgery scale function  $h(t)$  satisfying Assumption 3.9 and 3.10, the following holds for all  $t$  sufficiently large. Removing a finite set of incompressible tori and Klein bottles from  $M_t$  yields a manifold each component of which has a complete, locally homogeneous metric of finite volume.*

## 6 The Collapsing Theorem

### 6.1 First remarks

According to Theorem 1.17 in Section 1.6 of [1], a closed, connected 3-manifold admitting a flat metric is Seifert fibered and hence is a graph manifold. If a closed, orientable 3-manifold has a metric of non-negative sectional curvature then by [11] it is diffeomorphic to one of the following:

1. a spherical 3-dimensional space-form,
2. a manifold with a locally homogeneous metric modelled on  $S^2 \times \mathbb{R}$ , or
3. a flat 3-manifold.

Thus, without loss of generality we can make the following assumption.

**Assumption 1.** *For each  $n$ , no connected, closed component of  $M_n$  admits a Riemann metric of non-negative sectional curvature.*

The idea of the proof of Theorem 5.5 is to consider a sequence of balls of the form  $B_{g'_n(x)}(x, 1) \subset M_n$ ,  $n = 1, 2, \dots$ , where by definition  $g'_n(x) = \rho_n^{-2}(x)g_n$ . The hypotheses of the theorem and Assumption 1 imply that each of these balls is non-compact, but locally complete and of sectional curvature  $\geq -1$ . The general theory of Alexandrov spaces implies that given any such sequence there is a subsequence that converges in the sense of Gromov-Hausdorff to a ball of radius one in an Alexandrov space of curvature  $\geq -1$  and of dimension at least 1 and at most 3. The hypothesis that the volume of  $B_{g'_n(x)}(x, 1)$  is at most  $w_n$  and the fact that the  $w_n \rightarrow 0$  imply that the limit is a 1- or 2-dimensional. We then use results on the structure of Alexandrov spaces of dimension 1 and 2 to deduce strong topological and geometric information about the structure of these balls in  $M_n$  for all  $n$  sufficiently large. These local structures can then be pieced together to form a global result, proving the theorem stated above. We review this background material on Gromov-Hausdorff convergence and Alexandrov spaces in Sections 8, 9, and 10, but in this introduction we assume that these basic notions are understood and we formulate the precise structural results that will be proved. In Section 11 we deduce the local results, i.e., the possible structures of the balls  $B_{g'_n(x)}(x, 1)$ , and in Section 12 we piece the local results together proving the main topological decomposition result, Theorem 6.2 below. As we show below this result easily implies that the  $M_n$  are graph manifolds for all  $n$  sufficiently large.

### 6.1.1 Adjusting $\rho_n$

There is one simplification in Theorem 5.5 that is important to point out.

**Lemma 6.1.** *Let  $M_n$ ,  $w_n$  and  $\rho_n$  satisfy the hypotheses of Theorem 5.5 and suppose that the  $M_n$  satisfy Assumption 1. After passing to a subsequence, and replacing  $w_n$  and  $\rho_n$  by other constants and functions we can arrange that the hypothesis of Theorem 5.5 are satisfied and in addition the following hold:*

1. *For any connected component  $M_n^0$  of  $M_n$  and for any  $x \in M_n^0$  we have*

$$\rho_n(x) \leq \text{diam } M_n^0,$$

*and*

2. *if, for some  $0 < r_1, r_2 < 1$  we have  $B_{g'_n(x)}(x, r_1) \cap B_{g'_n(y)}(y, r_2) \neq \emptyset$  then*

$$\frac{1 - r_1}{1 + r_2} < \frac{\rho_n(y)}{\rho_n(x)} < \frac{1 + r_1}{1 - r_2}.$$

*Proof.* Without loss of generality we can assume that  $M_n$  is connected. If  $M_n$  is closed, then by assumption it is not the case that  $\text{Rm} \geq 0$  on all of  $M_n$ . If  $M_n$  has non-empty boundary, then also by assumption  $\text{Rm}$  is not everywhere positive. Thus, for each  $x \in M_n$ , there is a maximum  $r_n(x) \geq \rho_n(x)$  such that the  $\text{Rm} \geq -r_n(x)^{-2}$  on  $B(x, r_n(x))$ . Furthermore, by volume comparison (the Bishop-Gromov theorem)

$$\text{vol } B(x, r_n(x)) \leq \frac{V_{\text{hyp}}(1)}{V_{\text{Eucl}}(1)} w_n r_n^3(x),$$

where  $V_{\text{hyp}}(1)$ , resp.  $V_{\text{Eucl}}(1)$ , is the volume of the unit ball in hyperbolic, resp. Euclidean, 3-space. Thus, at the expense of changing the  $w_n$  by a factor independent of  $n$ , we define the function  $\rho_n$  so that  $\rho_n(x)$  is this maximum  $r_n(x)$ . Inequality 2 follows immediately for this choice.

Now suppose (after passing to a subsequence) that for each  $n$  there is  $x \in M_n$  with  $\rho_n(x) > \text{diam } M_n$ . This implies that  $\text{Rm}(x) \geq -(\text{diam } M_n)^{-2}$  for all  $x \in M_n$  and hence that  $\rho_n$  is a constant function; we denote its value by  $\rho_n$ . Passing to a subsequence we can assume that  $\text{vol}(M_n)/(\text{diam } M_n)^3$  tends to a limit (possibly  $+\infty$ ) as  $n \rightarrow \infty$ . First, we consider the case when this limit is non-zero. The fact that the volume divided by the cube of the diameter is bounded away from zero and the volume inequality assumed in Theorem 5.5 imply that  $\text{diam } M_n/\rho_n$  tends to 0 as  $n \rightarrow \infty$ . By the hypothesis about the boundary of  $M_n$ , this implies that  $M_n$  is closed. Rescaling  $M_n$  to make its diameter 1 yields a manifold whose sectional curvatures are bounded below by  $-(\text{diam } M_n)^2/\rho_n^2$  and whose volume is bounded away from zero. By Proposition 9.46 we see that passing to a subsequence there is a smooth limit which has non-negative sectional curvature. This is contrary to Assumption 1. Thus, we can suppose that  $\text{vol}(M_n)/(\text{diam } M_n)^3$  tends to zero as  $n$  goes to infinity. In this case we take  $w'_n = \text{vol}(M_n)/(\text{diam } M_n)^3$  and we take  $\rho_n$  to be the constant  $\text{diam } M_n$ . Obviously, Inequality 2 holds in this case.  $\square$

**Assumption 2 and notation:** Now we fix the constants  $w_n$  and the functions  $\rho_n: M_n \rightarrow (0, \infty)$  satisfying Lemma 6.1. For any  $n$  and any  $x \in M_n$  we denote by  $g'_n(x)$  the metric  $\rho_n(x)^{-2}g_n$ . Thus,  $B_{g_n}(x, \rho_n(x)) = B_{g'_n(x)}(x, 1)$  as subsets of  $M_n$ .

## 6.2 The collapsing theorem

Let us now state the topological theorem that is established using the compactness of Alexandrov spaces of curvature  $\geq -1$  and the volume collapsing hypotheses.

**Theorem 6.2.** *Suppose that we have a sequence of compact 3-manifolds satisfying the hypothesis of Theorem 5.5 and satisfying Assumption 1. Then, for every  $n$  sufficiently large there are compact, codimension-0, smooth submanifolds  $V_{n,1} \subset M_n$  and  $V_{n,2} \subset M_n$  with  $\partial M_n \subset V_{n,1}$  satisfying the following.*

1. *Each connected component of  $V_{n,1}$  is diffeomorphic to one of the following:*

- (a) *a  $T^2$ -bundle over  $S^1$  or a union of two twisted  $I$ -bundles over the Klein bottle along their common boundary;*

- (b)  $T^2 \times I$  or  $S^2 \times I$ , where  $I$  is a closed interval;
- (c) a compact 3-ball or the complement of an open 3-ball in  $\mathbb{R}P^3$ ;
- (d) a twisted  $I$ -bundle over the Klein bottle; or a solid torus.

In particular, every boundary component of  $V_{n,1}$  is either a 2-sphere or a 2-torus.

2.  $V_{n,2} \cap V_{n,1} = \partial V_{n,2} \cap \partial V_{n,1}$ .
3. If  $X_0$  is a 2-torus component of  $\partial V_{n,1}$ , then  $X_0 \subset \partial V_{n,2}$  if and only if  $X_0$  is not a boundary component of  $M_n$ .
4. If  $X_0$  is a 2-sphere component of  $\partial V_{n,1}$ , then  $X_0 \cap \partial V_{n,2}$  is diffeomorphic to an annulus.
5.  $V_{n,2}$  is the total space of a Seifert fibration and  $\partial V_{n,1} \cap \partial V_{n,2}$  is saturated under the induced  $S^1$ -fibration on  $\partial V_{n,2}$ .
6.  $M_n \setminus \text{int}(V_{n,2} \cup V_{n,1})$  is a disjoint union of solid cylinders, i.e., copies of  $D^2 \times I$ , and solid tori. The boundary of each solid torus is a boundary component of  $V_{n,2}$ , and each solid cylinder  $D^2 \times I$  meets  $V_{n,1}$  exactly in  $D^2 \times \partial I$ .

### 6.3 Proof that Theorem 6.2 implies Theorem 5.5

In deducing Theorem 5.5 from Theorem 6.2 we shall introduce several topological simplifications in the decomposition given in the conclusion of Theorem 6.2. While the decomposition given in Theorem 6.2 is deduced from the collapsing theory (in particular,  $V_{n,1}$  is the part of  $M_n$  close to a 1-dimensional space and  $V_{n,2}$  is the part close to a 2-dimensional space), as we modify the decomposition we work purely topologically and do not try to keep the connection with the collapsing geometry.

**Claim 6.3.** *It suffices to establish Theorem 5.5 under the assumption that we have a decomposition as given in Theorem 6.2 that satisfies the following additional properties:*

1.  $V_{n,1}$  has no closed components.
2. Each 2-sphere component of  $\partial V_{n,1}$  bounds a 3-ball component of  $V_{n,1}$ .
3. Each 2-torus component of  $\partial V_{n,1}$  that is compressible in  $M_n$  bounds a solid torus component of  $V_{n,1}$ .

*Proof.* By assumption, each closed component of  $V_{n,1}$  can be decomposed along a single incompressible  $T^2$  into Seifert fibered manifolds, and hence these satisfy the conclusion of Theorem 5.5. Thus, without loss of generality we can assume that there are no closed components of  $V_{n,1}$ . In the similar way, we can suppose that no component of  $M_n$  is the union of two solid tori, the union of a solid torus and a twisted  $I$ -bundle over the Klein bottle, or the union of two twisted  $I$ -bundles over the Klein bottle along a common boundary torus, since manifolds of the first two

types admit Riemannian metrics of non-negative sectional curvature and those of the third type decompose along an incompressible torus into pieces that are Seifert fibered.

Let  $C$  be a 2-sphere component of  $\partial V_{n,1}$ . If  $C$  bounds a component  $\hat{C}$  of  $V_{n,1}$  diffeomorphic to  $\mathbb{R}P^3 \setminus B^3$ , then we remove  $\hat{C}$  from  $M_n$  and from  $V_{n,1}$  and replace it in each with a 3-ball in each. This has the effect of removing a prime factor diffeomorphic to  $\mathbb{R}P^3$  from  $M_n$ . This allows us to assume that there are no components of  $V_{n,1}$  diffeomorphic to  $\mathbb{R}P^3 \setminus B^3$  and hence that the only components of  $V_{n,1}$  with boundary 2-spheres are either 3-balls or diffeomorphic to  $S^2 \times I$ .

Now let  $C$  be a 2-sphere component of  $\partial V_{n,1}$ , but not bounding a 3-ball component of  $V_{n,1}$ . We cut  $M_n$  open along  $C$  and cap off the resulting two copies of  $C$  with 3-balls. We add these balls to  $V_{n,1}$  forming  $V'_{n,1}$ , and we leave  $V_{n,2}$  unchanged. The resulting subsets  $V'_{n,1}$  and  $V_{n,2}$  satisfy all the conclusions of Theorem 6.2. If we can show that the result is a graph manifold, then the same is true for  $M_n$ . Induction then allows us to assume that every  $S^2$ -boundary component of  $V_{n,1}$  bounds a 3-ball component of  $V_{n,1}$ .

Next, we consider a 2-torus component  $T$  of  $\partial V_{n,1}$  that is a compressible 2-torus in  $M_n$ , but one that does not bound a solid torus component of  $V_{n,1}$ . By Dehn's lemma there is an embedded disk in  $M_n$  meeting  $T$  only along its boundary, that intersection being homotopically non-trivial in  $T$ . First, suppose that  $T$  separates  $M_n$ . We write  $M_n = P \cup_T N$ . A thickening of  $T \cup D$  has a 2-sphere boundary component  $S$ , which we can suppose (by reversing the labels of the sides if necessary) lies in  $P$ . Let  $R$  be the region between  $T$  and  $S$ ; it is diffeomorphic to the complement in a solid torus of a 3-ball. We form  $A = P \cup_T F$  where  $F$ , is a solid torus, glued in such a way that  $R \cup_T F$  is diffeomorphic to a 3-ball. We set  $V_{n,2}(A) = V_{n,2} \cap P$  and  $V_{n,1}(A) = (V_{n,1} \cap A) \cup F$ . We also form  $B = \hat{R} \cup_T N$  where  $\hat{R}$  is the solid torus obtained from  $R$  by attaching a 3-ball to its  $S^2$ -boundary. We set  $V_{n,2}(B) = V_{n,2} \cap N$  and  $V_{n,1}(B) = (V_{n,1} \cap N) \cup \hat{R}$ . It is easy to see that  $M_n$  is diffeomorphic to  $A \# B$  and that the given decompositions of  $A$  and  $B$  satisfy all the conclusions of Theorem 6.2 unless  $T$  bounds a component of  $V_{n,1}$  that is a twisted  $I$ -bundle over the Klein bottle. In this case, that component of  $V_{n,1}$  is  $N$  and  $\hat{R} \cup_T N$  is Seifert fibered, whereas the conclusions of Theorem 6.2 hold for  $A$ . By a straightforward induction argument, this allows us to assume that every compressible 2-torus component of  $\partial V_{n,1}$  that separates  $M_n$  bounds a solid torus component of  $V_{n,1}$ . If  $T$  does not separate  $M_n$  we cut  $M_n$  open along  $T$ , add a solid torus  $F$  as before to the copy of  $T$  bounding  $R$  and add a copy of  $\hat{R}$  to the other copy of  $T$ . Then  $M_n$  is diffeomorphic to the connected sum of the resulting manifold,  $M'_n$ , and  $S^2 \times S^1$ . Furthermore, adding  $\hat{R} \amalg F$  to  $V_{n,1}$  and leaving  $V_{n,2}$  unchanged produces a new decomposition satisfying the hypotheses of Theorem 6.2. Again a simple induction argument shows that repeated application of this operation removes all non-separating compressing tori boundary components of  $V_{n,1}$  without creating any new compressing tori boundary components that do not bound solid torus components of  $V_{n,1}$ . This completes the proof of the claim.  $\square$

With all these simplifying assumptions in place, we are ready to complete the proof that Theorem 6.2 implies Theorem 5.5. Let us consider the union,  $X$ , of the



$D^2 \times I$  components of the closure of  $M_n \setminus (V_{n,1} \cup V_{n,2})$  and the 3-ball components of  $V_{n,1}$ . Every 2-sphere boundary component of  $V_{n,1}$  bounds a 3-ball component of  $V_{n,1}$ , each  $D^2 \times I$  meets the disjoint union of the 3-balls exactly in  $D^2 \times \partial I$  and the boundary of each 3-ball contains exactly two disks in common with  $\coprod D^2 \times \partial I$ . It then follows from the fact that  $M_n$  is orientable that  $X$  is diffeomorphic to a disjoint union of a finite number of solid tori. Hence, the closure of  $M_n \setminus V_{n,2}$  is a finite collection of solid tori, components diffeomorphic to  $T^2 \times I$ , and components diffeomorphic to twisted  $I$ -bundles over the Klein bottle. Furthermore, all boundary components of the  $T^2 \times I$  and twisted  $I$ -bundles over the Klein bottle are incompressible in  $M_n$ . We remove from  $M_n$  all components of  $M_n \setminus V_{n,2}$  diffeomorphic to either  $T^2 \times I$  or to a twisted  $I$ -bundle over the Klein bottle. The result,  $W_n$ , is a manifold that is the union of  $V_{n,2}$  and a collection of solid tori glued in along boundary components. According to [35], since  $V_{n,2}$  is a Seifert fibration,  $W_n$  is a graph manifold. Since the tori boundary components that we cut along are incompressible,  $\partial W_n$  consists of incompressible boundary tori. It follows that each prime factor of  $W_n$  has the property that removing a disjoint union of submanifolds diffeomorphic to  $T^2 \times I$  and twisted  $I$ -bundles over the Klein bottle results in an open manifold each component of which admits complete homogeneous metrics of finite volume. The same is then true of  $M_n$ .

This completes the proof that Theorem 6.2 implies Theorem 5.5.

There is an addendum which will be important later

**Remark 6.4.** Suppose that every component of  $M_n$  is aspherical. Then no component of  $M_n$  is the union of a Seifert fibration and solid tori where at least one of the solid tori is glued in in such a way as to kill the homotopy class of the generic fiber of the Seifert fibration. The above argument implies that removing from  $M_n$  copies of  $T^2 \times I$  and twisted  $I$ -bundles over the Klein bottle yields a manifold each component of which is aspherical with incompressible boundary and is a Seifert fibration over a geometric 2-dimensional orbifold or is a  $T^2$ -bundle over the circle.

The rest of this paper is devoted to the proof of Theorem 6.2.

## 7 Overview of the rest of the argument

As we indicated above, the proof of Theorem 6.2 proceeds by finding local models for neighborhoods of every point of  $M_n$  for all  $n$  sufficiently large. This is done as follows. We show that given any sequence  $x \in M_n$ , after passing to a subsequence, the unit balls  $\rho^{-1}(x)B(x, \rho(x))$  converge in a Gromov-Hausdorff sense to an Alexandrov space of dimension 0, 1 or 2. The local structures of these spaces are fairly easy to understand. From these local structures we deduce local models for balls centered about  $x$  in the 3-manifolds  $M_n$ . We then show that these local models overlap in sufficiently nice ways that we can deduce the global topology of the  $M_n$  for all  $n$  sufficiently large.

Here we describe in outline the nature of the convergence in question and the nature of the limiting spaces (Alexandrov spaces). Then we turn to the local nature

of the limits and the consequences for the possible local natures of the 3-manifolds  $M_n$ . Finally, we indicate how to glue together the local structures on the  $M_n$  to produce the global collapsing results stated above.

The convergence that we deal with is Gromov-Hausdorff convergence, which is a notion of convergence for general metric spaces. Two metric spaces are *close in the Gromov-Hausdorff sense* if they can be isometrically embedded into a third metric space so that each is contained in a small neighborhood of the other. For example a  $n$ -dimensional manifold which is fibered over a  $k$ -manifold with all fibers having small diameter is close to the  $k$ -manifold base. In general, a Riemannian manifold can be close in this sense to a metric space that is not a Riemannian manifold. There are however some geometric properties that are preserved under Gromov-Hausdorff limits. One of the properties that we are interested in is a metric version of curvature  $\geq k$  for some constant  $k$ . The source of this idea is the theorem due to Toponogov [34] which says that in a complete Riemannian manifold of curvature  $\geq k$  given a geodesic triangle  $T = abc$  the following holds. Let  $\tilde{T} = \tilde{a}\tilde{b}\tilde{c}$  be a  $k$ -comparison triangle, i.e., a triangle in the complete, simply connected surface of constant curvature  $k$  with the same pairwise distances. Then the angle of  $\tilde{T}$  at  $\tilde{b}$  is no larger than the angle of  $T$  at  $b$ . This leads to the following notion. Let  $X$  be a metric space and  $a, b, c$  be three points in  $X$ . Given a real number  $k$ , we define the  $k$ -comparison angle,  $\tilde{\angle}_k abc$ , to be the angle at  $\tilde{b}$  of the  $k$ -comparison triangle  $\tilde{a}\tilde{b}\tilde{c}$ . Then we say that a metric space has curvature  $\geq k$  if for every 4 points  $x, a, b, c$  the sum of the three  $k$ -comparison angles at  $x$  formed from these points is at most  $2\pi$ . Such spaces are called *Alexandrov spaces of curvature  $\geq k$*  if in addition they are complete metric spaces and they are length spaces in the sense that every pair of points is joined by an isometric embedding of an interval. Toponogov's theorem immediately implies that a complete Riemannian manifold of Riemannian curvature  $\geq k$  is an Alexandrov space of curvature  $\geq k$ .

It is direct from the definition that the Gromov-Hausdorff limit of a sequence of Alexandrov spaces of curvature  $\geq k$  is again an Alexandrov space of curvature  $\geq k$ . It is also clear that the Hausdorff dimension of a Gromov-Hausdorff limit is no greater than the  $\liminf$  of the Hausdorff dimensions of the spaces in the sequence. Also, it turns out that an Alexandrov space of finite Hausdorff dimension has an open dense set that is a topological manifold and the dimension of this manifold is the Hausdorff dimension of the Alexandrov space, so that in particular, the Hausdorff dimension of an Alexandrov space is either  $\infty$  or a non-negative integer. Hence, a Gromov-Hausdorff limit of Riemannian  $n$  manifolds of curvature  $\geq k$  is an Alexandrov space of curvature  $\geq k$  and Hausdorff dimension at most  $n$ . There is also a sequential compactness result for Alexandrov spaces of curvature  $\geq k$  and dimension  $\leq n$ , and there are also local versions of these arguments that apply to metric balls instead of complete metric spaces. Thus, for any sequence  $x \in M_n$  as  $n \rightarrow \infty$ , after passing to a subsequence there is a Gromov-Hausdorff limit of the unit balls  $B_{\rho_n^{-2}(x)g}(x, 1)$ . This limit is an Alexandrov ball of curvature  $\geq -1$  and dimension  $\leq 3$ . In fact, because of the volume collapsing hypothesis the limit is an Alexandrov ball of dimension at most 2.

If the limit is a point, then it is an easy matter to rescale the manifolds  $M_n$

so that their diameters are 1 and then pass to a subsequence with a limit which is an Alexandrov space of curvature  $\geq 0$  and of dimension 1, 2, 3. If the limit has dimension 3, then the bounds on the derivatives of the curvature given in Condition 3 in Theorem 5.5 imply that the convergence is smooth and the limit is a manifold of curvature  $\geq 0$ . These are completely classified and all of them satisfy the Geometrization Conjecture. This allows us to assume that the Gromov-Hausdorff limit has dimension 1 or 2.

The next step is to study the local nature of these limits. Let us describe what happens when the limiting Alexandrov space is 1-dimensional. In this case it is either an interval (open, half-closed or closed) or a circle. The local structure of the 3-manifolds converging to such Alexandrov space near points converging to an interior point is a product of  $S^2 \times (0, 1)$  or  $T^2 \times (0, 1)$  where the surface fibers are of diameter converging to zero and the interval has length bounded away from zero. In fact we can view neighborhoods in the  $M_n$  as fibering over the limiting open interval or circle with fibers of small diameter which are either  $S^2$ -fibers or  $T^2$ -fibers. Near an end point the structure is either a 3-ball or a punctured  $\mathbb{R}P^3$  (when the fibers over interior points are  $S^2$ ) or a solid torus or a twisted  $I$ -bundle over the Klein bottle (when the fibers over the interior points are 2-tori).

We cut the manifold  $M_n$  open along central tori and 2-spheres, one in each almost 1-dimensional region to produce a manifold  $M'_n$  with boundary a disjoint union of 2-spheres and 2-tori.

Now we consider the second possibility when the limiting Alexandrov space is 2-dimensional. As we shall see, we fix  $\delta > 0$  sufficiently small and then we write a 2-dimensional Alexandrov space as a union four types of points:

- interior points that are the center of neighborhoods close to open balls in  $\mathbb{R}^2$ ,
- points at which the space is an almost circular cone of cone angle  $\leq 2\pi - \delta$ ,
- boundary points that are the center of neighborhoods close to open balls centered at boundary points of half-space, and
- boundary points at which is space is almost isometric to flat cone in  $\mathbb{R}^2$  of cone angle  $\leq \pi - \delta$ .

The next step is to transfer this local information about the 2-dimensional limits to local models for neighborhoods of  $x \in M_n$ . In the four cases just listed the local models are:

- $S^1 \times B(0, \epsilon^{-1})$  with a Riemannian metric which, after an overall change of scale, is almost a product of a flat metric of length 1 on  $S^1$  with a flat metric on the ball of radius  $\epsilon^{-1}$  in  $\mathbb{R}^2$ ;
- a solid torus;
- fibered over  $\mathbb{R}$  with each fiber a topological  $D^2$ ;
- a 3-ball.

It turns out that we have sufficient geometric control over these neighborhoods to show that they are glued together in completely standard ways. Thus, any compact subset of the open set of points of the first type is contained in a open set that is smoothly fibered by circles and the circle fibers of this fibration almost line up with the circles in the almost product structures. The solid tori over the interior cone points then are glued in and the circle fibration structure extends to a Seifert fibration with at most one exceptional fiber for each solid torus. This gives a large subset of the manifold that is Seifert fibered. The rest of the manifold is made out of union of cylinders,  $D^2 \times I$ , 3-balls and  $S^2 \times I$ , with each  $S^2 \times I$  containing a boundary component of  $M'_n$ . The cylinders meet end-on-end or meet the 3-balls or the  $S^2 \times I$  in 2-disks ends. Each boundary  $S^2$ -sphere of a 3-ball or of  $S^2 \times I$  that is not a boundary component of  $M'_n$  meets exactly 2 of the cylinders. Thus, the union of these regions is diffeomorphic to a disjoint union of punctured solid tori, one puncture for each  $S^2 \times I$ . Furthermore, the torus boundary of each of these regions is contained in the open subset of  $M'_n$  which is Seifert fibered and in this region these tori are isotopic to tori saturated under the fibration structure. Of course,  $M_n$  is obtained from  $M'_n$  by gluing together boundary components. It then is an elementary exercise in 3-dimensional topology to show that such a 3-manifold is in fact a graph manifold.

In Section 8 we introduce the basics of Gromov-Hausdorff convergence. In Section 9 we turn to the basics of Alexandrov spaces. In Section 10 we study the local structure of 2-dimensional Alexandrov spaces. Then in Section 11 we deduce the local structure of the 3-manifolds  $M_n$  that follow from the results about Alexandrov spaces of dimension 2. Finally, in Section 12 we show how to piece together the local results to give the global structure theorem. Lastly, in Section 13 we extend the result to an equivariant one for compact group actions, e.g., finite group actions.

## 8 Basics of Gromov-Hausdorff Convergence

### 8.1 Limits of compact metric spaces

We begin with a review of Hausdorff and Gromov-Hausdorff limits of metric spaces.

**Definition 8.1.** Let  $X$  and  $Y$  be compact metric spaces. Consider a metric space  $Z$  and isometric embeddings of  $X$  and  $Y$  into  $Z$ . The *Hausdorff distance in  $Z$*  between  $X$  and  $Y$  is the infimum of  $a > 0$  such that every point of  $Y$  is within distance  $a$  of  $X$  and every point of  $X$  is within distance  $a$  of  $Y$ . The *Gromov-Hausdorff distance* from  $X$  to  $Y$  is the infimum over all  $Z$  and all isometric embeddings of  $X$  and  $Y$  into  $Z$  of the Hausdorff distance in  $Z$  between  $X$  and  $Y$ . Equivalently, the Gromov-Hausdorff distance between  $X$  and  $Y$  is the infimum of the Hausdorff distance between  $X$  and  $Y$  in metrics on  $X \amalg Y$  extending the given metrics on  $X$  and  $Y$ . It is easy to see that two compact metric spaces are isometric if and only if the Gromov-Hausdorff distance between them is 0.

We say that a sequence  $X_n$  of compact metric spaces *converges in the Gromov-Hausdorff sense* to a compact metric space  $X_\infty$  if the Gromov-Hausdorff distance

between  $X_n$  and  $X_\infty$  goes to zero as  $n \rightarrow \infty$ . It is elementary to show that a sequence of compact metric spaces has at most one compact Gromov-Hausdorff limit up to isometry.

Suppose that the  $X_n$  converge in the Gromov-Hausdorff sense to  $X$ . Then a *realization* of this limit is a sequence of isometric embeddings  $X_n, X \rightarrow Z_n$  so that the Hausdorff distance from  $X_n$  and  $X$  in  $Z_n$  goes to zero as  $n \rightarrow \infty$ . Equivalently, a realization is a sequence of metrics  $d_n$  on  $X_n \amalg X_\infty$  extending the given metrics on the two factors so that the Hausdorff distance in  $d_n$  between the two factors goes to 0 as  $n \rightarrow \infty$ . Given a realization we say that a sequence  $x_n \in X_n$  converges to  $x \in X$ . If the distance in  $Z_n$  between  $x_n$  and  $x$  goes to zero as  $n \rightarrow \infty$ .

**Lemma 8.2.** *Given a realization of a Gromov-Hausdorff limit  $X_n, X \subset Z_n$ ,  $n = 1, 2, \dots$ , with the  $X_n$  and  $X$  being compact Hausdorff spaces, any sequence  $x_n \in X_n$  has a subsequence converging to a point  $x \in X$ .*

*Proof.* Let the Gromov-Hausdorff distance between  $X_n$  and  $X$  in  $Z_n$  be  $\epsilon_n$ . Of course, by definition  $\epsilon_n \rightarrow 0$  as  $n \rightarrow \infty$ . Then for each  $n$  there is a point  $\hat{x}_n \in X$  such that the distance in  $Z_n$  between  $x_n$  and  $\hat{x}_n$  is at most  $\epsilon_n$ . Passing to a subsequence, we can assume that the  $\hat{x}_n$  converge to a point  $x \in X$ . This is the limit of the corresponding subsequence of the  $x_n$ .  $\square$

It turns out that in the Gromov-Hausdorff distance every compact space is close to a discrete metric space.

**Definition 8.3.** An  $\epsilon$ -net  $L$  is a metric space with the property that  $d(\ell, \ell') \geq \epsilon$  for all  $\ell \neq \ell'$  in  $L$ . An  $\epsilon$ -net in a metric space  $X$  is an isometric image  $L \subset X$  of an  $\epsilon$ -net with the property that every point of  $X$  is within  $\epsilon$  of a point of  $L$ .

Every compact metric space has an  $\epsilon$ -net and any  $\epsilon$ -net in a compact metric space has finite cardinality. It is also clear that the Hausdorff distance between  $X$  and an  $\epsilon$ -net  $L$  in  $X$  is at most  $\epsilon$ . (Let  $Z = X$  with the natural embeddings of  $X$  and  $L$  into  $Z$ .) Thus, a compact metric space  $X$  is the Gromov-Hausdorff limit of any sequence  $L_n \subset X$  of  $\epsilon_n$ -nets in  $X$  provided  $\epsilon_n \rightarrow 0$  as  $n \rightarrow \infty$ . The following is immediate.

**Lemma 8.4.** *Fix  $\epsilon > 0$  and  $N < \infty$ . Suppose that  $L_n$  is a sequence of  $\epsilon$ -nets, with the cardinality of  $L_n$  being at most  $N$  for every  $n$ . Then after passing to a subsequence there is an  $\epsilon$ -net  $L_\infty$  of cardinality at most  $N$  which is the Gromov-Hausdorff limit of the  $L_n$ . Furthermore, for all  $n$  sufficiently large there is a bijection  $L_n \rightarrow L_\infty$  such that the push forwards of the metrics on the  $L_n$  converge uniformly to the limiting metric on  $L_\infty$ .*

One can characterize when a sequence of compact metric spaces of uniformly bounded diameter has a subsequence converging in the Gromov-Hausdorff sense in terms of the cardinalities of nets in the spaces. The following is elementary.

**Corollary 8.5.** *Let  $X_n$  be a sequence of compact metric spaces. Then every subsequence has a further subsequence converging in the Gromov-Hausdorff sense to a compact metric space if and only if for every  $\epsilon > 0$  there is  $N(\epsilon) < \infty$  and for each  $n$  for any  $\epsilon$ -net  $L_n(\epsilon) \subset X_n$  the cardinality of  $L_n(\epsilon)$  is at most  $N(\epsilon)$ .*

**Definition 8.6.** We shall need a based version of the Gromov-Hausdorff distance. Let  $(X, x)$  and  $(Y, y)$  be based, compact metric spaces. We say that the Gromov-Hausdorff distance from  $(X, x)$  to  $(Y, y)$  is the infimum of  $d$  such that there are isometric embeddings  $X, Y \subset Z$  such that  $X$  is in the  $d$ -neighborhood of  $Y$ ,  $Y$  is in the  $d$  neighborhood of  $X$  and  $d(x, y) \leq d$ .

In the based context all  $\epsilon$ -nets are assumed to contain the base point. A sequence  $(X_n, x_n)$  converges in the based Gromov-Hausdorff sense to a compact, based metric space  $(X_\infty, x_\infty)$  if and only if for every  $\epsilon$ -net  $L_\infty$  in  $(X_\infty, x_\infty)$  and a sequence of  $\epsilon'_n$  converging to  $\epsilon$  and  $\epsilon'_n$ -nets  $L_n$  in  $(X_n, x_n)$  and for all  $n$  sufficiently large bijections  $L_n \rightarrow L_\infty$  carrying  $x_n$  to  $x_\infty$  such that under these bijections the metrics on the  $L_n$  converge to the metric on  $L_\infty$ .

**Definition 8.7.** Let  $X$  and  $Y$  be compact metric spaces. A continuous function  $f: X \rightarrow Y$  is an  $\epsilon$ -approximation if there is a metric  $D$  on  $X \amalg Y$  extending the given metrics on  $X$  and  $Y$  such that (i)  $X$  is contained in the  $\epsilon$ -neighborhood of  $Y$ , (ii)  $Y$  is contained in the  $\epsilon$ -neighborhood of  $X$ , and (iii) for all  $y \in Y$  the fiber  $f^{-1}(y)$  is within  $\epsilon$  of  $y$ .

## 8.2 Limits of complete metric spaces

Gromov-Hausdorff convergence works well for compact metric spaces of bounded diameter, but using the same definition for complete metric spaces or more generally for sequences of compact metric spaces with unbounded diameter is much too restrictive. Here is the appropriate generalization to this case.

**Definition 8.8.** Let  $(X_n, x_n)$  be a sequence of based, complete, locally compact metric spaces. We say that  $(X_\infty, x_\infty)$  is the *Gromov-Hausdorff limit* of the  $(X_n, x_n)$  if for every  $R < \infty$  there is a sequence of  $\epsilon_n \rightarrow 0$  such that the closed balls  $(\overline{B(x_n, R + \epsilon_n)}, x_n)$  converge in the based Gromov-Hausdorff sense to  $(\overline{B(x_\infty, R)}, x_\infty)$ .

The results on Gromov-Hausdorff limits for compact metric spaces of bounded diameter immediately generalize in this context.

**Proposition 8.9.** *Let  $(X_n, x_n)$  be a sequence of complete, based metric spaces. Then every subsequence has a further subsequence converging to a complete, based metric space in the Gromov-Hausdorff sense if and only if for each  $\epsilon > 0$  and  $R < \infty$  there is a uniform bound  $N(\epsilon, R)$  to the cardinality of any  $\epsilon$ -net in  $B(x_n, R)$ .*

## 8.3 Manifolds with curvature bounded below

For any  $k \in \mathbb{R}$  we set  $H_k$  equal to the complete, simply connected surface of constant curvature  $k$ . Thus,  $H_k$  is a rescaling of the hyperbolic plane by  $\sqrt{|k|^{-1}}$  if  $k < 0$ , is  $\mathbb{R}^2$  if  $k = 0$ , and is the round sphere of radius  $\sqrt{k^{-1}}$  if  $k > 0$ .

Suppose that  $M$  is a complete, Riemannian manifold with locally convex boundary and with sectional curvature  $\geq k$ . We define the metric on  $M$  in the usual way: for any  $x, y \in M$ , the distance  $d(x, y)$  is the infimum of the lengths of all rectifiable

paths from  $x$  to  $y$ . Because the manifold is complete and the boundary is convex, there is a minimizing geodesic connecting  $x$  to  $y$ , i.e., a geodesic whose length is the distance between the points. This geodesic is an isometric embedding of an interval into  $M$ .

For any triple of points  $a, b, c$  in  $M$  take points  $\tilde{a}, \tilde{b}, \tilde{c}$  in  $H_k$  with the same pairwise distances<sup>5</sup> we define the  $k$ -comparison angle  $\tilde{\angle}_k abc$  to be the angle at  $\tilde{b}$  of the triangle  $\tilde{a}\tilde{b}\tilde{c}$  in  $H_k$ . It is a fundamental result of Toponogov theory ([34]) that the  $k$ -comparison angle at  $\tilde{\angle}_k abc$  is at most the angle in  $M$  between minimal geodesics  $\gamma$  from  $b$  to  $c$  and  $\alpha$  from  $b$  to  $a$ . Even more, as we move the point  $a$  along  $\alpha$  toward  $b$  and keep  $c$  fixed the  $k$ -comparison angle is a weakly monotone increasing function. From this we deduce:

**Lemma 8.10.** *Let  $M$  be a complete Riemannian manifold with locally convex boundary and with sectional curvatures  $\geq k$ . Let  $x; a, b, c$  be four distinct points in  $X$ . Then*

$$\tilde{\angle}_k axb + \tilde{\angle}_k bxc + \tilde{\angle}_k cxa \leq 2\pi.$$

*Proof.* Since the  $k$ -comparison angles are at most the angles between minimal geodesics to  $x$ , we need only see that given three geodesics emanating from  $x$  the sum of the 3 angles between them is at most  $2\pi$ . This is clear.  $\square$

**Lemma 8.11.** *There is a constant  $c_R = c_R(k, n)$  depending only on  $k$ , the dimension  $n$ , and a radius  $R$ , such that for any complete Riemannian  $n$ -manifold with locally convex boundary and with sectional curvature  $\geq k$  and any ball  $B$  of radius  $R$  in  $M$  the cardinality of any  $\epsilon$ -net in  $B$  is at most  $c_R(k, n)\epsilon^{-n}$ .*

*Proof.* We begin the proof with an elementary claim, whose proof we leave to the reader.

**Claim 8.12.** *There is a constant  $c(k, R) > 0$  such that for any triangle  $\tilde{a}\tilde{b}\tilde{c}$  in  $H_k$  with both  $|\tilde{a}\tilde{b}|$  and  $|\tilde{c}\tilde{b}|$  bounded above by  $R$ , and with  $|\tilde{a}\tilde{c}| \geq 2\left||\tilde{a}\tilde{b}| - |\tilde{c}\tilde{b}|\right|$  we have*

$$\tilde{\angle} \tilde{a}\tilde{b}\tilde{c} \geq c(k, R)d(\tilde{a}, \tilde{c}).$$

Also, for each  $n \geq 2$ , there is a constant  $d(n)$  such that for all  $\delta > 0$  there are at most  $d(n)\delta^{1-n}$  disjoint balls of radius  $\delta$  in  $S^{n-1}$  with the round metric of constant curvature 1.

Fix a complete Riemannian manifold of dimension  $n$  with locally convex boundary and with sectional curvature  $\geq k$ , and let  $B$  be a ball in  $M$  of radius  $R$  and center  $x$ . Now consider an  $\epsilon$ -net  $L \subset B$ . We divide  $B$  into  $N = \lceil 2R/\epsilon \rceil + 1$  disjoint annular rings  $A_1, \dots, A_N$  each of width  $\leq \epsilon/2$  and we consider  $L_i = L \cap A_i$ . For any  $\ell \neq \ell' \in L_i$ , the above claim implies that the comparison angle  $\tilde{\angle} \ell x \ell'$  is at least  $c(k, R)\epsilon/2$ . Thus, the angle at  $x$  between minimal geodesics from  $\ell$  and  $\ell'$  to  $x$  is at least  $c(k, R)\epsilon/2$ .

<sup>5</sup>If  $k > 0$ , then we require  $d(z, b) + d(b, c) + d(c, a) \leq 2\pi/\sqrt{k}$ . This will always be implicitly assumed.

Thus, there can be at most  $2^{n-1}d(n)c(k, R)^{1-n}\epsilon^{1-n}$  such points. Summing over all the annuli, we see that the cardinality of  $L$  is at most

$$2^{n-1}d(n)c(k, R)^{1-n}[2R + 1]\epsilon^{-n}.$$

This establishes the result.  $\square$

As a consequence, we have

**Corollary 8.13.** *Given a sequence of based, complete Riemannian manifolds of dimension  $n$  with locally convex boundary and with sectional curvature  $\geq k$ , there is a subsequence that converges in the Gromov-Hausdorff sense to a complete metric space.*

Let us examine some of the properties of this limiting metric space. The first involves the notion that arose in establishing the bounds on the cardinalities of lattices in balls.

**Definition 8.14.** Let  $X$  be a compact metric space. The  $n$ -dimensional rough volume of  $X$ , denoted  $Vr_n(X)$ , is defined as

$$\lim_{\epsilon \rightarrow 0} \beta_\epsilon(X) \epsilon^n,$$

where  $\beta_\epsilon(X)$  is the maximal cardinality of any  $\epsilon$ -net in  $X$ .

Notice that if  $X$  is a compact metric space then there is a unique  $d \in [0, \infty]$  such that  $Vr_n(X) = \infty$  for  $0 \leq n < d$  and  $Vr_n(X) = 0$  for  $d < n \leq \infty$ . The constant  $d$  is the *rough dimension* of  $X$ . It follows from the above that for a compact subset with non-empty interior in a complete Riemannian manifold with locally convex boundary and with sectional curvature  $\geq k$  its rough dimension is equal to its topological dimension.

An upper bound on rough dimension passes to Gromov-Hausdorff limits.

**Corollary 8.15.** *Let  $(X, x)$  be the Gromov-Hausdorff limit of a sequence of based, complete Riemannian  $n$ -manifolds with locally convex boundary and with sectional curvatures  $\geq k$ . Then any compact subset of  $X$  has rough dimension at most  $n$ .*

A second condition that passes to Gromov-Hausdorff limits is the fact that any two points are connected by a rectifiable path which is an isometric embedding of an interval into the space, and in particular whose length is equal to the distance between the endpoints. Such metric spaces are called *length spaces*. Notice that in a length space the Gromov-Hausdorff distance from  $B(x, R)$  and  $B(x, R')$  is at most  $|R - R'|$ .

**Lemma 8.16.** *The Gromov-Hausdorff limit of a sequence of based, complete Riemannian manifolds of dimension  $n$  with locally convex boundary and with sectional curvature  $\geq k$  is a length space.*



*Proof.* Let  $(X, x)$  be the limit of  $\{(M_i, p_i)\}$  and let  $y \neq z$  be points of  $X$  of distance  $d$  apart. Take sequences  $y_i, z_i \in M_i$  converging (in some fixed realization) to  $y$  and  $z$ , and let  $\gamma_i$  be a minimal geodesic in  $M_i$  from  $y_i$  to  $z_i$ , parametrized at unit speed by the interval  $[0, d_i]$ . Passing to a subsequence, we can arrange that there is a countable dense subset  $S$  of  $[0, d]$  such that the  $\gamma_i(s)$  converges to a point  $\gamma(s)$  of  $X$  for all  $s \in S$ . The completion of the set of these images is the required interval connecting  $y$  and  $z$ .  $\square$

The other condition that passes to limits is related to comparison angles. Let us first formulate the condition on Riemannian manifolds.

This leads to the following definition:

**Definition 8.17.** Let  $X$  be a metric space. We say that it has rough curvature  $\geq k$  if for every four points  $x; a, b, c$  the  $k$ -comparison angles satisfy

$$\tilde{Z}_k xab + \tilde{Z}_k bxc + \tilde{Z}_k cxa \leq 2\pi.$$

**Theorem 8.18.** Let  $(X, x)$  be the Gromov-Hausdorff limit of a sequence  $\{(M_i, p_i)\}$  of complete Riemannian  $n$ -manifolds with locally convex boundary and with sectional curvature  $\geq k$ . Then  $X$  is a length space whose rough dimension is  $\leq n$  with rough curvature  $\geq k$ .

This leads to the following definition:

**Definition 8.19.** An Alexandrov space of curvature  $\geq k$  is a complete length space of rough curvature  $\geq k$ . An Alexandrov space is an Alexandrov space of curvature  $\geq k$  for some  $k > -\infty$ . The *dimension* of an Alexandrov space is its rough dimension. A *geodesic* in an Alexandrov space is an isometric embedding of an interval into the Alexandrov space. We use this notion exclusively from now on, even when the Alexandrov space is a Riemannian manifold (and there is another notion of geodesics.)

We have shown:

**Corollary 8.20.** A sequence of complete Riemannian  $n$ -manifolds with locally convex boundary of sectional curvature  $\geq k$  has a subsequence with a Gromov-Hausdorff limit. Any Gromov-Hausdorff limit of such a sequence of manifolds is an Alexandrov space of dimension  $\leq n$  and curvature  $\geq k$ .

## 9 Basics of Alexandrov spaces

It is important to have results not just for Riemannian manifolds with curvature bounded below but also for Alexandrov spaces, so we translate the results above into results for Alexandrov spaces.

### 9.1 Properties of comparison angles

The condition on the comparison angles in Definition 8.17 is equivalent to other conditions on angles.

**Lemma 9.1.** *Suppose that  $X$  is a complete Alexandrov space with curvature  $\geq k$ . Let  $\gamma$  and  $\nu$  be geodesics (i.e., isometric embeddings of intervals) in  $X$  which begin at the same point  $x$ . Let the other endpoint of  $\gamma$ , resp.  $\nu$ , be  $y$ , resp.  $z$ , and let  $d_1$  and  $d_2$  be the lengths of  $\gamma$  and  $\nu$ . Then for any  $0 < s \leq d_1$  and  $0 < t \leq d_2$  denote by  $\gamma(s)$ , resp.  $\nu(t)$ , the point along  $\gamma$ , resp.  $\nu$ , at distance  $s$ , resp.  $t$ , from  $x$ . Then the comparison angle*

$$\tilde{\angle}_k \gamma(s) x \nu(t)$$

*is a weakly monotone decreasing function of either variable  $s, t$  when the other is held fixed. Also, for any  $0 < s < d_1$  the distance from  $z$  to  $\gamma(s)$  is at least as large as the corresponding distance in the comparison triangle in  $H_k$ .*

*Proof.* By symmetry in order to prove the first statement it suffices to take  $t = d_2$  and  $s < d_1$  and show that

$$\tilde{\angle}_k \gamma(s) x z \leq \tilde{\angle}_k y x z.$$

Applying the defining inequality to  $\{\gamma(s); x, y, z\}$ , yields  $\tilde{\angle}_k z \gamma(s) y + \tilde{\angle}_k z \gamma(s) x \leq \pi$ . (The fact that  $\gamma$  is a geodesic implies that  $\angle_k x \gamma(s) y = \pi$ .) This implies that  $d(z, \gamma(s))$  is at least as large as the distance in  $H_k$  between  $\tilde{\gamma}(s)$  and  $\tilde{z}$ , where  $\tilde{\gamma}$  is the geodesic in  $H_k$  from  $\tilde{x}$  to  $\tilde{y}$  and  $\tilde{\gamma}(s)$  is the point on this geodesic at distance  $s$  from  $\tilde{x}$ . But this implies that  $\tilde{\angle}_k \gamma(s) x z \geq \tilde{\angle}_k y x z$ , as claimed, as well as establishing the second statement in the lemma.  $\square$

Since all comparison angles are bounded above by  $\pi$ , it follows that there is a limit as  $s$  and  $t$  tend to zero of  $\tilde{\angle}_k \gamma(s) y \nu(t)$  which is called *the angle between  $\gamma$  and  $\nu$  at  $x$*  and is denoted  $\angle_k \gamma \nu$ . If the Alexandrov space is a Riemannian manifold then the angle between geodesics in the Alexandrov sense is the usual Riemannian angle between the geodesics.

The defining property of an Alexandrov space leads easily to unique extension of geodesics.

**Lemma 9.2.** *Let  $\gamma$  be a geodesic from  $x$  of positive length in an Alexandrov space. If  $\mu$  and  $\mu'$  are geodesics from  $x$  to points  $z$  and  $z'$  with  $\gamma \subset \mu \cap \mu'$  then either  $\mu \subset \mu'$  or  $\mu' \subset \mu$ .*

**Corollary 9.3.** *If  $\gamma$  is a geodesic from  $x$  to  $y$  and  $z$  is an interior point of  $\gamma$ , then there is a unique geodesic from  $x$  to  $z$ , namely the sub-geodesic of  $\gamma$  with endpoints  $x$  and  $z$ .*

**Lemma 9.4.** *Suppose that sequences of geodesics  $\alpha_n, \beta_n$  emanating from  $x_n$  converge to geodesics  $\alpha$  and  $\beta$  emanating from  $x$ . Then*

$$\liminf_{n \rightarrow \infty} \angle_k \alpha_n \beta_n \geq \angle_k \alpha \beta.$$

*Proof.* For any  $\epsilon > 0$  there are points  $y \in \alpha$  and  $z \in \beta$  such that  $\tilde{\angle}_k yxz = a \geq (\angle_k \alpha\beta) - \epsilon$ . By the convergence property there are  $y_n \in \alpha_n$  and  $z_n \in \beta_n$  converging to  $y$  and  $z$ . Thus,  $\tilde{\angle}_k y_n x_n z_n$  converges to  $a$  and hence by monotonicity the angle between  $\alpha_n$  and  $\beta_n$  at  $x_n$  is at least  $a - \epsilon$  for all  $n$  sufficiently large. Since this is true for every  $\epsilon > 0$ , this proves the result.  $\square$

There is a related fact for smooth manifolds:

**Lemma 9.5.** *Let  $M$  be a smooth Riemannian manifold with curvature  $\geq k$  and for any  $y \in M$  denote by  $S_y(M)$  the tangent sphere to  $M$  at  $y$ . Suppose that  $A \subset M$  is a compact set, and let  $U$  denote the complement of  $A$  in  $M$ . Then for each  $y \in U$  denote by  $A'_y \subset S_y(M)$  be the subset consisting of all tangent directions at  $y$  to geodesics (i.e. minimal geodesics) from  $y$  to  $A$ . This is a compact subset of  $S_y(M)$ . Then the function on  $TM|_U \rightarrow \mathbb{R}$  that associates to a unit tangent vector  $\tau$  at  $y$  the distance in  $S_y(M)$  of  $\tau$  to  $A'_y$  is lower semi-continuous.*

*Proof.* Suppose that  $\tau_n$  is a unit tangent vector at  $y_n \in U$ , the  $y_n$  converge to  $y$  and the  $\tau_n$  converge to  $\tau$ , a unit tangent vector at  $y$ . Let  $d_n$  be the distance in  $S_{y_n}(M)$  from  $\tau_n$  to  $A'_{y_n}$ . Passing to a subsequence we can suppose that the  $d_n$  converge to a limit  $d$ . We must show that  $d$  is greater than or equal to the distance from  $\tau$  to  $A'_y$ . For each  $n$  there is a geodesic  $\gamma_n$  from  $y$  to  $A$  whose tangent vector at  $y$  is distance  $d_n$  from  $\tau_n$ . Passing to a further subsequence, we can suppose that the  $\gamma_n$  converge to a geodesic  $\gamma$  from  $y$  to  $A$ . The tangent  $a$  to  $\gamma$  at  $y$  has the property that the distance from  $\tau$  to  $a$  is  $d$ . On the other hand,  $a \in A'_y$  so that  $d$  is greater than or equal to the distance from  $A'_y$  to  $\tau$ .  $\square$

The following is an elementary exercise.

**Lemma 9.6.** *For any  $\epsilon > 0$  there is  $\beta > 0$  such that the following holds. Suppose that we have three points  $a, b, c$  in an Alexandrov space of curvature  $\geq -1$  and suppose that  $d(a, b), d(b, c)$  are each between  $1/10$  and  $1$  suppose furthermore that  $d(a, b) + d(b, c) \leq d(a, c) + \beta$ . Then the comparison angle  $\tilde{\angle}_k abc \geq (1 - \epsilon)\pi$ .*

### 9.1.1 Effect of scaling

Let  $X$  be an Alexandrov space of curvature  $\geq k$ . The rescaled metric space  $rX$  is an Alexandrov space of curvature  $\geq r^{-2}k$ , and for any  $x, y, z \in X$  the  $k$ -comparison angles in  $X$  agree with the  $r^{-2}k$ -comparison angles in  $rX$ . As we rescale we always implicitly rescale the lower bound for the curvature.

**Claim 9.7.** *Let  $(X, x)$  be a based Alexandrov space of curvature  $\geq k$ . Suppose that  $r_n$  is a sequence of positive constants converging to 0, then after passing to a subsequence, the based Alexandrov spaces  $r_n^{-1}(X, x)$  converge to a limit  $(Y, y)$  that is a based Alexandrov space of curvature  $\geq 0$ . Furthermore, under this convergence the comparison angles also converge when the comparison angles in  $r_n^{-1}X$  are  $\angle_{r_n^2 k}$ .*

From now on we simplify the notation by dropping the  $k$  from the notation for comparison angles since  $k$  will always be clear from the context.

## 9.2 The Product Theorem for Alexandrov Spaces of Curvature $\geq 0$

**Theorem 9.8.** *Suppose that  $X$  is a complete Alexandrov space of dimension  $n$  and of curvature  $\geq 0$  and that  $\gamma$  is an isometric embedding of  $\mathbb{R}$  into  $X$ . Then there is a complete Alexandrov space  $Y$  of dimension  $n - 1$  and of curvature  $\geq 0$  and an isometry  $Y \times \mathbb{R} \cong X$  in such a way that  $\gamma$  is the image of  $\{y_0\} \times \mathbb{R}$  for some point  $y_0 \in Y$ .*

*Proof.* Let  $\gamma^\pm$  be the opposite geodesic rays in  $\gamma$  with endpoint  $x \in X$ . Consider sequences  $\{x_{n,-}\}$  and  $\{x_{n,+}\}$ , equidistant from  $x$ , tending to the two ends of  $\gamma$ , (with  $x_{n,+} \in \gamma^+$ ). For any  $y \in X$  consider the comparison angle  $\tilde{\angle} x_{n,-}yx_{n,+}$ . Since  $d(x_{n,+}, y)$  and  $d(x_{n,-}, y)$  tend to  $\infty$  and  $d(x_{n,+}, y) + d(x_{n,-}, y) - d(x_{n,+}, x_{n,-})$  is bounded above by  $2d(x, y)$ , it follows that the comparison angles converge  $\pi$  as  $n \rightarrow \infty$ . This means that, possibly after passing to a subsequence, the geodesics  $\mu_{n,\pm}$  from  $y$  to  $x_{n,\pm}$  converge to geodesics  $\gamma_y^\pm$  whose union is a geodesic line  $\gamma_y$  in  $X$  (i.e., a geodesic embedding  $\mathbb{R} \subset X$ ) passing through  $y$ . In this way we construct for each  $y \in X$  an isometric embedding of  $\mathbb{R} \rightarrow X$  passing through  $y$  parallel, in some sense, to  $\gamma$ . The end of  $\gamma_y$  determined by  $\gamma_y^+$  is called the  $+$ -end and the other end is the  $-$  end.

**Claim 9.9.** *For any choice of sequences  $x_{n,\pm}$  tending to infinity in  $\gamma^\pm$  and any geodesics  $\mu_{n,\pm}$  from  $y$  to  $x_{n,\pm}$  there are limiting geodesic rays  $\gamma_y^\pm$  whose union is an isometric copy of  $\mathbb{R}$  in  $X$  passing through  $y$ . This isometric copy of  $\mathbb{R}$  is independent of the choice of the sequences  $x_{n,\pm} \subset \gamma$  tending to the  $\pm$ -end of  $\gamma$  and of the geodesics  $\mu_{n,\pm}$ . Furthermore, for any  $y' \in \gamma_y$  we have  $\gamma_{y'} = \gamma_y$ .*

*Proof.* Fix a sequence in  $\gamma$  going to  $\infty$  in the negative direction and geodesics from  $y$  to these points with a limiting geodesic ray  $\gamma_y^-$  beginning at  $y$  and consider two sequences in  $\gamma$  going to infinity in the positive direction and geodesics from  $y$  to these points with limiting geodesic rays. Each of these rays completes  $\gamma_y^-$  to a complete geodesic, and hence by the unique continuation of geodesics these limiting geodesic rays in the positive direction are equal. The symmetric argument shows all limiting geodesic rays from  $y$  in the negative direction are identical. This proves the first statement.

Suppose that  $y_n \mapsto y$  and  $\mu_{n,\pm}$  are geodesics connecting  $x_{n,\pm}$  to  $y_n$ . Again the comparison angles converge to  $\pi$  so that, after passing to a subsequence, these geodesics converge to a geodesic copy of  $\mathbb{R}$  passing through  $y$ . The above argument proves that this copy of  $\mathbb{R}$  is  $\gamma_y$ . Now suppose that  $y' \in \gamma_y$  and has distance  $d$  from  $y$ . By symmetry we can suppose that  $y'$  is further toward the positive end of  $\gamma_y$  than  $y$ . Let  $\mu_n$  be a geodesic from  $x_{n,+}$  to  $y$ , let  $x'_n$  be the point on  $\mu_n$  at distance  $d$  from  $y$ , and let  $\mu_{n,0}$  be the subgeodesic of  $\mu_n$  with endpoints  $x_{n,+}$  and  $x'_n$ . By the above the  $x'_n$  converge to  $y'$  and the geodesics  $\mu_{n,0}$  converge to the geodesic ray in  $\gamma_{y'}$  emanating from  $y'$  in the positive direction. On the other hand, the  $\mu_{n,0}$  converge to the geodesic sub-ray of  $\gamma_y$  emanating from  $y'$  in the positive direction. This proves that  $\gamma_y = \gamma_{y'}$ , proving the second assertion in the claim.  $\square$

This means that given the isometric copy  $\gamma$  of  $\mathbb{R}$  in  $X$  we have a well-defined foliation of  $X$  by geodesics of the form  $\gamma_y$  for  $y \in X$ . We denote this foliation by  $\mathcal{F}(\gamma)$ . Now let us establish a strong notion of parallelism among the geodesics in  $\mathcal{F}(\gamma)$ .

**Claim 9.10.** *Suppose that  $y', y'' \in \gamma_y$ . Then the distance from  $y'$  to  $\gamma$  is equal to the distance from  $y''$  to  $\gamma$ .*

*Proof.* By symmetry it suffices to show that the distance from  $y'$  to  $\gamma$  is greater than or equal to the distance from  $y''$  to  $\gamma$ . Let  $d$  be the distance from  $y'$  to  $y''$  and, by symmetry we can suppose that  $y''$  lies closer to the  $+$ -end of  $\gamma_y$ . For each  $n$  sufficiently large we take a geodesic  $\mu'_n$  from  $x_{n,+}$  to  $y'$  and we set  $z''_n$  equal to the point at distance  $d$  from  $y'$  on this geodesic. As  $n \mapsto \infty$  the points  $z''_n$  converge to  $y''_n$ . Let  $\tilde{D}_n$  be the length of  $\mu'_n$ . Fix a point  $x''_n$  on  $\gamma$  closest to  $z''_n$ , let  $D_n$  be the distance from  $x_n$  to  $x''_n$ , set  $d' = dD_n/\tilde{D}_n$  and let  $x'_n$  be point at distance  $d'$  from  $x''_n$  along  $\gamma$  toward the negative end. The distance from  $x'_n$  to  $y'$  is bounded independent of  $n$  and hence passing to a subsequence we can suppose that  $x'_n$  converge to a point  $x' \in \gamma$ . Construct the planar comparison triangle  $\tilde{y}'\tilde{x}_{n,+}\tilde{x}'_n$  and let  $\tilde{z}''_n$ , resp.  $\tilde{x}''_n$ , be the point along the side  $\tilde{y}'\tilde{x}_{n,+}$ , resp.  $\tilde{x}'_n\tilde{x}_{n,+}$ , at distance  $d$ , resp.  $d'$ , from  $\tilde{y}'$ , resp.  $\tilde{x}'_n$ . Then by of planar triangles  $|\tilde{z}''_n\tilde{x}''_n| = \frac{\tilde{D}_n-d}{\tilde{D}_n}|\tilde{y}'\tilde{x}'_n|$  and by the fundamental comparison result for Alexandrov spaces we have  $|z''_n x''_n| \geq |\tilde{z}''_n \tilde{x}''_n|$ . Thus, in the limit as  $n \rightarrow \infty$  we have that the distance from  $y''$  to  $\gamma$  is  $\geq d(y', x')$ , which in turn is greater than or equal to the distance from  $y'$  to  $\gamma$ . This completes the proof of the claim.  $\square$

**Claim 9.11.** *Given  $y \in X$  there is a constant  $C < \infty$  such that the distance from any  $x' \in \gamma$  to  $\gamma_y$  is at most  $C$ .*

*Proof.* Take a sequence of points  $\{y_n\}_{n=-\infty}^{\infty}$  equally spaced at distance 1 along  $\gamma_y$  and for each  $n$  let  $x_n \in \gamma$  be a closest point on  $\gamma$  to  $y_n$ . Then the distance from  $x_n$  to  $x_{n+1}$  is at most  $2d + 1$ , where  $d$  is the distance from any point of  $\gamma_y$  to  $\gamma$ . It follows that as  $n \rightarrow \pm\infty$  the  $x_n$  converge to the  $\pm$ -end of  $\gamma$ . Hence given any  $x' \in \gamma$  there  $n$  such that  $d(x', x_n) \leq 2d + 1$ , and hence the distance from  $x'$  to  $\gamma_y$  is bounded above by  $3d + 1$ .  $\square$

**Corollary 9.12.** *Let  $y_{n,\pm}$  be a sequence of points converging to the plus and minus ends of  $\gamma_y$ . Then geodesic arcs  $\mu_{n,\pm}$  from  $y_{n,\pm}$  to  $x$  converge to opposite geodesic rays on  $\gamma$ , in particular  $\gamma$  is an element of  $\mathcal{F}(\gamma_y)$ . More generally,  $\mathcal{F}(\gamma) = \mathcal{F}(\gamma_y)$ .*

*Proof.* Since there is a constant  $C < \infty$  such that every  $x \in \gamma$  is within distance  $C$  of  $\gamma_y$ , it follows the comparison angle between  $\mu_{n,+}$  and  $\gamma^-$  tends to  $\pi$ , and similarly with  $+$  and  $-$  reversed. This proves the first statement. Now we see that for any element  $\gamma'$  of  $\mathcal{F}(\gamma)$  there is a constant  $C'$  depending only on  $\gamma'$  such that every point of  $\gamma'$  is within a distance  $C'$  of  $\gamma_y$  and hence by the same argument it follows that  $\gamma'$  is also an element of  $\mathcal{F}(\gamma_y)$ .  $\square$

It follows that for any two elements  $\gamma_1, \gamma_2$  of  $\mathcal{F}(\gamma)$  there is a distance  $d$  such that every point  $x_1 \in \gamma_1$  is exactly distance  $d$  from  $\gamma_2$ ; that is to say any two elements of  $\mathcal{F}(\gamma)$  are parallel in the sense that they are constant distance apart.

Now we define a function  $f^+$  by  $f^+(y) = \lim_{n \rightarrow \infty} d(x_{n,+}, y) - d(x_{n,+}, x)$  and similarly we define  $f^-$  using the points  $x_{n,-}$  instead of  $x_{n,+}$ . By the usual argument, limits of this type are affine linear on geodesics in flat space. By the comparison property this implies that  $f^\pm$  are convex on any geodesic in  $X$ , meaning that if  $\mu$  is a geodesic arc with endpoints  $a, b$  and  $c$  is a point on the arc such that  $d(b, c)/d(a, b) = t$ , then  $f^+(c) \geq tf^+(a) + (1-t)f^+(b)$ , and analogously for  $f^-$ . Thus,  $f^+ + f^-$  is a convex function on each geodesic and clearly  $f^+ + f^- \geq 0$  everywhere. Of course,  $f^+ + f^-$  is identically zero along  $\gamma$ .

**Proposition 9.13.**  *$f^+ + f^-$  is identically zero and  $f^+$  is affine linear on each geodesic.*

*Proof.* For each  $n$  let  $y_n \in \gamma_y$  be the point equidistant from  $x_{n,+}$  and  $x_{n,-}$ . We claim that after passing to a subsequence we can arrange that the  $y_n$  converge to a point  $y_0 \in \gamma_y$ . Let  $x_n$  be a closest point on  $\gamma$  to  $y_n$ . Then the difference  $d(x_{n,+}, x_n) - d(x_{n,-}, x_n)$  is bounded by twice the distance from  $x_n$  to  $\gamma$ , and hence, by the previous claim, this difference is bounded independent of  $n$ . It then follows that the  $x_n$  are within a bounded distance of  $x$  and hence so are the  $y_n$ . Thus, the  $y_n$  have a subsequence converging to  $y_0 \in \gamma_y$ .

**Claim 9.14.** *Let  $y_0 \in \gamma_y$  be the limit of a subsequence of points  $y_n \in \gamma_y$  equidistant from the  $x_{n,+}$  and  $x_{n,-}$ . Then  $f^+(y_0) = f^-(y_0) = 0$  and  $x$  is the unique closest point of  $\gamma$  to  $y_0$ .*

*Proof.* Let  $\tilde{D}_n = d(x_{n,+}, y_0)$ . Since  $2\tilde{D}_n = d(x_{n,+}, y_0) + d(x_{n,-}, y_0) \geq d(x_{n,+}, x_{n,-}) = 2d(x_{n,+}, x)$ , we have  $\tilde{D}_n \geq d(x_{n,\pm}, x)$ . Taking limits we see that

$$\lim_{n \rightarrow \infty} (d(x_{n,\pm}, y_0) - d(x_{n,\pm}, x)) \geq 0. \quad (9.1)$$

On the other hand, let  $x_0 \in \gamma$  be a closest point of  $\gamma$  to  $y_0$ , and consider all geodesics from  $y_0$  to  $x_0$ . If any one of these geodesics makes an angle at  $x_0$  less than  $\pi/2$  with one of the directions along  $\gamma$ , then, since angles between geodesics are greater than the comparison angles, the point  $x_0$  is not a closest point on  $\gamma$  to  $y_0$ . Thus, any geodesic from  $y_0$  to  $x_0$  makes angle at least  $\pi/2$  at  $x_0$  with both directions along  $\gamma$ . Since the sum of the angles at  $x_0$  to the two directions along  $\gamma$  is at most  $\pi$ , it follows that the angle at  $x_0$  between any geodesic from  $y_0$  to  $x_0$  and each direction along  $\gamma$  is  $\pi/2$ . This means that for every  $n$  the comparison angle  $\tilde{\angle}_{y_0 x_0 x_{n,\pm}}$  is at most  $\pi/2$ . Using comparison triangles we see that

$$\lim_{n \rightarrow \infty} (d(x_{n,\pm}, y_0) - d(x_{n,\pm}, x_0)) \leq 0. \quad (9.2)$$

By symmetry we can suppose that  $x_0$  lies in  $\gamma^+$  so that  $d(x_{n,+}, x_0) \leq d(x_{n,+}, x)$ . The only way that Inequalities 9.1 and 9.2 are consistent with this is if both those inequalities are equalities and in addition  $x = x_0$ . Equality in Inequality 9.1 means that  $f^+(y_0) = f^-(y_0) = 0$ .  $\square$

Since  $f^+ + f^- \geq 0$  and is convex on  $\gamma_y$ , the fact that it is zero at  $y_0 \in \gamma_y$  implies that it is identically zero on  $\gamma_y$ . Since this is true for every element of the foliation  $\mathcal{F}(\gamma)$ , we see that  $f^+ + f^- = 0$ , and hence  $f^+$  is both concave and convex on each geodesic. Consequently,  $f^+$  is affine linear on each geodesic. This completes the proof of the proposition.  $\square$

A similar argument shows that given any  $y \in X$  any closest point on  $\gamma$  to  $y$  is the unique point of  $x' \in \gamma$  with  $f^+(y) = f^+(x')$ . Also, notice that this argument implies that if  $y' \in X \setminus \gamma$ , if  $x' \in \gamma$ , and if  $f^+(y') = f^+(x')$ , then  $\lim_{n \rightarrow \infty} \tilde{\angle} y' x' x_{n,\pm} = \pi/2$ . Since the comparison angles are monotone increasing as we move in along  $\gamma$  toward  $x'$ , it follows that for any  $x'' \in \gamma$ , distinct from  $x'$ , we have  $\tilde{\angle} y' x' x'' = \pi/2$ . Of course, there is nothing distinguished about  $\gamma$  so in fact given  $a \neq b \in X$  with  $f^+(a) = f^+(b)$  for any  $c \in \gamma_b$  distinct from  $b$  we have  $\tilde{\angle} abc = \pi/2$ . This proves:

**Corollary 9.15.** *Let  $a, b$  be distinct points of  $X$  with  $f^+(a) = f^+(b)$  and let  $c \in \gamma_b$  be a point distinct from  $b$ . Then  $\tilde{\angle} abc = \pi/2$ .*

Now we consider the fibers of  $Y_t = (f^+)^{-1}(t)$  for  $t \in \mathbb{R}$ . Since  $f^+$  is affine linear on each geodesic, for each  $t$  the fiber  $Y_t$  is geodesically convex: any geodesic in  $X$  with endpoints in  $Y_t$  lies completely in  $Y_t$ . Also, for each  $t \in \mathbb{R}$ ,  $Y_t$  is a complete metric space since  $X$  is a complete. Hence, for each  $t \in \mathbb{R}$ , the fiber  $Y_t$  is a complete Alexandrov space of curvature  $\geq 0$  and of dimension one less than the dimension of  $X$ . Of course,  $Y_t$  meets each geodesic in  $\mathcal{F}(\gamma)$  in exactly one point. Thus, for each  $t \in \mathbb{R}$ , flowing along the leaves of the foliation  $\mathcal{F}(\gamma)$  defines an identification of  $Y_t$  with  $Y_0$ .

**Claim 9.16.** (i) *For each  $t \in \mathbb{R}$  the identification  $Y_t$  with  $Y_0$  given by flowing along the leaves of  $\mathcal{F}(\gamma)$  is an isometry.*

(ii) *Given  $t, t'$  and  $a \in Y_t$  the distance from  $a$  to  $Y_{t'}$  is  $|t' - t|$  and the unique closest point of  $Y_{t'}$  to  $a$  is the intersection of  $\gamma_a \cap Y_{t'}$ .*

*Proof.* Let  $a, b \in Y_t$ . Let  $\gamma_a$  and  $\gamma_b$  be the elements of  $\mathcal{F}(\gamma)$  through  $a$  and  $b$ , and let  $a_0$  and  $b_0$  be the intersections of these geodesics with  $Y_0$ . Then the distance between  $a$  and  $b$  is also the distance between  $a_0$  and  $b_0$  and the unique closest point of  $\gamma_b$  to  $a_0$  is  $b_0$ . This proves the first statement.

For the second, note that  $(f^+)'$  has norm 1 and for every  $b$  the only directions  $\tau \in S_b(X)$  with  $(f^+)'(\tau) = \pm 1$  are the two directions along  $\gamma_b$ . Since  $f^+(Y_{t'}) - f^+(a) = t' - t$ , it follows that any geodesic from  $a$  to  $Y_{t'}$  has length  $\geq |t' - t|$  and the length is strictly greater than  $|t' - t|$  unless the geodesic lies in  $\gamma_a$ . The second result follows.  $\square$

We endow  $Y_0 \times \mathbb{R}$  with the product metric:

$$d((y, t), (y', t')) = \sqrt{d_{Y_0}(y, y')^2 + (t - t')^2}.$$

We define a map  $\Phi: Y_0 \times \mathbb{R} \rightarrow X$  by sending  $(y, t)$  to the unique point on  $\gamma_y \cap (f^+)^{-1}(t)$ . We claim that  $\Phi$  is an isometry. Clearly it is a homeomorphism and for each  $t \in \mathbb{R}$  it is an isometry from  $Y_0 \times \{t\}$  onto  $Y_t$ . Let us consider the distance

between  $a = (y, t)$  and  $c = (y', t')$  for  $t \neq t'$ . Let  $b = (y', t)$ . Since  $b$  and  $c$  lie on the same element of  $\mathcal{F}(\gamma)$  and  $f^+(a) = f^+(b)$ , it follows from Corollary 9.15 that  $\tilde{\angle}abc = \pi/2$ , which means that

$$d(a, c) = \sqrt{d_{Y_t}(y, y')^2 + (t - t')^2}.$$

Of course, we already have established that  $d_{Y_t} = d_{Y_0}$ . This proves that  $\Phi$  is an isometry.  $\square$

**Corollary 9.17.** *Suppose that  $X$  is an Alexandrov space of curvature  $\geq 0$  containing an isometric copy of  $\mathbb{R}^m$  for some  $m > 0$ . Then there is an Alexandrov space  $Y$  and an isometric product decomposition  $X = \mathbb{R}^m \times Y$  with the property that the given copy of  $\mathbb{R}^m$  is identified with  $\mathbb{R}^m \times \{y_0\}$  for some  $y_0 \in Y$ .*

### 9.3 Strainers

A crucial concept for Alexandrov spaces is that of a strainer<sup>6</sup>. Let  $X$  be an Alexandrov space of curvature  $\geq k$ . Fix  $\delta > 0$ . A  $(n, \delta)$ -strainer at a point  $x \in X$  is a set  $\{a_1, b_1, \dots, a_n, b_n\}$  such that:

1.  $\tilde{\angle}a_i x a_j \geq \pi/2 - \delta$  for all  $i \neq j$ .
2.  $\tilde{\angle}b_i x b_j \geq \pi/2 - \delta$  for all  $i \neq j$ .
3.  $\tilde{\angle}a_i x b_j \geq \pi/2 - \delta$  for all  $i \neq j$ .
4.  $\tilde{\angle}a_i x b_i \geq \pi - \delta$  for all  $i$ .

The size of an  $(n, \delta)$ -strainer is the minimum of the  $2n$  distances  $\{d(x, a_i), d(x, b_i)\}_{i=1}^n$ .

Notice that it follows from the defining property that all the angles in the first 3 items are  $\leq \pi/2 + 2\delta$ . We say that an Alexandrov space  $X$  has *strainer dimension*  $n$  at  $x \in X$  if:

- for every neighborhood  $U$  of  $x$  and every  $\delta > 0$ ,  $X$  there is an  $(n, \delta)$ -strainer at some point of  $U$ , and
- there is a  $\delta_0 > 0$  and a neighborhood  $U_0$  of  $x$  so that no point of  $U_0$  has an  $(n + 1, \delta_0)$ -strainer.

The following two results are elementary and are proved using the defining property of comparison angles and Lemma 9.1, see Theorem 9.4 of [3].

**Lemma 9.18.** *Given  $n$ , the following holds for all  $\delta > 0$  sufficiently small.*

- Suppose that  $x \in X$  has an  $(n, \delta)$ -strainer  $\{a_1, b_1, \dots, a_n, b_n\}$  of size  $s$  and that the strainer dimension of  $X$  at  $x$  is  $n$ . Then there is a constant  $r > 0$  depending only on  $s$  and  $\delta$  and a constant  $\epsilon > 0$  depending only on  $\delta$  and going to zero as  $\delta$  does such that the map  $B(x, r) \rightarrow \mathbb{R}^n$  defined by  $y \mapsto (d(a_1, y), \dots, d(y, a_n))$  is a  $(1 + \epsilon)$ -bilipschitz homeomorphism from  $B(x, r)$  to an open subset of  $\mathbb{R}^n$ .

---

<sup>6</sup>Called “burst points” in [3].



- If there is a  $(n, \delta)$ -strainer for  $X$  at  $x$ , then the strainer dimension of  $X$  at  $x$  is at least  $n$ .

The strainer dimension of  $X$  is the same at every point of  $X$ .

The strainer dimension of  $X$  is its strainer dimension at any of its points.

**Proposition 9.19.** *If  $X$  has strainer dimension  $n$ , then  $X$  is locally compact and every compact neighborhood in  $X$  has rough dimension  $n$ . If  $X$  has strainer dimension  $\infty$ , then  $X$  is not locally compact.*

#### 9.4 Alexandrov Balls

For any  $0 < R \leq \infty$  an Alexandrov ball  $B(x, R)$  of curvature  $\geq k$  is a metric space with the property that:

- It is a metric ball centered at  $x$  of radius  $R$ .
- For every  $0 < R' < R$  the sub-ball  $B(x, R') \subset B(x, R)$  has compact closure in  $B(x, R)$ .
- for any  $p, q \in B(x, R)$  with  $d(x, p) \geq d(x, q)$  if  $d(x, p) + d(p, q)/2 < R$ , then there is a geodesic joining  $p$  and  $q$  in  $B(x, R)$ .
- For any points  $p; a, b, c \in B(x, R)$  with

$$\max(d(p, a), d(p, b), d(p, c)) < R - d(x, p),$$

the  $k$  comparison angles satisfy

$$\tilde{\angle} apb + \tilde{\angle} bpc + \tilde{\angle} cpa \leq 2\pi.$$

The first condition is a type of uniform local completeness for balls. One can think of the second condition in this way. Since we are not assuming any convexity for balls, the second condition is a weaker but uniform condition replacing the existence of geodesics for the ball.

**Example:** 1. Suppose that  $M$  is a complete Riemannian manifold with locally convex boundary and with the sectional curvatures on  $B(x, R) \subset M$  bounded below by  $k$ . Then  $B(x, R)$  is an Alexandrov ball of curvature  $\geq k$ .

2. An Alexandrov ball of radius  $\infty$  and curvature  $\geq k$  is a complete Alexandrov space of curvature  $\geq k$ .

**Lemma 9.20.** *Suppose that  $B(x, R)$  is an Alexandrov ball and that  $\gamma$  and  $\nu$  are geodesics emanating from  $p \in B(x, R)$  of lengths,  $d_1, d_2$  which are at most  $(R - d(x, p))/3$ . Then for  $0 < s \leq d_1$  and  $0 < t \leq d_2$  the comparison angle  $\tilde{\angle} \gamma(s)p\nu(t)$  is a monotone increasing function of either variable when the other is held fixed.*

*Proof.* Let  $T$  be any triangle in  $B(x, R)$  (with geodesic sides and vertices  $v_1, v_2, v_3$ ) with the property if  $a$  is a point on a side of  $T$ , then  $\max(d(a, v_i))_{i=1}^3 < R - d(x, a)$ . Then the defining property holds for  $a; v_1, v_2, v_3$ . Suppose that  $a$  is on the side  $v_1v_2$ . This implies that by the argument given in the case of complete Alexandrov spaces that  $\tilde{\angle}av_1v_3 \geq \tilde{\angle}v_2v_1v_3$ , and hence that the monotonicity statement holds for the comparison angles along the geodesics  $v_1v_2$  and  $v_1v_3$ . Given  $p \in B(x, R)$  and geodesics  $\gamma$  and  $\nu$  emanating from  $p$  of length at most  $(R - d(x, p))/3$  and ending at  $v$  and  $w$ , for any point  $a$  on  $\gamma$  there maximum of the distances from  $a$  to  $p, v, w$  is less than  $R - d(a, x)$  so that the above applies. The result follows.  $\square$

**Definition 9.21.** Fix  $0 < R \leq \infty$  and  $k$ . Suppose that  $R_n \rightarrow R$  and  $k_n \rightarrow k$ . We say that a sequence of Alexander balls  $B(x_n, R_n)$  of curvature  $\geq k_n$  converge in the based Gromov-Hausdorff sense to  $B(x, R)$  if (i)  $R_n \rightarrow R$  as  $n \rightarrow \infty$  and for each  $S < R$  the closed balls  $\overline{B(x_n, S)}$  converge to  $\overline{B(x, S)}$ . Then  $B(x, R)$  is an Alexandrov ball of curvature  $\geq k$  and the  $k_n$ -comparison angles in the  $B(x_n, R_n)$  converge to the  $k$ -comparison angles in  $B(x, R)$ . Implicitly when we discuss Alexandrov balls they are considered based at the central point of the ball and the Gromov-Hausdorff distance and/or convergence is the based version.

**Lemma 9.22.** Fix positive numbers  $a, b$  with  $a + b < 1 - 2\epsilon$ . Suppose that  $B(x, 1)$  and  $B(x', 1)$  are Alexandrov balls within distance  $\epsilon$  of each other in the Gromov-Hausdorff distance, say that we have a distance function  $d$  on  $B(x, 1) \amalg B(x', 1)$  extending the given distance functions on the balls with the property that each ball is in the  $\epsilon$ -neighborhood of the other. Suppose that  $y \in B(x, a)$ . Then for any point  $y' \in B(x', 1)$  with  $d(y, y') < \epsilon$ , then the balls  $B(y, b)$  and  $B(y', b)$  are within  $4\epsilon$  of each other in the Hausdorff distance defined by  $d$ .

#### 9.4.1 Limits that are products

We need a product result for Alexandrov balls.

**Proposition 9.23.** Fix  $r > 0$ . Let  $\lambda_n \rightarrow \infty$  and  $\delta_n \rightarrow 0$  as  $n \rightarrow \infty$ . Suppose that  $X_n = B(p_n, R)$  is a sequence of Alexandrov balls of dimension  $N$  and curvature  $\geq k$ . Suppose that for each  $n$  there are points  $x_n \in X_n$  and compact sets  $\{A_n^+, A_n^-\}$  with

$$d(x_n, A_n^+), d(x_n, A_n^-) \geq 2r,$$

$$A_n^+ \cup A_n^- \cup B(x_n, r) \subset B(p_n, R/3).$$

We also suppose that the comparison angle<sup>7</sup>  $\tilde{\angle}A_n^-x_nA_n^+ > \pi - \delta_n$ . Suppose that the  $(\lambda_n X_n, x_n)$  converge in the Gromov-Hausdorff sense to an  $N$ -dimensional Alexandrov space  $(X, x)$ . Then there is a based Alexandrov space  $(Y, y)$  of dimension  $\leq N-1$  and isometry  $(X, x) \cong (Y, y) \times (\mathbb{R}, 0)$  with the property that for any sequence of points  $z_n \in X_n$  converging to a point  $z \in X$  and geodesics  $\gamma_n^\pm$  from  $z_n$  to  $A_n^\pm$ , the  $\gamma_n^\pm$  converge to the geodesic rays from  $z$  in the positive and negative  $\mathbb{R}$ -directions in the product.

<sup>7</sup>Meaning the angle of the  $k$ -comparison triangle with side lengths  $d(A_n^-, x_n), d(x_n, A_n^+), d(A_n^-, A_n^+)$

*Proof.* Denote by  $g_n$  the metrics on  $X_n$ ; the rescaled metrics are  $\lambda_n^2 g_n$ . Let  $\zeta_n^\pm$  be geodesics from  $x_n$  to  $A_n^\pm$  and let  $y_n^\pm$  be the other endpoint of  $\zeta_n^\pm$ . Since the comparison angle  $\widetilde{\angle} y_n^+ x_n y_n^- \geq \widetilde{\angle} A_n^- x_n A_n^+$  is greater than  $\pi - \delta_n$ , by monotonicity for any points  $u_n^\pm$  on  $\zeta_n^\pm$  the comparison angle  $\angle u_n^- x_n u_n^+$  is greater than  $\pi - \delta_n$ . Hence, rescaling by the  $\lambda_n$  and taking limits we see that for points  $u^\pm$  on the limiting geodesic rays  $\zeta^\pm$  the comparison angle  $\widetilde{\angle} u^- x u^+ = \pi$ , meaning that  $\zeta = \zeta^- \cup \zeta^+$  is a geodesic line. Since the  $X_n$  have curvature  $\geq k$  and the  $\lambda_n \rightarrow \infty$ , the limit  $X$  has curvature  $\geq 0$ . Hence, by Theorem 9.8 it splits as a product  $Y \times \mathbb{R}$  in such a way that  $\zeta$  is the factor in the  $\mathbb{R}$ -direction through the base point. Furthermore, it also follows from this proposition that, letting  $f_n$  be the function

$$d_{\lambda_n^2 g_n}(A_n^-, \cdot) - d_{\lambda_n^2 g_n}(A_n^-, x_n)$$

the  $f_n$  converge to a function  $f: X \rightarrow \mathbb{R}$  whose level sets are the parallel copies of  $Y$  in the product structure. Let  $z_n \in B(x_n, r)$  be a sequence of points converging to  $z \in X$ , and let  $\gamma_n^\pm$  be a geodesic from  $z_n$  to  $A_n^\pm$ . It is easy to see that the  $\gamma_n^+$  converge to rays in the positive  $\mathbb{R}$ -direction. Symmetrically, the  $\gamma_n^-$  converge to rays in the negative  $\mathbb{R}$ -direction.  $\square$

**Addendum 9.24.** Analogous arguments work to show the following: Given a sequence of constants and balls as in the previous proposition and sequences of compact sets  $A_n^+, A_n^-, (A'_n)^+, (A'_n)^-$  with each pair  $\{A_n^+, A_n^-\}$  and  $\{(A'_n)^+, (A'_n)^-\}$  satisfying the hypothesis of the previous proposition and with the angles  $\angle A_n^\pm x_n (A'_n)^\pm$  converging to  $\pi/2$ , the limit can be written isometrically as a product of  $(Y, y) \times (\mathbb{R}^2, 0)$  where the limiting geodesics to the four compact sets form the  $x$ - and  $y$ -axes in the  $\mathbb{R}^2$ -direction through the central point  $(y, 0)$ .

## 9.5 The Tangent Cone

Let  $x \in X$  be a point in a complete Alexandrov space or in an Alexandrov ball. We define the metric space of germs of geodesics at  $x$  as follows. The underlying set is the set of equivalence classes of geodesics emanating from  $x$ , with  $\gamma$  and  $\nu$  being equivalent if and only if their intersection is a non-trivial geodesic. We define a metric by  $d([\gamma], [\nu])$  is the angle at  $x$  between  $\gamma$  and  $\nu$ . It is easy to see that this distance depends only on the equivalence classes and that it is a metric on the set of equivalence classes of geodesics emanating from  $x$ . The tangent sphere  $S_x(X)$  is the metric completion of this metric space, cf [3].

**Proposition 9.25.** (See [3].) *Suppose that  $X$  is a complete Alexandrov space or an Alexandrov ball and  $x \in X$ . Then  $S_x(X)$  is a compact metric space of diameter  $\leq \pi$ .*

Fix an Alexandrov ball  $X = B(y, R)$  and curvature  $\geq k$  and of dimension  $n$ , and fix  $x \in X$ . Consider a sequence of constants  $\lambda_\ell \rightarrow \infty$  as  $\ell \rightarrow \infty$ . Then the based Alexandrov spaces  $(\lambda_\ell X, x)$  are of dimension  $n$  and curvature  $\geq k/\lambda_\ell^2$ . Hence, passing to a subsequence there is a limit  $T_x X$  which is an Alexandrov space of dimension  $\leq n$  and curvature  $\geq 0$ .

The monotonicity of angles along geodesics easily implies the following:

**Claim 9.26.**  $T_x X$  is isometric to the cone over the tangent sphere  $S_x X$ .

**Corollary 9.27.** Suppose that  $X$  is an Alexandrov ball of curvature  $\geq k$  and of dimension  $n$ . Then,  $S_x X$  is a compact Alexandrov space of dimension  $n - 1$ , curvature  $\geq 1$  and diameter  $\leq \pi$ , and  $(\lambda X, x)$  converges in the Gromov-Hausdorff sense to  $T_x X$ , the cone on  $S_x X$ , as  $\lambda \rightarrow \infty$ .

**Definition 9.28.**  $T_x X$  is the tangent cone of  $X$  at  $x$ .

## 9.6 Consequences of the existence of Tangent Cones

Now using the tangent cone we can establish;

**Theorem 9.29.** Suppose that  $X$  is a complete Alexandrov space or an Alexandrov ball. Suppose also that  $X$  is of dimension  $n$ . Then for every  $\delta > 0$ , the subset of points  $x \in X$  at which  $X$  has an  $(n, \delta)$ -strainer is an open dense set.

*Proof.* If  $n = 1$ , then  $X$  is isometric to either a line, a half-line, a compact interval, or a circle. All points of  $X$  except its endpoints have  $(1, \delta)$ -strainers for every  $\delta > 0$ .

Suppose by induction that we know the result for  $n' < n$  and fix  $x \in X$  and  $\delta > 0$ . Then the tangent sphere  $S_x X$  is an Alexandrov space of dimension  $n - 1$  and hence has an open dense subset  $U$  of points at which  $S_x X$  has an  $(n - 1, \delta)$ -strainer. It follows that every point of  $T_x X$  contained in the cone on  $U$  except the cone point has a  $(n, \delta)$ -strainer. By the above convergence result, it follows that there are points of  $X$  arbitrarily close to  $x$  at which  $X$  has an  $(n, \delta)$ -strainer. This proves the subset of points at which  $X$  has an  $(n, \delta)$ -strainer is dense.

Clearly from the definition, the set of points with an  $(n, \delta)$  strainer is open in  $X$ .  $\square$

**Lemma 9.30.** For each natural number  $n$  there is a constant  $c(n)$  so that for any  $n$ -dimensional compact Alexandrov space  $S$  with curvature  $\geq 1$  the  $n$ -dimensional rough volume  $Vr_n(S)$  is at most  $c(n)$ . For any  $\epsilon > 0$  sufficiently small, every  $\epsilon$ -net in  $S$  has cardinality at most  $c(n)\epsilon^{-n}$ .

*Proof.* It is easy to see that any such Alexandrov space has diameter  $\leq \pi$ . (Actually, we shall make use of this result only for tangent spheres where we have this bound immediately.) From this and an induction on dimension it is straightforward to establish the result.  $\square$

**Corollary 9.31.** There is a constant  $c(n, k, R)$  such that the following holds. Let  $X$  be a complete  $n$ -dimensional Alexandrov space of curvature  $\geq k$ . Then for any  $x \in X$  and any  $R < \infty$  the cardinality of an  $\epsilon$ -net in  $B(x, R)$  is at most  $c(n, k, R)\epsilon^{-n}$ .

This leads immediately to a sequential compactness result for Alexandrov spaces.

**Corollary 9.32.** Let  $(X_i, x_i)$  be a sequence of complete Alexandrov spaces of dimension  $\leq n$  and curvature  $\geq k$ . Then, after passing to a subsequence there is a Gromov-Hausdorff limit. Any such limit is a complete Alexandrov space of dimension at most  $n$  and curvature  $\geq k$ .

*Proof.* This is direct from the previous corollary and Corollary 8.5.  $\square$

There is also a version of this result for Alexandrov balls.

**Corollary 9.33.** *Let  $B(x_i, R_i)$  be a sequence of Alexandrov balls of curvature  $\geq k$  with  $R_i \rightarrow R$  with  $0 < R \leq \infty$  as  $i \rightarrow \infty$ . Then, after passing to a subsequence, the balls  $B(x_i, R_i)$  converge in the Gromov-Hausdorff sense to a limit  $B(x_\infty, R)$  that is an Alexandrov ball of curvature  $\geq k$ .*

*Proof.* The above arguments show that for any  $R' < R$  there is a uniform bound to the cardinality of any  $\epsilon$ -net in  $\overline{B(x_i, R')}$ , so that passing to a subsequence we can arrange that these compact balls converge. Taking a sequence of  $R'_n \rightarrow R$  and passing to a diagonal sequence we construct a Gromov-Hausdorff limit of the  $B(x_i, R_i)$ . It is immediate to see that the limit is an Alexandrov ball of curvature  $\geq k$ .  $\square$

**Remark 9.34.** Gromov-Hausdorff limits of manifolds, or Alexandrov spaces, of a given dimension can have strictly smaller dimension. From example, a sequence of  $n$ -spheres of radii  $r_i \rightarrow 0$  is a sequence of  $n$ -manifolds with curvature  $\geq 0$ . This sequence converges in the Gromov-Hausdorff sense to a point, which is an Alexandrov space of rough dimension 0.

**Definition 9.35.** The boundary of an Alexandrov ball is defined inductively on dimension. Let  $X$  be a one-dimensional Alexandrov ball. Then it is either isometric to either an interval or a circle. Its boundary as an Alexandrov ball is its topological boundary. More generally, we define the boundary of a higher dimensional Alexandrov ball by induction. For  $X$  an  $n$ -dimensional Alexandrov ball, we define  $\partial X$  to be the subset of  $X$  consisting of points  $p$  for which  $\Sigma_p$  is an  $(n-1)$ -dimensional compact Alexandrov space (and hence an Alexandrov ball) with non-empty boundary. Then  $\partial X$  is a closed subset. Its complement is denoted  $\text{int } X$ .

### 9.6.1 Bounding the number of small loops

We give a general result which allows us to bound the number of homotopy classes represented by small loops.

**Proposition 9.36.** *There is  $\ell_0 > 0$  such that the following holds. For any choice of positive constants  $\ell, r, \epsilon$ , each at most  $\ell_0$ , there is a constant  $N_n(\ell, r, \epsilon) < \infty$  depending on these constants and the dimension  $n$  such that the following holds. Suppose that  $B = B(x, 1)$  is an Alexandrov ball of dimension  $n$  and curvature  $\geq -1$ , that  $y \in B$  with  $d(x, y) = \ell$ . Let  $\Gamma \subset \pi_1(B, x)$  denote the image of  $\pi_1(B(y, r), y) \rightarrow \pi_1(B, x)$  defined by sending a loop  $\alpha$  based at  $y$  to  $\gamma^{-1}\alpha\gamma$  where  $\gamma$  is a (fixed) geodesic from  $x$  to  $y$ . Then, for any group  $H$  and surjective homomorphism  $f: \pi_1(B, x) \rightarrow H$  that corresponds to a covering space of  $B$ , the number of cosets in  $C = H/f(\Gamma)$  represented by loops based at  $x$  of length at most  $\epsilon$  is at most  $N_n(\ell, r, \epsilon)$ .*

*Proof.* Let  $p: \tilde{B} \rightarrow B$  be the covering corresponding to  $f: \pi_1(B, x) \rightarrow H$ . Fix a lift  $\tilde{x}$  of  $x$ , and let  $\tilde{y}$  be the lift of  $y$  that is connected to  $\tilde{x}$  by a lift of  $\gamma$ . We define

a metric on  $\tilde{B}$  as follows: given  $\tilde{a}, \tilde{b} \in \tilde{B}$  we set  $\tilde{d}(\tilde{a}, \tilde{b}) = \inf \{\ell(p(\omega))\}$  as  $\omega$  ranges over all paths in  $\tilde{B}$  connecting  $\tilde{a}$  and  $\tilde{b}$  with rectifiable image under  $p$ . (Here,  $\ell(p(\omega))$  denotes the length of the path  $p(\omega)$ .) Every point of  $\tilde{B}$  has a neighborhood that projects homeomorphically under  $p$  and with the property that the metric  $\tilde{d}$  agrees on this neighborhood with the pull back under  $p$  of the metric on  $B$ .

The pre-image  $p^{-1}(B(y, r))$  is a disjoint union  $\coprod_{c \in C} U_c$  where  $p$  induces a covering map  $U_c \rightarrow B(y, r)$ . The component  $U_e$  is the one that contains  $\tilde{y}$ . Under the action of  $\pi_1(B, x)$  on  $\tilde{B}$ , an element  $a \in \pi_1(B, x)$  sends  $U_e$  to  $U_{[f(a)]}$  where  $[f(a)]$  denotes the coset  $f(a) \cdot f(\Gamma) \in C$ . Notice that for  $c \neq c'$  in  $C$  we have  $U_c \cap U_{c'} = \emptyset$ . Also, notice that since  $U_c$  projects onto  $B(x, r)$ ,  $U_c$  contains the ball of radius  $r$  about any lift of  $y$  contained in  $U_c$ . This implies that if  $a, a' \in \pi_1(B, x)$  and  $[f(a)] \neq [f(a')]$ , then  $d(a\tilde{y}, a'\tilde{y}) \geq 2r$ . Let  $a_1, \dots, a_n$  be elements of  $\pi_1(B, x)$  represented by loops of length at most  $\epsilon$  based at  $x$  and suppose that the associated cosets  $f(a_1)f(\Gamma), \dots, f(a_n)f(\Gamma)$  in  $C$  are distinct. We label these cosets  $c_1, \dots, c_n$ . Then the  $a_i\tilde{x}$  all within distance  $\epsilon$  of  $\tilde{x}$  and hence  $\ell - \epsilon < \tilde{d}(\tilde{x}, a_i\tilde{y}) < \ell + \epsilon$ . Set  $A = \max([1 + (4\epsilon/r)], [t])$  where  $[t]$  denotes the greatest integer less than or equal to  $t$ . Then divide the interval  $[\ell - \epsilon, \ell + \epsilon]$  into  $A$  subintervals each of length at most  $r/2$ . Then for one of these intervals there are at least  $n/A$  of the  $\tilde{y}_i$  whose distance to  $\tilde{x}$  lies in this interval. Hence, we have  $n' = [n/A]$  points, which after relabelling we can take to be  $\{\tilde{y}_1, \dots, \tilde{y}_{n'}\}$  in  $B(\tilde{x}, \ell + r)$  with the property that  $d(\tilde{y}_i, \tilde{y}_j) \geq 2r$  for all  $i \neq j$  and  $|d(\tilde{x}, \tilde{y}_i) - d(\tilde{x}, \tilde{y}_j)| < r/2$ . This implies that for every  $1 \leq i < j \leq n'$  the comparison angle  $\tilde{\angle}_{\tilde{y}_i\tilde{x}\tilde{y}_j}$  is bounded below by a positive constant depending only on  $\ell, r$  and  $\ell_0$  (provided that  $\ell_0$  is sufficiently small).

Notice that since  $\tilde{B}$  is a local Alexandrov space, every point of  $\tilde{B}$  has a tangent sphere which is compact and of curvature  $\geq 1$ .

**Claim 9.37.** *For each  $i = 1, \dots, n'$  let  $\gamma_i$  be a geodesic from  $\tilde{y}_i$  to  $\tilde{x}$ . Then for each  $i \neq j$  the angle between  $\gamma_i$  and  $\gamma_j$  at  $\tilde{x}$  is at least as large as the comparison angle  $\tilde{\angle}_{\tilde{y}_i\tilde{x}\tilde{y}_j}$ .*

Given this claim the result is immediate from the uniform lower bound on the comparison angles and Lemma 9.30.

*Proof.* (of the claim) This is the standard monotonicity result on angles and follows if we can show that the Alexandrov property holds for all quadruples  $\{a; b, c, d\}$  in  $B(\tilde{x}, 2\ell_0) \subset \tilde{B}$ . Here is what we know about  $\tilde{B}$ :

1. It is local Alexandrov space of curvature  $\geq -1$ .
2. The ball  $B(\tilde{x}, 2/3)$  has compact closure in  $\tilde{B}$ .
3. Every pair of points in  $B(\tilde{x}, 1/3)$  is joined by a geodesic in  $B(\tilde{x}, 2/3)$ .

The reasons for these are: (i)  $\tilde{B}$  is locally isometric to  $B$ ; (ii)  $\overline{B(\tilde{x}, 2/3)}$  is a closed and bounded subset of  $p^{-1}(\overline{B(x, 2/3)})$  and the latter is a complete metric space being a covering of a complete metric space with the covering projection being distance non-increasing; (iii) Follows from the second and the usual curve shortening arguments.

Then according to Remark 3.5 of [3], these three properties imply that there is an  $\ell_0 > 0$  such that the Alexandrov property holds for all 4-tuples in  $B(\tilde{x}, 2\ell_0)$ . This completes the proof of the claim.  $\square$

This completes the proof of the proposition.  $\square$

## 9.7 Directional Derivatives

Let  $X$  be either a complete Alexandrov space or an Alexandrov ball, and let  $f: X \rightarrow \mathbb{R}$  be a Lipschitz function. We say that  $f$  has a *directional derivative at  $x$* , if there is a continuous function  $f': S_x X \rightarrow \mathbb{R}$  such that for any geodesic  $\gamma$  emanating from  $x$  and parametrized by arc length we have

$$\lim_{t \rightarrow 0} \frac{f(\gamma(t)) - f(x)}{t} = f'([\gamma]).$$

The main example of this is the following:

**Lemma 9.38.** *Let  $X$  be a complete Alexandrov space. Let  $A$  be a compact subset of  $X$  and let  $y \in X \setminus A$ . Let  $d: X \rightarrow \mathbb{R}$  be the distance function from  $A$  and let  $A' \subset S_y X$  be the set of tangent directions to geodesics from  $y$  to  $A$ . Then  $d$  is a Lipschitz function and  $d$  has a directional derivative  $d'$  at  $y$  given by*

$$d'(\alpha) = -\cos(d(\alpha, A')).$$

**Remark 9.39.** The same result holds when  $X$  is an Alexandrov ball  $B(x, R)$  provided that  $A$  and  $y$  are contained in  $B(x, R - d)$  where  $d$  is the distance from  $A$  to  $y$ .

**Definition 9.40.** Let  $X$  be a complete Alexandrov space or an Alexandrov ball, let  $U \subset X$  be an open set and let  $f: U \rightarrow \mathbb{R}$  be a Lipschitz function with a directional derivative at every point of  $U$ . We say that  $f$  is *regular* at  $x \in U$  if there is a direction  $\tau \in S_x X$  such that  $f'_x(\tau) > 0$ .

**Lemma 9.41.** *Let  $X$  be a complete Alexandrov space. Suppose that  $A$  is a compact set and  $U$  is an open subset of  $X$ , disjoint from  $A$ . Then the subset  $V \subset U$  of points at which  $d = d(A, \cdot)$  is regular is an open subset.*

*Proof.* Suppose that  $v \in V$ . Then there is a geodesic  $\gamma$  emanating from  $v$  such that  $(d(A, \gamma(s)) - d(A, v))/s$  has limit  $> 0$  at  $s = 0$ . This means that for any minimal geodesic  $\alpha$  connecting  $v$  to  $A$ , the angle at  $v$  between  $\alpha$  and  $\gamma$  is greater than  $\pi/2$ . Denote by  $w$  the other endpoint of  $\gamma$ . By choosing  $\gamma$  sufficiently short, we can assume that  $d(A, w) > d(A, v)$  and that there is a unique geodesic from  $v$  to  $w$ . Let  $v_n$  be a sequence of points in  $U$  converging to  $v$ , and let  $\gamma_n$  be a geodesic from  $v_n$  to  $w$ . Then  $\gamma_n$  converge to  $\gamma$  as  $n \rightarrow \infty$ . Suppose that for each  $n$  there is a geodesic  $\mu_n$  from  $A$  to  $v_n$  such that the angle at  $v_n$  between  $\mu_n$  and  $\gamma_n$  is at most  $\pi/2$ . Passing to a subsequence we can suppose that the  $\mu_n$  converge to a geodesic  $\mu$  from  $A$  to  $v$ , and we know the  $\gamma_n$  converge to  $\gamma$ . Thus, by Lemma 9.4 the angle between  $\mu$  and  $\gamma$  is at most  $\pi/2$ , which is a contradiction.  $\square$

Similarly, one shows:

**Corollary 9.42.** *Suppose that we have a sequence of Alexandrov balls  $B_n = B(x_n, R_n)$  of curvature  $\geq k$  converging in the Gromov-Hausdorff topology to a limit  $B = B(x, R)$ . Suppose that there are compact subsets  $A_n \subset B_n$  converging to a compact subset  $A \subset B$  and open subsets  $V_n$  converging to  $V$ . Suppose that there is a geodesic from  $A_n$  to each point of  $V_n$ , and suppose that  $q \in V$  and  $q_n \in V_n$  is a sequence converging to  $q$ . Then if  $d(A, \cdot)$  is regular at  $q$ , then for all  $n$  sufficiently large,  $d(A_n, \cdot)$  is regular at  $q_n$ .*

Likewise, we have:

**Lemma 9.43.** *Suppose that  $f: B \rightarrow \mathbb{R}$  is a Lipschitz function with directional derivatives and that  $q_n \in f^{-1}(f(q))$  is a sequence converging to  $q$ . Let  $\gamma_n$  be a geodesic from  $q$  to  $q_n$ . Suppose that the unit tangent vectors to the  $\gamma_n$  at  $q$  converge to a tangent direction  $\tau$ . Then  $f'_q(\tau) = 0$ .*

*Proof.* This is elementary from the comparison results, see §11.3 of [3].  $\square$

### 9.7.1 Regular functions on smooth manifolds

We shall need information about level sets of regular functions on smooth manifolds.

**Lemma 9.44.** *Suppose that  $X$  is a locally complete Riemannian manifold and that  $f$  is the distance function from a compact set  $A$  and that  $f$  is regular (in the Alexandrov sense) at  $q_0 \in X \setminus A$ . Then there is a neighborhood  $U$  of  $q_0$  and a smooth unit vector field  $\tau$  on  $U$  with the property that  $f'_q(\tau) > 0$  for all  $q \in U$ . Furthermore, there is an open interval  $J$ , an open subset  $U'$  of  $\mathbb{R}^{n-1}$ , and a bi-Lipschitz homeomorphism  $U \cong U' \times J$  with the property that the level sets of  $f|_U$  are identified with the subsets  $U' \times \{j\}$  for  $j \in J$ . In particular, the level sets of  $f$  are topologically locally flat, codimension-1 submanifolds near  $q$ .*

*Proof.* Consider the subset of the unit tangent bundle of  $X$  consisting of directions  $\chi_q \in T_q X$  with the property that  $f'_q(\chi_q) > 0$  as  $q$  varies over an open neighborhood  $U$  of  $q_0$ . Arguments similar to the above show that this is an open subset  $\mathcal{O}$  of  $TX$ . If we take  $U$  small enough, the fiber of  $\mathcal{O}$  over every  $q \in U$  is non-empty. Hence, after shrinking  $U$ , there is a smooth unit vector field  $\tau$  defined in a neighborhood  $U$  of  $q$  and  $\alpha > 0$  such that  $f'_q(\tau(q)) \geq \alpha$  for all  $q \in U$ . Now we integrate  $\tau$  to define a smooth local coordinate system  $(x^1, \dots, x^n)$  near  $q_0$  such that  $\tau = \partial/\partial x^1$ . We replace  $U$  by a smaller open set which is the product of an open ball in  $(x^2, \dots, x^n)$ -space with an interval in the  $x^1$ -direction. Since  $f'(\partial/\partial x^1) > 0$  everywhere, we see that the level sets of  $f$  meet each interval in the  $x^1$ -direction in at most one point. That is to say, near  $q_0$  these level sets are given by the graphs of functions  $x^1 = \varphi(x^2, \dots, x^n)$ . Elementary arguments show that the map  $(x^1, \dots, x^n) \mapsto (f(x^1, \dots, x^n), x^2, \dots, x^n)$  is the required bi-Lipschitz homeomorphism.  $\square$

We also need a fairly restricted version of an analogous result for maps to the plane. The following is an elementary lemma.



**Lemma 9.45.** *Given  $\epsilon' > 0$ , the following holds for all  $\epsilon > 0$  sufficiently small. Let  $B(0, \epsilon^{-1})$  be the ball of radius  $\epsilon^{-1}$  in the Euclidean plane centered at the origin. We denote by  $(x, y)$  the Euclidean coordinates on this ball and by  $\theta$  the usual coordinate along the circle. Let  $g$  be a Riemannian metric on  $U = B(0, \epsilon^{-1}) \times S^1$  that is within  $\epsilon$  in the  $C^N$ -topology (where  $N = \lfloor \epsilon^{-1} \rfloor$ ) of the product of the usual Euclidean metric on  $B(0, \epsilon^{-1})$  and the Riemannian metric of length 1 on the circle. Suppose that  $F = (f_1, f_2): U \rightarrow \mathbb{R}^2$  is a map with the property that  $f_1$  and  $f_2$  are 1-Lipschitz with respect to  $g$  with directional derivatives at all points of  $U$ . Suppose further that the directional derivatives of  $f_i$  with respect to  $g$  satisfy:*

$$|f'_1(\partial_x) - 1| < \epsilon$$

$$|f'_2(\partial_y) - 1| < \epsilon$$

$$\max(|f'_1(\pm\partial_y)|, |f'_2(\pm\partial_x)|, |f'_1(\pm\partial_\theta)|, |f'_2(\pm\partial_\theta)|) < \epsilon.$$

*Then any fiber  $F^{-1}(p)$  that meets  $B(0, \epsilon^{-1}/2)$  is a circle that is  $\epsilon'$ -orthogonal to the family of horizontal spaces  $B(0, \epsilon^{-1}) \times \{\theta\}$  in the sense that, fixing  $a \in F^{-1}(p)$ , as  $b \in F^{-1}(p)$  approaches  $a$  the angle (measured with respect to product metric) of the geodesic (in the product metric) from  $a$  to  $b$  with the horizontal space through  $a$  is within  $\epsilon'$  of  $\pi/2$ . Furthermore, any fiber  $F^{-1}(p)$  that meets  $B(0, \epsilon^{-1}/2)$  intersects each horizontal space  $\{\theta\} \times B(0, \epsilon^{-1})$  in a single point.*

### 9.7.2 A smooth limit result

As we have already indicated, the entire argument revolves around considering sequences  $\{x_n \in M_n\}_{n=1}^\infty$ , rescaling the metrics  $g_n$ , and, after passing to a subsequence, extracting a limit (usually a Gromov-Hausdorff limit) of the metric unit balls in the rescaled metrics. In general, a limit like this can be of dimension 1, 2, or 3 (although when we use  $\rho_n^{-2}(x_n)$  to rescale the limit, the volume collapsing hypothesis implies that the limit has dimension 1 or 2) and depending on which it is we get a different structure for balls. The easiest case to treat is when the limit is 3-dimensional. As the next theorem shows, because of the assumption on bounds on the curvature and its derivatives in the statement of Theorem 5.5, such limits are automatically smooth limits, rather than the more general Gromov-Hausdorff limits that occur in the other two cases.

**Proposition 9.46.** *Let  $(M_n, g_n)$  and  $w_n$  be as in the statement of Theorem 5.5. Suppose that we have a sequence of points  $x_n \in M_n$  such that  $B_n = B_{g_n}(x_n, \rho_n(x_n))$  is disjoint from  $\partial M_n$  and a sequence of constants  $\lambda_n^2$  with a Gromov-Hausdorff limit of a subsequence of  $(B_n, \lambda_n^2 g_n, x_n)$ , which is a 3-dimensional Alexandrov space. Then, passing to a further subsequence, there is a smooth limit of the  $(B_n, \lambda_n^2 g_n, x_n)$ , which is a complete manifold of non-negative curvature.*

*Proof. First step:*

**Claim 9.47.** *If  $(B_n, \lambda_n^2 g_n, x_n)$  converges to a 3-dimensional Alexandrov space, then there is a sequence of points  $y_n \in B_n$  converging to a point  $y$  in the limit and*

constants  $r > 0$  and  $\kappa > 0$  such that for all  $n$  sufficiently large  $\text{Vol } B_{\lambda_n^2 g_n}(y_n, r) \geq \kappa r^3$ .

*Proof.* Fix  $\delta > 0$  sufficiently small. Let  $X$  be the limiting 3-dimensional Alexandrov space. By Corollary 6.7 of [3] the subset  $R_\delta(X)$  consisting of points with a  $(3, \delta)$ -strainer is dense. Choose  $y \in R_\delta(X)$  and let  $y_n \in M_n$  be a sequence converging to  $y$ . Then there is a  $(3, \delta)$ -strainer  $\{a_1, b_1, a_2, b_2, a_3, b_3\}$  at  $y$ . Let  $d$  be the size of this strainer. Hence for all  $n$  sufficiently large, there is a  $(3, \delta)$ -strainer of size  $d/2$  at  $y_n$  in  $\lambda_n^2 B_n$ . According to Lemma 9.18 this means that for some  $r \ll d/2$ , but depending only on  $d$ , there is an almost bilipschitz homeomorphism from  $B_{\lambda_n^2 g_n}(y_n, r)$  to the ball of radius  $r$  in Euclidean space, where the error estimate goes to zero with  $\delta$ . Hence, there is  $\epsilon_0 > 0$  such that for any  $0 < \epsilon \leq \epsilon_0$  and for all  $n$  sufficiently large, the cardinality of a maximal  $\epsilon$ -net in  $B_{\lambda_n^2 g_n}(y_n, r)$  is at least  $\alpha \epsilon^{-3} r^3$  for a universal constant  $\alpha > 0$ . If we choose  $\epsilon > 0$  sufficiently small depending on  $n$  then the volume in  $\lambda_n^2 g_n$  of any ball of radius  $\epsilon/2$  centered at a point of  $B_{\lambda_n^2 g_n}(y_n, r)$  is at least  $(1/2)\omega_0(\epsilon/2)^3$  where  $\omega_0$  is the volume of the unit ball in Euclidean 3-space. Hence,  $\text{Vol } B_{\lambda_n^2 g_n}(y_n, (r+\epsilon)) \geq \alpha \omega_0 r^3/16$ . Taking the limit as  $\epsilon \rightarrow 0$  gives the uniform lower bound to the volume of the ball of radius  $B_{\lambda_n^2 g_n}(y_n, r)$ .  $\square$

**Second Step:** Suppose that  $y_n \in B_n$  is as in the previous claim. Then, the  $(B_n, \lambda_n^2 g_n)$  are uniformly volume non-collapsed at  $y_n$ . That is to say for some  $r > 0$  and  $w' > 0$ , for all  $n$  the volume of  $B_{\lambda_n^2 g_n}(y_n, r)$  is at least  $w' r^3$ . Since the ball  $B(y_n, \rho(y_n))$  has volume is at most  $w_n \rho(y_n)^3$  where  $w_n \rightarrow 0$  as  $n \rightarrow \infty$ , it follows from Bishop-Gromov volume comparison that  $\rho(y_n) \lambda_n \mapsto \infty$  as  $n$  tends to infinity. Hence, for any  $A < \infty$ , for all  $n$  sufficiently large, we have  $4A < \rho(y_n) \lambda_n$ . Thus, by our assumption, for all  $n$  sufficiently large, the sectional curvatures of  $\lambda_n^2 g_n$  on  $B_{\lambda_n^2 g_n}(y_n, 4A)$  are  $\geq -\lambda_n^{-2} \rho(y_n)^{-2} > -(4A)^{-2}$ . Again invoking the Bishop-Gromov inequality, we see that there is a constant  $w''(w', A) > 0$  such that for any  $s \leq A$  and any  $z \in B_{\lambda_n^2 g_n}(y_n, A)$  we have  $\text{Vol}(B_{\lambda_n^2 g_n}(z, s)) \geq w'' s^3$ . Taking  $r = \min(A/\lambda_n, \bar{r}(w''))$ , where  $\bar{r}(w'')$  is the constant from Condition 3 of Theorem 5.5, we see that for any  $z_n \in B_{\lambda_n^2 g_n}(y_n, A)$  the volume of  $B_{\lambda_n^2 g_n}(z_n, r) \geq w'' r^3$  and the sectional curvatures on this ball are bounded below by  $-r^{-2}$ . Since  $r \leq \bar{r}(w'') \lambda_n$ , it follows from Proposition 2.9 we have uniform bounds on the curvature and all of its derivatives at every point of  $B_{\lambda_n^2 g_n}(y_n, A)$  depending only on  $A$  and  $w'$ . Hence, we can pass to a subsequence, so that the  $B_{\lambda_n^2 g_n}(y_n, A)$  have a smooth limit. Taking a sequence of  $A$  tending to infinity and a diagonal subsequence allows us to pass to a subsequence so that the  $(M_n, \lambda_n^2 g_n, y_n)$  have a smooth, complete limit. Since  $\rho(y_n) \lambda_n$  tends to infinity, the curvature of the limiting manifold is  $\geq 0$ .  $\square$

This result about the 3-dimensional limits will be important as we study the 1- and 2-dimensional limits.

## 9.8 Blow-up results

We need two special results about rescaling Alexandrov spaces so as to construct higher dimensional limits. We need these results in order to handle sequences of

points  $x_n \in M_n$  converging to a singular point of a 1- or 2-dimensional limit. The following two results are reformulations in our context of Lemma 3.6 of [32].

**Proposition 9.48.** *Suppose that  $B_n = B(x_n, 1)$  is a sequence of Alexandrov balls of dimension  $d$ , of radius 1, and with curvature  $\geq k$ . Suppose that the  $B_n$  are non-compact and converge to an interval  $J$  with the  $x_n$  converging to the endpoint  $x$  of  $J$ . Fix  $\pi/2 < \alpha < \pi$ . Then, after passing to a subsequence, there are points  $\hat{x}_n \in B_n$  with  $d(x_n, \hat{x}_n) \rightarrow 0$  as  $n \rightarrow \infty$  such that one of the following holds:*

1. *At every point of  $B(\hat{x}_n, 1/2) \setminus \{\hat{x}_n\}$  there is a direction in which the directional derivative of the distance function  $f_n = d(\hat{x}_n, \cdot)$  is greater than  $\alpha$ . In this case for every  $0 < r' < 1/2$  the metric ball  $B(\hat{x}_n, r')$  is homeomorphic to the tangent cone at  $\hat{x}_n$  and if  $B_n$  are smooth manifolds then the  $B(\hat{x}_n, r')$  is diffeomorphic to a smooth ball in  $\mathbb{R}^d$ .*
2. *There is a sequence of positive constants  $\zeta_n \rightarrow 0$  as  $n \rightarrow \infty$  such that:*
  - (a) *Every point in  $B(\hat{x}_n, 1/2)$  at which the maximum value of the directional derivative of  $f_n$  is at most  $\alpha$  is within distance  $\zeta_n$  of  $\hat{x}_n$ , and*
  - (b) *there is a point  $q_n$  at distance  $\zeta_n$  from  $\hat{x}_n$  at which the maximum value of the directional derivative is at most  $\alpha$ .*

*In this case, passing to a subsequence there is a limit  $(X, z)$  of the  $3\zeta_n^{-1}B(\hat{x}_n, 1/2)$ . This limit is a complete Alexandrov space of curvature  $\geq 0$  and of dimension strictly greater than 1. The distance from  $z$  has no critical points at distance greater than  $1/3$  from  $z$ . If the limit is 2-dimensional then the area of any unit ball  $B(y, 1)$  for any  $y \in B(x, 1/2)$  has area at least a positive constant  $a(\alpha)$  depending only on  $\alpha$ . Furthermore, the restriction of  $f$  to subset  $\{w \in X \mid f(w) > 1/3\}$  is the topological projection mapping of a product with fibers being either closed intervals or circles.*

*Proof.* We fix  $\alpha$  with  $\pi/2 < \alpha < \pi$  and consider a sequence  $B_n$  as in the hypothesis. We take a point  $y \in J$  at distance  $3/4$  from the endpoint  $x$  of  $J$ , and we take a sequence  $y_n \in B_n$  converging to  $y$ . For each  $n$  sufficiently large there is a maximum  $\hat{x}_n$  for  $d(y_n, \cdot)$ . Then  $\hat{x}_n \rightarrow x$  as  $n \rightarrow \infty$  so that  $d(x_n, \hat{x}_n) \rightarrow 0$  as  $n \rightarrow \infty$ . One possibility is that there is a subsequence of  $n$  for which  $f_n = d(\hat{x}_n, \cdot)$  has no points outside of  $\hat{x}_n$  in  $B_n$  at which the maximum of  $f'_n$  is  $\leq \alpha$ . In this case, the first conclusion stated in the proposition holds. Otherwise, we can pass to a subsequence such that for all  $n$  there are points distinct from  $\hat{x}_n$  at which the directional derivative of  $f_n$  is bounded above by  $\alpha$ . Now consider any sequence (in  $n$ ) of points  $q_n \in B_n \setminus \{\hat{x}_n\}$  with the maximum value of the directional derivative for  $f_n$  at  $q_n$  being at most  $\alpha$ . Passing to a subsequence we see that the sequence converges to the endpoint  $x$  of  $J$ . In particular, there is a sequence  $\zeta_n \rightarrow 0$  as  $n \rightarrow \infty$  such that the maximum distance of the set of all  $q_n$  in  $B(x_n, 1/2)$  with the property that the maximal value of the directional derivative of  $f_n$  at  $q_n$  is at most  $\alpha$  is  $\zeta_n$ . Fix a point  $q_n$  with this property at distance  $\zeta_n$  from  $\hat{x}_n$ , and rescale the balls by  $3\zeta_n^{-1}$ . Passing to a subsequence there is a limiting based Alexandrov space  $(X, z)$  of curvature  $\geq 0$ .

This space is complete and non-compact. Clearly, it has the property that all critical points of  $f = d(z, \cdot)$  are at distance  $\leq 1/3$  from  $z$ .

Next, we show that  $X$  has dimension at least 2. If not, then  $X$  is a non-compact interval and there is a point at distance  $1/3$  from  $z$  at which the maximum of the directional derivative is at most  $\alpha$  and no such points at distances more than  $1/3$  from  $z$ . This means that  $X$  is a ray with  $z$  being at distance  $1/3$  from the endpoint, and the  $q_n$  converge to the endpoint of  $X$ . We claim that this contradicts the fact that  $\hat{x}_n$  maximizes the distance from  $y_n$ . Fix  $1/3 < D$  and consider the interval of length  $D$  (in the rescaled metric) on any geodesic from  $y_n$  to  $q_n$  with one endpoint being  $q_n$ . These compact intervals converge to an interval of length  $D$  in  $X$  with one endpoint being the endpoint of  $X$ . It follows that the points at distance  $1/3$  from  $q_n$  on these geodesics converge to  $z$ , and hence for  $n$  large are arbitrarily close to  $x_n$ . It then follows that the distance from  $y_n$  to  $q_n$  is greater than the distance from  $y_n$  to  $\hat{x}_n$ , which is a contradiction.

This shows that  $X$  has dimension at least 2. Clearly, by construction,  $f$  has only regular values on  $f^{-1}(1/3, \infty)$ . It then follows from the theory of Alexandrov spaces that  $f$  is the projection mapping of a topological product structure. Since the  $B_n$  converge to an interval  $J$ , for all  $n$  sufficiently large the distance function from  $\hat{x}_n$  is regular on the complement of the ball of radius  $1/25$  centered at  $x_n$ ; furthermore, the fibers of this map are connected. This means that all the fibers of  $f_n$  at distance more than  $\zeta_n$  from  $\hat{x}_n$  are connected. Thus, if the dimension of  $X$  is 2 then all the fibers of  $f$  at distance more than  $1/3$  are either closed intervals or circles.

Still supposing that the dimension of  $X$  is 2, we shall show that there is a positive lower bound to the area of  $B(z, 1)$  depending only on  $\alpha$ . If this fails for a given value of  $\alpha$ , then there is a sequence of examples  $B_{n,k}$ , constants  $\zeta_{n,k}$  (for the given value of  $\alpha$ ) going to zero as  $n \rightarrow \infty$  and converging as  $n \rightarrow \infty$  to limits  $(X_k, z_k)$  such that the area of the  $B(z_k, 1)$  go to zero as  $k \rightarrow \infty$ . Taking a subsequence in  $k$  we can assume that the  $(X_k, z_k)$  converge in the Gromov-Hausdorff sense to a limit  $(X, z)$ . Since the areas of the  $B(z_k, 1)$  are converging to zero,  $X$  has dimension 1. Taking an appropriate diagonal sequence we have a sequence  $B_{n(k),k}$  and constants  $\zeta_{n(k),k}$  converging to zero such that the rescaled balls  $3\zeta_{n(k),k}^{-1}$  converge to  $(X, z)$ . This contradicts what we just established.

Once we have a universal lower bound to the area of  $B(z, 1)$  it follows by volume comparison (since the curvature of  $X$  is  $\geq 0$ ), that the area of any  $B(y, 1)$  for any  $y \in B(z, 1/2)$  is also universally bounded below by a positive constant depending only on  $\alpha$ .  $\square$

The following result is a 2-dimensional analogue. The statement and proof are taken from [33] and are included for completeness.

**Proposition 9.49.** *Suppose that  $B_n = B(x_n, 1)$  is a sequence of Alexandrov balls of radius 1 with curvature  $\geq k$ . Suppose that the  $B_n$  are non-compact and converge to an Alexandrov ball  $B = B(x, 1)$ . Suppose that  $\dim B$  is 2 and  $\text{diam} T_x B < \pi$ . Then, after passing to a subsequence, there are points  $\hat{x}_n \in B_n$  with  $d(x_n, \hat{x}_n) \rightarrow 0$  as  $n \rightarrow \infty$  and  $r > 0$  independent of  $n$  such that one of the following holds:*

1.  $d(\hat{x}_n, \cdot)$  has no critical points in  $B(\hat{x}_n, r) \setminus \{\hat{x}_n\}$ . In this case  $B(\hat{x}_n, r')$  is homeomorphic to the tangent cone at  $\hat{x}_n$  and if  $B_n$  is a smooth manifold then  $\overline{B(\hat{x}_n, r')}$  is diffeomorphic to a closed ball in Euclidean space for every  $0 < r' < r$ .
2. There is a sequence of positive constants  $\zeta_n \rightarrow 0$  as  $n \rightarrow \infty$  such that:
  - (a) Every critical point of  $d(\hat{x}_n, \cdot)$  in  $B(\hat{x}_n, r)$  is within distance  $\zeta_n$  of  $\hat{x}_n$ , and
  - (b) there is a critical point  $q_n$  for  $d(\hat{x}_n, \cdot)$  at distance  $\zeta_n$  from  $\hat{x}_n$ .

In this case, passing to a subsequence there is a limit of the  $\zeta_n^{-1}B(\hat{x}_n, r)$ . This limit is a complete Alexandrov space of curvature  $\geq 0$  and of dimension strictly greater than 2.

*Proof.* For any  $\epsilon > 0$  there is an  $\epsilon$ -net in  $T_x B$  consisting of tangent vectors to geodesics  $\gamma_1, \dots, \gamma_N$  in  $B$  with other end points  $y^1, \dots, y^N$ . Choosing these geodesics to be short enough we can assume that  $\tilde{Z}y^i x y^j \geq \epsilon/2$  for all  $i \neq j$ . We let  $f = \frac{1}{N} \sum_{i=1}^N d(y^i, \cdot)$ . Also, assuming that  $\epsilon > 0$  is sufficiently small and the geodesics are sufficiently short,  $f$  has a local max at  $x$  and this is the only local max for  $f$  in  $B(x, r)$  for some  $r > 0$ .

For each  $1 \leq j \leq N$ , let  $y_n^j$  be a sequence converging to  $y^j$ . We define  $f_n = \frac{1}{N} \sum_{j=1}^N d(y_n^j, \cdot)$ . Then for all  $n$  sufficiently large there is  $\hat{x}_n \in B_n$  which is a local maximum for  $f_n$  and  $\hat{x}_n \rightarrow x$  as  $n \rightarrow \infty$ . In particular,  $d(x, \hat{x}_n) \rightarrow 0$  as  $n \rightarrow \infty$ . We consider the distance function  $d(\hat{x}_n, \cdot)$ . Fix  $r > 0$  less than  $1/2$  the minimum length of the  $\gamma_i$  and chosen such that the distance function  $d(x, \cdot)$  is regular on  $B(x, r') \setminus \{x\}$ . Let  $C(n)$  be set of critical points for  $d(\hat{x}_n, \cdot)$  contained in  $B(\hat{x}_n, r) \setminus \{\hat{x}_n\}$ . Then, passing to a subsequence, either  $C(n) = \emptyset$  for all  $n$  or there is a sequence  $\zeta_n > 0$  tending to zero such that  $C(n) \subset \overline{B(\hat{x}_n, \zeta_n)}$  and there is a point  $q_n \in C(n)$  at distance  $\zeta_n$  from  $\hat{x}_n$ . In the first case  $B(\hat{x}_n, r)$  satisfies the first conclusion of the proposition for all  $n$  and the proof is complete. In the second case, we choose geodesics  $\nu_n$  from  $\hat{x}_n$  to  $q_n$ . Now consider the sequence  $\zeta_n^{-1}(B(\hat{x}_n, r), \hat{x}_n)$  and pass to a subsequence with a limit  $(Z, z)$ . This is a complete Alexandrov space of curvature  $\geq 0$ . Our goal is to show that the dimension of  $Z$  is greater than 2, so we suppose that  $Z$  has dimension 2. Passing to a further subsequence we can suppose that the  $q_n$  converge to a point  $q \in Z$  at distance 1 from  $z$ , and the geodesics  $\nu_n$  converge to a geodesic  $\nu$  from  $z$  to  $q$ . Denote by  $\tau$  the direction of  $\nu$  at  $z$ .

Consider the set  $R^j$  all geodesic rays emanating from  $z$  that are limits as  $n \rightarrow \infty$  of geodesics from  $\hat{x}_n$  to  $y_n^j$ . Fix momentarily geodesics  $\gamma^j \in R^j$ . For  $a < \infty$  let  $u^j$  be the point on  $\gamma^j$  at distance  $a$  from  $z$ . Since  $q$  is a critical point for  $d(z, \cdot)$  we have  $\tilde{Z}z q u^j \leq \pi/2$  for all  $j$ . Since  $\tilde{Z}u^j z q \leq \text{Arcsin}(1/a)$  and since  $Z$  has curvature  $\geq 0$ , this implies that  $\tilde{Z}u^j q z \geq \pi/2 - \text{Arcsin}(1/a)$  for all  $j \leq N$ . This shows that the distance in  $S_z Z$  between  $\tau$  and the directions  $[R^j]$  tangent to the geodesics  $R^j$  is at least  $\pi/2 - \text{Arcsin}(1/a)$ . Also, by construction the distance in  $S_z Z$  between  $[R^j]$  and  $[R^{j'}]$  is  $\geq \epsilon/2$  for all  $j \neq j'$ .

Given  $\epsilon > 0$  we can choose  $a$  sufficiently large, so that there are at most two points separated by distance at least  $\epsilon/2$  contained in the closed annular region

$\overline{B(\tau, \pi/2 + 2\text{Arcsin}(1/a))} \setminus B(\tau, \pi/2 - \text{Arcsin}(1/a))$  in  $S_z Z$ . Thus, all but at most two of the  $[R^j]$  have distance greater than  $\pi/2 + 2\text{Arcsin}(1/a)$  from  $\tau$ , and the remaining (at most two) have distance at least  $\pi/2 - \text{Arcsin}(1/a)$  from  $\tau$ . This means for all  $n$  sufficiently large, all but two of the  $N$  the distances at  $S_{\hat{x}_n} B_n$  between the direction  $[\nu_n]$  and the direction  $[\gamma_n^j]$  of any geodesic from  $\hat{x}_n$  to  $y_n^j$  are greater than  $\pi/2 + 2\text{Arcsin}(1/a)$  and for the remaining at most two values of  $j$  these distances of  $[R^j]$  from  $\tau$  are at least  $\pi/2 - 2\text{Arcsin}(1/a)$ . This implies that for all  $n$  sufficiently large, the directional derivative of  $f_n$  at  $\hat{x}_n$  is positive in the  $\tau$  direction and hence  $f_n$  does not have a maximum at  $\hat{x}_n$ , contrary to assumption.  $\square$

### 9.9 Gromov-Hausdorff limits of balls in the $M_n$

Now we turn from generalities about Alexandrov spaces to special properties of Gromov-Hausdorff limits of balls in the  $M_n$ . Recall that we have a sequence of constants  $w_n \rightarrow 0$  as  $n \rightarrow \infty$  and functions  $\rho_n: M_n \rightarrow [0, \infty)$  with the property that  $\rho_n(x) \leq \text{diam}(M_n^0)$  for every  $n$  and every  $x$  in the connected component  $M_n^0$  of  $M_n$ . Thus, for every  $n$  and every  $x \in M_n$ , the ball  $B_{g_n}(x, \rho_n(x))$  is non-compact. Since  $M_n$  is itself compact, it follows that for every  $0 < r < \rho_n(x)$ , the ball  $B_{g_n}(x, r)$  has compact closure in  $B(x, \rho_n(x))$ . It then follows that the  $B_{g_n}(x, \rho_n(x))$  are Alexandrov balls. Rescaling the metric by  $\rho_n(x)^{-2}$ , that is to say replacing the metric  $g_n$  on this ball by the metric  $g'_n(x) = \rho_n^{-2}(x)g_n$  we obtain non-compact Alexandrov balls  $B_{g'_n(x)}(x, 1)$  of radius 1 with the property that their sectional curvatures are bounded below by  $-1$ , and their volumes are bounded above by  $w_n$ . Since  $w_n \rightarrow 0$  as  $n \rightarrow \infty$ , the following is then immediate from Proposition 9.33 and Claim 9.47.

**Proposition 9.50.** *Let  $x_n \in M_n$  be given for every  $n \geq 1$ . Then, after passing to a subsequence, the  $B_{\rho_n^{-2}(x_n)g_n}(x_n, 1)$  converge to an Alexandrov ball  $B = B(\bar{x}, 1)$  of curvature  $\geq -1$  and of dimension 1 or 2. The limiting ball contains points at every distance  $< 1$  from  $\bar{x}$ .*

This leads immediately to the following corollary.

**Corollary 9.51.** *There is a decreasing sequence of constants  $\epsilon_n > 0$  tending to zero as  $n \rightarrow \infty$  such that for every  $n$  and for any  $x_n \in M_n$  there is an Alexandrov ball  $B$  of radius 1, of curvature  $\geq -1$ , and of dimension 1 or 2, such that  $B_{\rho^{-2}(x_n)g_n}(x_n, 1)$  is within  $\epsilon_n$  in the Gromov-Hausdorff distance of  $B$ .*

## 10 2-dimensional Alexandrov spaces

In order to get enough information about the structure of balls in the  $M_n$  limiting (after rescaling) to a 2-dimensional Alexandrov ball, we need fairly delicate information about 2-dimensional Alexandrov balls. We shall fix an appropriate  $\delta > 0$  sufficiently small and show that there is always a cover a 2-dimensional ball by four types of neighborhoods (for more details, see Theorem 10.30):

1. balls near flat balls in  $\mathbb{R}^2$ ,

2. balls near flat circular cones of cone angle  $\leq 2\pi - \delta$ ,
3. balls near flat cones in  $\mathbb{R}^2$  of cone angle  $\leq \pi - \delta$ , and
4. balls near flat boundary points.

The important facts about 2-dimensional Alexandrov balls that will be used in establishing the results are the following:

1. If a 2-dimensional Alexandrov ball is nearly flat at one scale then it is nearly flat at all smaller scales.
2. If a 2-dimensional Alexandrov ball is close to a circular cone of angle  $\geq \alpha$  then it has an annular region which is fibered by the circles that are metric spheres. This annular region is nearly flat on scales depending only on  $\alpha$ .
3. If a 2-dimensional Alexandrov ball has nearly flat boundary at some point on one scale then the same is true on all smaller scales.
4. If a 2-dimensional Alexandrov ball is close to a flat cone of angle  $\geq \alpha$  then there is a region that is a topological product foliated by metric spheres that are intervals with endpoints in the boundary. Further, every point of intersection of this region with the boundary is a nearly flat boundary point and the interior point of this annular region are nearly flat on a scale which is determined by  $\alpha$  and the distance to the boundary.

Establishing these results is the subject of this section.

### 10.1 Basics

**Claim 10.1.** *A 2-dimensional Alexandrov ball  $X$  is a topological 2-manifold, possibly with boundary. The topological boundary of  $X$  is Alexandrov boundary  $\partial X$ .*

*Proof.* For a proof, see §12.9.3 of [3]. □

Let  $X$  be a 2-dimensional Alexandrov ball. We define the *cone angle* at any point  $p \in X$  to be the total length of the tangent sphere  $\Sigma_p$ . It follows from the Alexandrov space axioms that if  $p \in \text{int } X$  then the cone angle at  $p$  is at most  $2\pi$  and the tangent cone is a flat circular cone of this cone angle. If  $p \in \partial X$ , then the cone angle at  $p$  is at most  $\pi$ , and the tangent cone is a sub-cone of  $\mathbb{R}^2$  of this cone angle.

**Lemma 10.2.** *Suppose that  $(X_n, x_n)$  is a sequence of 2-dimensional Alexandrov balls converging to a 2-dimensional Alexandrov ball  $(X, x)$  and suppose that  $y_n \in X_n$  converges to  $y \in X$ . Then:*

1. *If  $y_n \in \partial X_n$  for all  $n$ , then  $y \in \partial X$ .*
2. *Conversely, if  $y \in \partial X$ , then there is a sequence  $z_n \in \partial X_n$  converging to  $y$ .*

*Proof.* We begin with a proof of the first statement. Let us suppose to the contrary that  $y_n \in \partial X_n$  for all  $n$  and that  $y \in \text{int } X$ . Let  $d_n$  be the Gromov-Hausdorff distance from  $(X_n, y_n)$  to  $(X, y)$ . Choose constants  $\lambda_n \rightarrow \infty$  such that  $\lambda_n d_n \rightarrow 0$ . Then the Gromov-Hausdorff distance from  $(\lambda_n X_n, y_n)$  to  $(\lambda_n X, y)$  goes to zero and the  $(\lambda_n X, y)$  converge to the tangent cone to  $X$  at  $y$ . This allows us to assume that the  $(X_n, y_n)$  converge to  $(C, y)$  where  $C$  is a circular cone and  $y$  is the cone point.

**Claim 10.3.** *Given the cone  $C$  there is a positive function  $s(d)$  defined for  $0 < d < \infty$  and for each  $\epsilon > 0$  there is  $\delta > 0$  such that at each point  $z$  in the metric sphere  $S(y, d) = \{w \in C \mid d(y, w) = d\}$  there is a  $(2, \delta)$ -strainer  $\{a_1(z), a_2(z), b_1(z), b_2(z)\}$  with  $a_1(z) = y$  and with the following property. Setting  $f = d(a_1(z), \cdot)$  and  $g = d(b_1(z), \cdot)$ , then  $(f, g)$  defines a homeomorphism from a neighborhood of  $z \in C$  to an open square  $R$  in  $\mathbb{R}^2$  of side length  $s(d(y, z))$ , a homeomorphism that is a  $(1 + \epsilon)$  almost isometry.*

*Proof.* This follows from direct computation in the flat cone  $C$ . □

By compactness, for any  $t > 0$  and any  $\epsilon > 0$  we can find a finite number of points  $z_1, \dots, z_k$  in  $S(y, t)$  with  $(2, \delta)$ -strainers  $\{a_1(z_i) = y, a_2(z_i), b_1(z_i), b_2(z_i)\}$  and  $(1 + \epsilon)$ -almost isometries  $(f_i, g_i): U(z_i) \rightarrow R$  as in the claim with the  $U(z_i)$  covering  $S(y, t)$ .

Now we pass from the cone to the sequence  $X_n$ . We choose sequences  $z_{n,i} \in X_n$  converging to  $z_i$  and points  $a_1(z_{n,i}) = y_n, a_2(z_{n,i}), b_1(z_{n,i}), b_2(z_{n,i})$  in  $X_n$  converging to  $a_1(z_i) = y, a_2(z_i), b_1(z_i), b_2(z_i)$ . For all  $n$  sufficiently large we have a neighborhood  $U(z_{n,i})$  and a function  $(f_{n,i}, g_{n,i}): U(z_{n,i}) \rightarrow R$ , both defined analogously to the ones for  $C$ . For all  $n$  sufficiently large the functions  $(f_{n,i}, g_{n,i})$  are a  $(1 + \epsilon)$ -almost isometries for every  $i$ . Taking limits we see that for all  $n$  sufficiently large,  $\cup_{i=1}^k U(z_{n,i})$  covers the metric sphere  $S(y_n, t)$ . The following results now follow easily from this by standard arguments.

**Claim 10.4.** *For any  $0 < d < 1$  and any  $\epsilon > 0$  the following holds for all  $n$  sufficiently large.*

1. *For each  $t \in (d, 1)$  the metric sphere  $S(y_n, t)$  is a simple closed, rectifiable curve whose length is between  $(1 - \epsilon)$  and  $(1 + \epsilon)$  of the length of  $S(y, t) \subset C$ .*
2.  *$d(y_n, \cdot): B(y_n, 1) \setminus B(y_n, d) \rightarrow [d, 1)$  is the projection of a product structure. The fibers of this projection are the metric spheres  $S(y_n, t)$ ,  $d \leq t < 1$ .*

*Proof.* The first statement is clear from the existence of the boxes  $U(z_{n,i})$  converging to the  $U(z_i)$  and the almost isometries to  $R$ . Let us consider the second statement. For any  $t > 0$ , for all  $n$  sufficiently large the intersection of  $U(z_{n,i})$  with the metric spheres  $S(y_n, s)$  are vertical lines. This provides a local product structure in  $U(z_{n,i})$  whose projection onto a factor is given by  $d(y_n, \cdot)$ . It is easy to patch these local product structures together to give a local product structure in a neighborhood of  $S(y_n, t)$  whose projection to a factor is given by  $d(y_n, \cdot)$ . For any  $0 < d < 1$ , provided that  $n$  is sufficiently large, this result holds for all  $t \in [d, 1)$  and the local product



structures around the  $S(y_n, t)$  fit together to give a product structure as required on  $B(y_n, 1) \setminus B(y_n, d)$ .  $\square$

The previous claim has two important consequences:

**Claim 10.5.** *Fix  $0 < \delta < 1$ . Then the following hold for all  $n$  sufficiently large. There is an infinite cyclic covering  $\tilde{B}$  of  $B(y_n, 1)$  determined by a surjective homomorphism  $\pi_1(B(y_n, 1), y_n) \rightarrow \mathbb{Z}$  and an element  $a \in \pi_1(B(y_n, 1), y_n)$  that maps to a generator of  $\mathbb{Z}$  and is represented by a loop based at  $y_n$  of length less than  $\delta$ .*

*Proof.* Take a geodesic  $L$  in  $X_n$  joining  $y_n$  to the metric sphere  $S(y_n, 1)$ . This geodesic represents a relative homology class in  $Y_n = \bar{B}(y_n, 1)$  modulo the union of  $\partial(X_n \cap B(y_n, 1))$  and  $S(y_n, 1)$ . Taking the intersection number with this geodesic defines a homomorphism  $\iota$  from  $H_1(Y_n)$  to  $\mathbb{Z}$  and hence an infinite cyclic covering  $p: \tilde{Y}_n \rightarrow Y_n$ . Let  $\partial^0(X_n)$  denote the boundary component of  $X_n$  that contains  $y_n$ . Since it is contained in the  $1/2$ -neighborhood of  $y_n$ , it is a circle and has intersection number 1 with  $L$ , showing that  $\iota$  is surjective. For any  $0 < d$  and, given  $d$ , for all  $n$  sufficiently large, it also unwraps the metric spheres near the non-compact end of  $B(y_n, 1)$  to copies of  $\mathbb{R}$ . Now fix  $r > 0$  sufficiently small. For any  $0 < d$  for all  $n$  sufficiently large (given  $d$ ), the class of the metric sphere  $S(y_n, d)$  also generates the covering transformation. The length of this circle is at most twice the length of the corresponding circle in the cone (and hence is at most  $4\pi d$ ). Thus, the generating covering transformation moves any point on the pre-image of  $S(y_n, d)$  a distance at most  $4\pi d$ . Hence, there is an element  $a \in \pi_1(B(y_n, 1), y_n)$  that maps to a generator under the homomorphism to  $\mathbb{Z}$  and is represented by a loop based at  $y_n$  of length at most  $2d + 4\pi d$ . Choosing  $d > 0$  less than  $\delta/(2 + 4\pi)$  gives the last statement in the claim.  $\square$

On the other hand, for  $\ell_0 > 0$  as in Proposition 9.36 there is  $r < \ell_0$  and a point  $z \in C$  within distance  $\ell_0$  of the cone point such that  $B(z, r)$  is simply connected. Then, for every  $n$  sufficiently large there is a point  $z_n \in B(y_n, \ell_0)$  such that the composition  $\pi_1(B(z_n, r)) \rightarrow \pi_1(B(y_n, 1), y_n) \rightarrow \mathbb{Z}$  is trivial. Thus, fixing  $\epsilon > 0$  and  $N > N_2(\ell_0, r, \epsilon)$  from Proposition 9.36 we take  $\delta = \epsilon/N$ . Then for all  $n$  sufficiently large the powers  $a^k$  for  $-N \leq k \leq N$  are represented by loops based at  $y_n$  of length less than  $\epsilon$ . These map to distinct elements in  $\mathbb{Z}$ . Now applying Proposition 9.36 we see get a contradiction for all  $n$  sufficiently large. This completes the proof of the first statement.

We turn now to the second statement. Suppose that  $y \in \partial X$  and  $y_n \in X_n$  converges to  $y$ . We shall show that  $d(y_n, \partial X_n)$  goes to zero. If that is true we simply replace the sequence  $y_n$  with a sequence  $z_n \in \partial X_n$  with the same limit. So suppose to the contrary that there is  $d > 0$  such that  $d(y_n, \partial X_n) \geq d$ . By rescaling exactly as above, we can assume that the  $X_n$  converge to a flat cone  $C$  in  $\mathbb{R}^2$ , the  $y_n$  converge to the cone point, and that the distance from  $y_n$  to  $\partial X_n$  goes to infinity. The distance function from the cone point is regular in the complement of the cone point and the level sets are arcs with endpoints in the boundary of the cone. Fix  $d > 0$ . It follows that for all  $n$  sufficiently large, the distance function

from  $y_n$  is regular on  $B(y_n, 2d) \setminus B(y_n, d)$ . According to Theorem 12.7 of [3] the level sets of the distance function from  $y_n$  in this range are topological one-manifolds with boundary in the boundary of  $X_n$ . Since the distance from  $y_n$  to  $\partial X_n$  goes to infinity as  $n \rightarrow \infty$ , this implies that for all  $n$  sufficiently large, the metric sphere  $S(y_n, t)$  is disjoint union of simple closed curves for any  $t \in (d, 2d)$ . On the other hand, the exact same arguments as above constructing boxes almost isometric to squares in  $\mathbb{R}^2$  apply away from the endpoints of  $S(y, t)$ <sup>8</sup>. This means that for any  $\epsilon > 0$  sufficiently small and, given  $\epsilon > 0$ , for all  $n$  sufficiently large there is an open covering of  $S(y_n, t)$  consisting of  $U_+, U_-, J$  where  $U_+$  and  $U_-$  are the (disjoint) subsets of points of  $S(y_n, t)$  within  $\epsilon$  of the endpoints  $p_+$  and  $p_-$  of  $S(y, t)$  and  $J$  is an interval with one end in  $U_+$  and the other in  $U_-$ . Clearly, this is a contradiction, since no disjoint union of simple closed curves has such an open cover.  $\square$

**Corollary 10.6.** *Suppose that  $X_n$  are 2-dimensional Alexandrov balls converging to a 2-dimensional Alexandrov ball  $X$ . Suppose that  $x_n \in X_n$  converge to  $x \in X$ . Let  $d_n$  be the distance from  $x_n$  to  $\partial X_n$  and let  $d$  be the distance from  $x$  to  $\partial X$ . Then  $d = \lim_{n \rightarrow \infty} d_n$ .*

There is another consequence of this result that will be important later. It is established by a standard limiting argument.

**Lemma 10.7.** *Given  $a > 0$  there is  $\delta = \delta(a) > 0$  such that the following holds. Suppose that  $B(\bar{y}, 1/2)$  and  $B(\bar{y}', 1/2)$  are 2-dimensional Alexandrov balls of area  $\geq a/8$  and curvature  $\geq -1$  with  $\bar{y} \in \partial B(\bar{y}, 1/2)$ . If the Gromov-Hausdorff distance between  $B(\bar{y}, 1/2)$  and  $B(\bar{y}', 1/2)$  is less than  $\delta$ , then  $\bar{y}$  is within distance  $(0.1)$  of  $\partial B(\bar{y}', 1/2)$ .*

Lastly, we need a uniform area estimate for sub-balls of a ball with given area.

**Lemma 10.8.** *Given  $a > 0$  there is  $a' = a'(a)$  with  $0 < a'(a) \leq a$  such that the following holds for any 2-dimensional Alexandrov ball  $B(x, 1)$  of curvature  $\geq -1$  and area  $\geq a$ . For any  $y \in B(x, 15/16)$  and any  $r < 1/16$  the area of  $B(y, r)$  is at least  $a'r^2$ .*

*Proof.* It suffices to prove this result for  $r = 1/16$ , since by the Bishop-Gromov volume comparison, it then follows for any  $r \leq 1/16$  (with a different constant  $a'$ ). The result for  $r = 1/16$  follows by the usual limiting argument.  $\square$

## 10.2 The Interior

We approximate interior points by cones, including flat cones.

**Definition 10.9.** Fix  $\mu > 0$ . Let  $X$  be an 2-dimensional Alexandrov ball of curvature  $\geq -1$ . Then  $X$  is *interior  $\mu$ -good at a point  $y \in \text{int } X$  of angle  $\alpha$  and on scale  $r$*  if  $B_{r-2g}(y, 1)$  is within  $\mu$  in the Gromov-Hausdorff distance of the unit ball centered

<sup>8</sup>It follows from the second item of Proposition 10.18 that this argument works up to the boundary.

at the cone point in the circular cone of cone angle  $\alpha$ . We say that  $X$  is *interior  $\mu$ -flat at  $y$  on scale  $s$*  if  $B_{s^{-2}g}(y, 1)$  is within  $\mu$  in the Gromov-Hausdorff distance of the unit ball in  $\mathbb{R}^2$ .

We need to establish the relationship between being  $\mu$ -flat and having a  $(2, \delta)$ -strainer.

**Lemma 10.10.** *1. Given  $\delta > 0$  there are  $\mu > 0$  and  $d > 0$  such that if  $B$  is an Alexandrov ball of curvature  $\geq -1$  that is interior  $\mu$ -flat at  $y \in B$  on some scale  $d' \leq d$  then there is a  $(2, \delta)$ -strainer of size  $d'$  centered at  $y$ .*

*2. Given  $\mu > 0$  there is  $\delta > 0$  and  $R < \infty$  such that if  $B$  is an Alexandrov ball of curvature  $\geq -1$  and if there is a  $(2, \delta)$ -strainer of size  $d$ , for some  $0 < d \leq 1$ , at  $y \in B$ , then  $B$  is interior  $\mu$ -flat at  $y$  on scale  $d/R$ .*

*Proof.* If the first does not hold for some  $\delta > 0$  there are sequences  $\mu_k, d'_k \rightarrow 0$  and counter examples  $y_k \in B_k$  at scale  $d'_k$ . The unit balls  $(d'_k)^{-1}B(y_k, d'_k)$  converge to the unit ball in  $\mathbb{R}^2$  and the  $(d'_k)^{-1}B(y_k, d'_k)$  are Alexandrov spaces of curvature  $\geq -(d'_k)^2$ . Of course, there is a  $(2, \delta)$ -strainer of size 1 at the origin in the unit ball in  $\mathbb{R}^2$ . Using the upper semi-continuity of comparison angles under limits we see that for all  $k$  sufficiently large there is a  $(2, \delta)$ -strainer of size 1 at  $y_k$ . The contradiction establishing the first result follows by rescaling.

If the second does not hold for some  $\mu > 0$ , then there are sequences  $\delta_k \rightarrow 0$  and  $R_k \rightarrow \infty$  and  $0 < d_k \leq 1$  and counter examples  $y_k \in B_k$  for these values. The balls  $(R_k/d_k)B(y_k, d_k)$  have  $(2, \delta_k)$ -strainers of size  $R_k/2$  and hence these balls converge to  $\mathbb{R}^2$ . This means that the  $(R_k/d_k)B(y_k, d_k/R_k)$  converge to the unit ball in  $\mathbb{R}^2$  and hence  $B$  is interior  $\mu$ -flat at  $y_k$  on scale  $d_k/R_k$  for all  $k$  sufficiently large, which is a contradiction.  $\square$

The next thing to notice is that being interior  $\mu$ -flat at one scale implies interior flatness at all smaller scales.

**Lemma 10.11.** *Given  $\mu > 0$  there is  $\nu > 0$  such that the following holds. If an Alexandrov ball  $X = B(x, 1)$  of curvature  $\geq -1$  is interior  $\nu$ -flat at  $x$  on scale  $\ell$  for some  $0 < \ell \leq 1$ , then the ball  $X$  is interior  $\mu$ -flat at  $x$  on all positive scales  $\leq \ell$ .*

*Proof.* Suppose that the result does not hold for some  $\mu$ . Then there are sequences  $\nu_n$  and  $\ell_n$  with  $\nu_n$  tending to 0 as  $n \rightarrow \infty$  and  $X_n = B(x_n, 1)$  which are interior  $\nu_n$ -flat at  $x_n$  on scale  $\ell_n$  but not interior  $\mu$ -flat at some scale  $0 < s_n < \ell_n$ . Since  $\nu_n \rightarrow 0$ , the sequence  $\ell_n^{-1}B(x_n, \ell_n)$  converges to the unit ball in  $\mathbb{R}^2$ . Passing to a subsequence we arrange that the  $s_n/\ell_n$  converge to a limit  $s$ , with  $0 \leq s \leq 1$ . If  $s > 0$  then the  $s_n^{-1}B(x_n, \ell_n)$  converge to the ball in  $\mathbb{R}^2$  of radius  $s^{-1}$ , which implies that the  $s_n^{-1}B(x_n, s_n)$  converge to the unit ball in  $\mathbb{R}^2$ , which is a contradiction. If the  $s_n/\ell_n$  converge to 0, then for each  $\delta > 0$  and  $d < \infty$  for all  $n$  sufficiently large, in  $s_n^{-1}B(x_n, \ell_n)$  there is a  $(2, \delta_n)$ -strainer of size  $d$  centered at  $x_n$ , where  $\delta_n \rightarrow 0$  as  $n \rightarrow \infty$ . This means that the  $s_n^{-1}(X_n, x_n)$  converge to  $(\mathbb{R}^2, 0)$ , and hence the unit balls converge to the unit ball in  $\mathbb{R}^2$ . This is a contradiction.  $\square$

Now, we show that interior good at a point implies locally interior flat in a nearby annular region where the constants depend on the area. See FIG. 1.

**Proposition 10.12.** *Given  $\mu > 0$  and  $a' > 0$ , there are positive constants  $s_0 = s_0(a')$  and  $\mu'(\mu, a')$  such that for all  $0 < \mu' \leq \mu'(\mu, a')$  the following holds. Suppose that a 2-dimensional Alexandrov ball  $X = B(x, 1)$  of curvature  $\geq -1$  and area  $\geq a'$  is interior  $\mu'$ -good at  $x$  on scale 1. Then  $X$  is interior  $\mu$ -flat at every point  $y \in B(x, 7/8) \setminus B(x, 1/8)$  on all scales  $\leq s_0$ . Furthermore, for every  $b \in (1/8, 7/8)$  the metric sphere  $S(x, b)$  is a simple closed curve, and the closed metric ball  $\overline{B(x, b)}$  is homeomorphic to a disk.*

*Proof.* Fix  $\mu > 0$ . Suppose we have a sequence  $\mu'_n \rightarrow 0$  and Alexandrov balls  $B(x_n, 1)$  as in the statement. Then passing to a subsequence these converge to a flat circular cone  $(C, p)$  of area  $\geq a'$ . There is  $0 < s_0$ , depending only on  $a'$ , such that at every point of  $\overline{B(p, 7/8)} \setminus B(p, 1/8)$  the cone  $C$  is interior flat at all scales  $\leq s_0$ . Suppose that for each  $n$  there is a point  $y_n \in (B(x_n, 7/8) \setminus B(x_n, 1/8))$  at which  $B_n$  is not interior  $\mu$ -flat on all scales  $\leq s_0$ . Then  $s_0^{-1}(B(y_n, s_0))$  converges to a unit ball in  $\mathbb{R}^2$  and arguing as in the previous result, Lemma 10.11, we see that for all  $n$  sufficiently large  $B_n$  is  $\mu$ -flat at  $y_n$  on all scales  $\leq s_0$ . This is a contradiction, and the first statement follows immediately.

The function  $d(p, \cdot)$  is regular on the annular region  $A = B(p, 7/8) \setminus B(p, 1/8)$  in the cone  $C$ , and in fact for every  $y \in A$  there is a direction  $\tau$  at  $y$  with the directional derivative of the distance from  $p$  in the  $\tau$ -direction equal to 1. Thus, given  $\delta > 0$ , provided that  $\mu'$  sufficiently small, the distance  $d(x, \cdot)$  is regular on the annular region  $A' = B(x, 7/8) \setminus B(x, 1/8)$ , and indeed at every  $y' \in A'$  there is a direction  $\tau'$  so that the directional derivative of  $d(x, \cdot)$  in the  $\tau'$ -direction is at least  $1 - \delta$ . It then follows from §11 of [3] and the arguments given in the proof of Lemma 10.2 that, provided that  $\mu'$  is sufficiently small,  $S(x, b)$  is a simple closed curve and the closed region bounded by  $S(x, 1/8)$  and  $S(x, 7/8)$  is homeomorphic to a product  $S^1 \times I$ .

Now let us show that, possibly after making  $\mu' > 0$  smaller, the closed metric balls  $\overline{B(x, b)}$  are homeomorphic to closed disks. If not then there is a sequence of counter examples  $B(x_k, 1)$  within distance  $\mu'_k$  of circular cones for a sequence of  $\mu'_k \rightarrow 0$  as  $k \rightarrow \infty$ . Passing to a subsequence we can assume that the  $B(x_k, 1)$  converge to an Alexandrov space  $B_\infty$ . By the uniform lower bound on the areas,  $B_\infty$  is 2-dimensional and hence is a circular cone. If the cone angle of  $B_\infty$  is less than  $2\pi$ , invoking Proposition 9.49 we see that there are points  $\hat{x}_k \in B(x_k, 1)$  also converging to the cone point such that for all  $k$  sufficiently large the distance function  $d(\hat{x}_k, \cdot)$  has no critical points in  $B(\hat{x}_k, 1/2)$ . (The other possible result according to Proposition 9.49 is that there is a rescaling of the balls that converges to a limit of dimension greater than 2. But, this is absurd since the balls in the sequence all have dimension 2.) It follows that the closed metric balls  $\overline{B(\hat{x}_k, b)}$  are homeomorphic to disks for all  $b \in (0, 1/2)$ . Now for  $k$  sufficiently large,  $S(x_k, 3/8)$  separates  $S(\hat{x}_k, 1/4)$  and  $S(\hat{x}_k, 1/2)$  and hence the region between  $S(\hat{x}_k, 1/4)$  and  $S(x_k, 3/8)$  is homeomorphic to a product. This implies that  $\overline{B(x_k, 3/8)}$  is homeomorphic to a disk. Since  $d(x_k, \cdot)$  is regular on  $(1/8, 7/8)$  all the closed metric balls  $\overline{B(x_k, b)}$  for

$b \in (1/8, 7/8)$  are homeomorphic to closed disks. This is a contradiction, proving the result follows in this case.

Now suppose that  $B_\infty$  has cone angle  $2\pi$ , i.e., suppose that it is a disk in  $\mathbb{R}^2$ . Fix  $\delta > 0$  sufficiently small. Then for all  $k$  sufficiently large there is a  $(2, \delta)$ -strainer at  $x_k$  of size  $1/2$ . Hence, for all these  $k$  there is a bi-Lipschitz homeomorphism from a ball in  $\mathbb{R}^2$  whose radius is independent of  $k$  to a neighborhood of  $x_k$  whose image contains a fixed size metric ball about  $x_k$ . It then follows that this fixed size metric ball has closure that is homeomorphic to a disk. Since as  $k \rightarrow \infty$  all critical points for the distance function from  $x_k$  are arbitrarily close to  $x_k$ , again we achieve a contradiction for all  $k$  sufficiently large, proving the result in this case.  $\square$

**Definition 10.13.** If  $B(x, 1)$  satisfies the statement in the above proposition, then we say that  $B(x, 7/8) \setminus B(x, 1/8)$  is a  $(\mu, s_0)$ -good annular region. (See FIG. 1). Notice that for any  $s'_0 < s_0$  a  $(\mu, s_0)$ -good annular region is automatically a good  $(\mu, s'_0)$ -good annular region.

### 10.3 The boundary

We turn to the analogues for the boundary of interior flatness and interior goodness.

**Definition 10.14.** Fix  $\mu > 0$ . Let  $B(x, 1)$  be a 2-dimensional Alexandrov ball of curvature  $\geq -1$  and let  $y \in X$ . We say that  $X$  is *boundary  $\mu$ -good of angle  $\alpha$  and on scale  $r$  near  $y \in X$*  if the rescaled ball  $r^{-1}B(y, r)$  is within  $\mu$  in the based Gromov-Hausdorff distance of the unit ball centered at the cone point in a (flat) 2-dimensional cone in  $\mathbb{R}^2$  of cone angle  $\alpha$ . We say that  $X$  is *boundary  $\mu$ -flat near  $y \in X$  on scale  $r$*  if  $r^{-1}B(y, r)$  is within  $\mu$  in the based Gromov-Hausdorff distance to the unit ball centered at a boundary point of  $\mathbb{R} \times [0, \infty)$ .

**Lemma 10.15.** *Given  $\mu > 0$  and  $a' > 0$  there is  $0 < \mu''_0(\mu, a') \leq \mu$  such that the following holds for all  $0 < \mu'' \leq \mu''_0(\mu, a')$ . Suppose that a 2-dimensional Alexandrov ball  $X = B(x, 1)$  of curvature  $\geq -1$  and area  $\geq a'$  is boundary  $\mu''$ -good near  $x$  on scale 1. Then for any  $b \in [1/64, 7/8]$  the metric sphere  $S(x, b)$  is an arc with endpoints in  $\partial X$  and the closed metric ball  $\overline{B(x, b)}$  is homeomorphic to a 2-disk.*

*Proof.* Fix  $\mu > 0$  and  $a' > 0$  and suppose  $\mu''_0$  is sufficiently small, and suppose that  $X = B(x, 1)$  satisfies the hypothesis of the lemma. Since the distance function from the cone point in a flat cone is regular on the corresponding annular region, assuming that  $\mu''$  is sufficiently small, the distance function from  $x$  is regular on  $B(x, 7/8) \setminus B(x, 1/64)$ . It follows from §11 of [3] and the arguments given in the proof of Lemma 10.2 that, provided that  $\mu''$  is sufficiently small, for any  $b \in (1/64, 7/8)$  the metric sphere  $S(x, b)$  is an arc with endpoints in  $\partial X$ .

We must also show that, provided that  $\mu'' > 0$  is sufficiently small, the closed metric ball  $\overline{B(x, b)}$  is homeomorphic to a disk. If there is no such  $\mu'' > 0$  with this property, then we take a sequence of counter-examples  $B_n = B(x_n, 1)$  for  $\mu''_n \rightarrow 0$ . Passing to a subsequence we can take a limit  $B = B(x, 1)$  which is a flat cone in  $\mathbb{R}^2$ . If the cone angle is less than  $\pi$  then arguing as in Lemma 10.12 we obtain a contradiction. It remains to consider the case when the limit is a flat cone of cone

angle  $\pi$ . In this case, for all  $n$  sufficiently large the distance function from  $x_n$  has no critical points outside a fixed size metric ball around  $x_n$ , the size of the ball going to zero as  $n \rightarrow \infty$ . On the other hand, the distance function  $F_n$  from a point of  $\partial B_n$  at distance  $7/8$  from  $x_n$  is regular on a fixed size metric ball about  $x_n$ . Hence, a smaller metric ball about  $x_n$  is contained in a compact region  $R_n$  of  $B_n$  that is fibered by the intersection of  $R_n$  with level sets of  $F_n$ , each of these being intervals. Thus,  $R_n$  is homeomorphic to a disk and contains a fixed size metric closed ball  $A$  about  $x_n$ , which consequently is also homeomorphic to a disk. For  $n$  sufficiently large all the critical points of the distance function from  $x_n$  within distance  $7/8$  of  $x_n$  are contained in  $A$ . Hence, the region between  $B(x_n, t) \setminus A$  is a product region for any  $t < 7/8$ . The result follows in this case as well.  $\square$

The next observation is that boundary flatness near a boundary point at one scale implies boundary flatness near that point at all smaller scales.

**Lemma 10.16.** *Given  $\mu > 0$  for all  $\nu' > 0$  sufficiently small, the following holds for any  $0 < \ell \leq 1$ . Suppose that an Alexandrov ball  $X = B(x, 1)$  of curvature  $\geq -1$  is boundary  $\nu'$ -flat near  $x$  on scale  $\ell$ .*

1. *For any  $0 < r \leq \ell$  if  $d(x, \partial X) < r\nu'$ , then the ball  $X$  is boundary  $\mu$ -flat near  $x$  on scale  $r$ . In particular, if  $x \in \partial X$ , the  $X$  is boundary  $\mu$ -flat near  $x$  on all positive scales  $\leq \ell$ .*
2. *If  $y \in \text{int} X \cap B(x, 7\ell/8)$ , then  $X$  is interior  $\mu$ -flat at  $y$  on all positive scales  $\leq \min(\ell/8, d(y, \partial X))$ .*

*Proof.* It follows from Lemma 10.15 that provided that  $\nu$  is sufficiently small  $\partial B(x, 15/16)$  is an arc and each end of this arc is at distance  $15/16$  from  $x$ .

**Claim 10.17.** *Fix  $\beta > 0$ . The following holds for all  $\nu' > 0$  sufficiently small. Suppose an Alexandrov  $X = B(x, 1)$  of curvature  $\geq -1$  with  $x \in \partial X$  is boundary  $\nu'$ -flat near  $x$  on scale 1. Then for any  $0 < r < 7/8$ , fixing  $e_+$  and  $e_-$  on  $\partial X$  at distance  $r$  from  $x$ , and on opposite sides of  $x$  on  $\partial X$ , the comparison angle  $\tilde{\angle} e_+ x e_-$  is greater than  $\pi - \beta$ .*

*Proof.* If  $\nu' > 0$  is sufficiently small, then there are points at distance  $\max(r, 7/8)$  from  $x$  with this property. The result follows from the fact that as we move points  $e_+, e_-$  toward  $x$  along  $\partial X$ , the comparison angle is weakly monotone increasing.  $\square$

Let us prove the first statement in the lemma. If this statement does not hold, then there are sequences  $\nu'_n \rightarrow 0$  as  $n \rightarrow \infty$  and  $\ell_n \leq 1$ , constants  $r_n \in (0, \ell_n]$ , and examples  $X_n = B(x_n, 1)$  boundary  $\nu'_n$ -flat near  $x_n$  on scale  $\ell_n$  with  $d(x_n, \partial X_n) \leq r_n \nu'_n$  yet  $X_n$  is not boundary  $\mu$ -flat near  $x_n$  on scale  $r_n$ . Passing to a subsequence we can suppose that the  $r_n/\ell_n \rightarrow d$  with  $0 \leq d \leq 1$ . Clearly, since the  $\nu'_n \rightarrow 0$ , the  $\ell_n^{-1} B(x_n, \ell_n)$  converge to a unit ball in half-space centered around a boundary point. If  $d > 0$ , then the  $(1/r_n) B(x_n, \ell_n)$  converge to a ball of radius  $d^{-1}$  in half-space centered about the boundary point and the result is established. On the other hand, if the  $d = 0$ , then, after passing to a subsequence, the sequence  $r_n^{-1} B(x_n, \ell_n)$

converges to a complete Alexandrov space  $(X, x)$  of curvature  $\geq 0$ . Since the  $\nu'_n$  go to zero, by the previous claim there is a geodesic line in  $X$  through  $x$ . On the other hand, since the distance from  $x_n$  to  $\partial X_n$  is at most  $r_n \nu'_n$  it follows that  $x \in \partial X$ . Hence,  $X$  is the product of  $\mathbb{R}$  with complete Alexandrov space  $Y$  of dimension 1 with a boundary. Clearly,  $Y$  is non-compact. Since  $\partial Y \neq \emptyset$ , it must be the case that  $Y = [0, \infty)$  and hence  $X$  is isometric to a closed half-space in  $\mathbb{R}^2$ . This proves that for all  $n$  sufficiently large,  $X_n$  is boundary flat at  $x_n$  on scale  $r_n$ . The result now follows.

The second statement is proved by a similar argument, using the first part for sequences for which  $d(y_n, \partial X_n)/\ell_n$  tends to zero.  $\square$

Putting this all together we obtain the analogue of Proposition 10.12 producing good annular regions. See FIG. 2.

**Proposition 10.18.** *Given  $\mu > 0$  and  $a' > 0$  there are positive constants  $s_2 = s_2(a')$  and  $s_1 = s_1(a')$  with  $0 < s_2 < s_1$  and  $\mu''_0(\mu, a') \leq \mu$  such that the following holds for all  $0 < \mu'' \leq \mu''_0(\mu, a')$ . Suppose that a 2-dimensional Alexandrov ball  $X = B(x, 1)$  of curvature  $\geq -1$  and area  $\geq a'$  is boundary  $\mu''$ -good near  $x$  on scale 1. Then:*

1. *For every point  $y \in \partial X \cap (B(x, 15/16) \setminus B(x, 1/100))$  the ball  $X$  is boundary  $\mu$ -flat near  $y$  on all scales  $\leq s_1$ .*
2. *For any  $z \in \text{int } X \cap (B(x, 7/8) \setminus B(x, 1/64))$ , the ball  $X$  is interior  $\mu$ -flat at  $z$  on all scales  $\leq \min(s_2, d(z, \partial X))$ .*
3. *For any  $b \in [1/64, 7/8]$  the metric sphere  $S(x, b)$  is an arc with endpoints in  $\partial X$  and the closed metric ball  $\overline{B(x, b)}$  is homeomorphic to a 2-disk.*

*Proof.* Fix  $\mu > 0$  and  $a' > 0$ . Let  $\nu > 0$  and  $\nu' > 0$  be the constants associated to  $\mu$  by Lemmas 10.11 and 10.16, respectively. We choose  $s_1$  so that the following holds. For any flat cone  $C$  in  $\mathbb{R}^2$  with cone point  $p$  and with the property that the area of  $B(p, 1)$  is at least  $a'$ , near every point of  $\partial(B(p, 15/16) \setminus B(p, 1/100))$  the cone  $C$  is boundary flat on scale  $s_1$ . Now we show that for  $\mu'' > 0$  sufficiently small and for every point  $y \in \partial X \cap (B(x, 15/16) \setminus B(x, 1/100))$  the ball  $X$  is boundary  $\nu'$ -flat near  $y$  on scale  $s_1$ . Suppose not. Then there are a sequence of  $\mu''_k \rightarrow 0$  and examples  $X_k = B(x_k, 1)$  of area  $\geq a'$  that are boundary  $\mu''_k$ -good near  $x_k$  on scale 1, for which there are points  $y_k \in \partial X_k \cap (B(x_k, 15/16) \setminus B(x_k, 1/100))$  near which  $X_k$  is not boundary  $\nu'$ -flat on scale  $s_1$ . Passing to a subsequence, we can suppose that the  $X_k$  converge to a limit which is a flat cone  $C$  of area  $\geq a'$  with cone point  $p$ . We can also assume that the  $y_k$  converge to  $\bar{y}$  in  $\overline{B(p, 15/16)} \setminus B(p, 1/100)$ , and by Lemma 10.2, we have  $\bar{y} \in \partial C$ . Thus,  $C$  is boundary flat near  $\bar{y}$  on scale  $s_1$ . Hence, for all  $k$  sufficiently large,  $X_k$  is boundary  $\nu'$ -flat near  $y_k$  on scale  $s_1$ , and hence by Lemma 10.16 boundary  $\mu$ -flat near  $y_k$  on all scales  $\leq s_1$ . This is a contradiction, proving the first statement

We fix  $s_2$  so that for any flat cone  $C$  in  $\mathbb{R}^2$  with cone point  $p$  and with the area of  $B(p, 1)$  being at least  $a'$ , any point  $y$  in the interior of the annular region  $B(p, 15/16) \setminus B(p, 1/100)$  of  $C$  has the property that  $C$  is interior flat at  $y$

on scale  $\min(s_2, d(y, \partial C))$ . We claim that provided that  $\mu''$  is sufficiently small then every point in the interior of  $B(x, 7/8) \setminus B(x, 1/64)$  is interior  $\nu$ -flat on scale  $\min(s_2, d(y, \partial X))$ . Again if it does not hold there is a sequence  $\mu_k'' > 0$  converging to 0 and  $X_k = B(x_k, 1)$  with points  $y_k \in \text{int } X_k \cap (B(x_k, 7/8) \setminus B(x_k, 1/64))$  at distance  $d_k$  from  $\partial X_k$  satisfying the hypothesis of the second statement for  $\mu_k''$  such that  $X_k$  is not interior  $\mu$ -flat near  $y_k$  of scale  $\min(s_2, d_k)$ . Since  $\mu_k'' \rightarrow 0$ , the  $X_k$  converge to a flat cone  $C$  in  $\mathbb{R}^2$  of area  $\geq a'$ . Passing to a subsequence we can assume that the  $d_k$  converge to  $d \geq 0$ . If  $d$  is positive, then passing to a further subsequence we can assume that the  $y_k$  converge to  $\bar{y} \in \text{int } C$  at distance  $d$  from  $\partial C$ . Since  $C$  is interior flat at  $y$  on all scales  $\leq \min(s_2, d)$ , for all  $k$  sufficiently large  $X_k$  is interior  $\nu$ -flat at  $y_k$  on scale  $\min(s_2, d_k)$ . Hence, by Lemma 10.11,  $X_k$  is interior  $\mu$ -flat at  $y_k$  on all scales  $\leq \min(s_2, d_k)$ . Suppose now that  $d = 0$ . For each  $k$  let  $z_k \in \partial X_k$  be a closest point to  $y_k$  on  $\partial X_k$ . Of course, for all  $k$  sufficiently large, we have  $1/100 < d(x_k, z_k) < 15/16$ . By the first part of this result, for every  $\nu > 0$ , for all  $k$  sufficiently large,  $X_k$  is boundary  $\nu$ -flat near  $n_k$  on all scales  $\leq 2d_k$ . It follows that the  $(1/2d_k)B(z_k, 2d_k)$  converge to the unit ball in half-space  $B(\bar{z}, 1)$  centered about a boundary point. Passing to a subsequence we arrange that the points  $y_k$  converge to a point  $\bar{y}$  at distance  $1/2$  from  $\bar{z}$  and also at distance  $1/2$  from  $\partial B(\bar{z}, 1)$ . This means that  $B(\bar{z}, 1)$  is interior flat at  $\bar{y}$  on all scales  $\leq 1/2$ . It then follows that for all  $k$  sufficiently large  $(1/2d_k)B(z_k, 2d_k)$  is interior  $\nu$ -flat on scale  $1/2$  at  $y_k$  and hence by Lemma 10.11, interior  $\mu$ -flat at  $y_k$  on all scales  $\leq 1/2$ . Hence, for all  $k$  sufficiently large,  $B(x_k, 1)$  is interior  $\mu$ -flat at  $y_k$  on all scales  $\leq d_k$ . This is a contradiction, establishing the second item.

The last statement is contained in Lemma 10.15.  $\square$

**Definition 10.19.** A 2-dimensional ball  $B(x, 1)$  satisfying the conclusion of the previous proposition is said to have a  $(\mu, s_1, s_2)$ -good collar. Notice that if  $s'_1 < s_1$  and  $s'_2 < s_2$  the a  $(\mu, s_1, s_2)$ -good collar is automatically a  $(\mu, s'_1, s'_2)$ -collar.

#### 10.4 The covering

In the previous subsection we studied balls that are interior good and boundary good on various scales. Now we show that there is a covering of any ball whose area is bounded below by a positive constant by such balls where the scales are uniformly bounded.

Recall from Lemma 10.8 that given any 2-dimensional Alexandrov ball  $B(\bar{x}, 1)$  of curvature  $\geq -1$  and area  $\geq a$ , for any  $\bar{y} \in B(\bar{x}, 15/16)$  and any  $r \leq 1/16$  the area of  $B(\bar{y}, r)$  is at least  $a'(a)r^2$ .

**Lemma 10.20.** *Given positive constants  $a, \mu$ , and  $r_0$  there is a positive constant  $r_1 = r_1(a, \mu, r_0) < r_0$  such that for any 2-dimensional Alexandrov ball  $B = B(x, 1)$  of curvature  $\geq -1$  and of area  $\geq a$  and any  $y \in \partial B \cap B(x, 15/16)$ , the ball  $B$  is boundary  $\mu_0'' = \mu_0''(\mu, a'(a))$ -good near  $y$  on some scale  $r(y)$  satisfying  $r_1 \leq r(y) \leq r_0$ .*

*Proof.* Fix positive constants  $a, \mu$ , and  $r_0$ , and suppose that there is no  $r_1 > 0$  as required. Then there is a sequence  $r_{1,n} \rightarrow 0$  as  $n \rightarrow \infty$  and Alexandrov balls



$B_n = B(x_n, 1)$  of curvature  $\geq -1$  and of area  $\geq a$  with  $y_n \in \partial B_n \cap B(x_n, 15/16)$  with the property that  $B_n$  is not boundary  $\mu''_0$ -good at  $y_n$  on any scale between  $r_{1,n}$  and  $r_0$ . Passing to a subsequence we can assume that the  $(B(x_n, 1), y_n)$  converge to  $(B(\bar{x}, 1), \bar{y})$  with  $\bar{y} \in \partial B(\bar{x}, 1)$ . The Alexandrov ball  $B(\bar{x}, 1)$  is of curvature  $\geq -1$  and has area  $\geq a$ . Now for any sequence  $\lambda_n \rightarrow \infty$  the Alexandrov balls  $\lambda_n(B(\bar{x}, 1), \bar{y})$  converge to the tangent cone of  $B(\bar{x}, 1)$  at  $\bar{y}$ . Since  $y \in \partial B(\bar{x}, 1)$ , this tangent cone is a flat cone in  $\mathbb{R}^2$ . It follows that there is  $0 < r(y) < r_0$  such that  $B(\bar{x}, 1)$  is boundary  $\mu''_0$ -good near  $y$  on scale  $r(y)$ . Thus, for all  $n$  sufficiently large  $B_n$  is boundary  $\mu$ -good at  $y_n$  on scale  $r(y)$ . Since  $r_{1,n} < r(y) < r_0$  for all  $n$  sufficiently large, this is a contradiction.  $\square$

**Corollary 10.21.** *Given positive constants  $a, \mu$ , and  $r_0$  there is a positive constant  $0 < r_1 = r_1(a, \mu, r_0) < r_0$  and positive constants  $\delta_0(a, \mu) > 0$ , and  $c(a)$  such that setting  $s_1 = s_1(a'(a))$  and  $s_2 = s_2(a'(a))$ , for any 2-dimensional Alexandrov ball  $B = B(x, 1)$  of curvature  $\geq -1$  and of area  $\geq a$  and for any  $y \in \partial B \cap B(x, 15/16)$ , either:*

1.  *$B$  is boundary  $\mu$ -good near  $y$  on scale  $r(y)$  and of angle  $\theta$ , where  $c(a) \leq \theta \leq \pi - \delta_0$  and where  $r_1 \leq r(y) \leq r_0$ , and furthermore  $(1/r(y))B(y, r(y))$  has a  $(\mu, s_1, s_2)$ -good collar, or*
2.  *$B$  is boundary  $\mu$ -flat near  $y$  on all scales  $\leq r_1$ .*

*Proof.* Given  $\mu > 0$  fix  $0 < \nu' \leq \mu$  as in Lemma 10.16. Then choose  $\delta_0 > 0$  such that any flat unit cone in  $\mathbb{R}^2$  of cone angle between  $\pi - \delta_0$  and  $\pi$  is within  $\nu'/2$  in the Gromov-Hausdorff distance of the flat unit cone of cone angle  $\pi$ . Then chose  $r_1 = r_1(a, \nu'/2, r_0)$ . From the previous result for any  $y \in \partial B \cap B(x, 15/16)$  there is  $r(y)$  with  $r_1 \leq r(y) \leq r_0$  such that the ball  $B$  is boundary  $\mu''_0(\nu'/2, a'(a))$ -good near  $y$  on some scale  $r(y)$ . Suppose the angle of the comparison cone is  $\leq \pi - \delta_0$ . Then, since the area of  $(1/r(y))B(y, r(y))$  is at least  $a'(a)$ , we see from Proposition 10.18 that  $(1/r(y))B(y, r(y))$  has a  $(\nu'/2, s_1, s_2)$ -good collar. Since  $\nu' \leq \mu$ , we see that  $(1/r(y))(y, r(y))$  has a  $(\mu, s_1, s_2)$ -good collar. Since  $\nu/2 < \mu$ , this completes the proof that Case 1 holds when the comparison angle is less than  $\pi - \delta_0$ , except for the uniform positive lower bound on the angle. The lower bound on the cone angle is immediate from the lower bound  $a'(a)$  on the area of the rescaled balls. This completes the proof when the cone angle is less than  $\pi - \delta_0$ .

Now suppose the cone angle is  $\geq \pi - \delta_0$ . Since  $\mu''_0 \leq \nu'/2$ , this implies that  $B$  is boundary  $\nu'/2$ -good at  $y$ . But in this case by the choice of  $\delta_0$ , the cone is within  $\nu'/2$  of the flat cone of cone angle  $\pi$ , and hence  $B$  is boundary  $\nu'$ -flat near  $y$  at scale  $r(y)$ . It then follows from Lemma 10.16 that  $B$  is boundary  $\mu$ -flat near  $y$  on all scales  $\leq r(y)$ . Since  $r_1 \leq r(y)$  this establishes the result in this case as well.  $\square$

**Proposition 10.22.** *Given positive constants  $a, \mu$ , and  $r_0$ , let  $s_1 = s_1(a'(a))$ ,  $s_2 = s_2(a'(a))$ ,  $\delta_0 = \delta_0(a, \mu)$ , and  $r_1 = r_1(a, \mu, r_0)$  be as in the previous lemma. Set  $s_0 = s_0(a'(a))$  from Proposition 10.12. Then for any  $d > 0$ , there is a positive constant  $r_2 = r_2(a, \mu, r_0, r_1, d) < r_1$  such that for any 2-dimensional Alexandrov ball*

$B(\bar{x}, 1)$  of curvature  $\geq -1$  and area  $\geq a$  and for any  $y \in \partial B \cap B(x, 15/16)$  one of the following holds:

1.  $B$  is boundary  $\mu$ -good near  $y$  of angle  $\leq \pi - \delta_0$  on some scale  $r(y)$  with  $r_1 \leq r(y) \leq r_0$ . Furthermore,  $(1/r(y))B(y, r(y))$  has a  $(\mu, s_1, s_2)$ -good collar region.
2.  $B$  is boundary  $\mu$ -flat near  $y$  at all scales  $\leq r_1$ .

If  $z \in B(x, 7/8)$  and  $d(z, \partial B) \geq d$ , then one of the following holds:

3.  $B$  is interior  $\mu$ -good at  $z$  of angle  $\leq 2\pi - \delta_0$  on some scale  $r(z)$  with  $r_2 \leq r(z) \leq r_1$  and  $(1/r(z))B(z, r(z))$  has a  $(\mu, s_0)$ -good annular region.
4.  $B$  is interior  $\mu$ -flat at  $z$  on all scales  $\leq r_2$ .

*Proof.* According Corollary 10.21 one of the first two possibilities holds for every  $y \in \partial B \cap B(x, 15/16)$ .

Now let  $\nu$  be the constant of Lemma 10.11 for  $\mu$  and let  $\mu_0$  be the minimum of  $\nu/2$  and  $\mu'(\mu, a'(a))$  as in Proposition 10.12, and let  $\delta' > 0$  be such that the flat circular cone of angle  $2\pi - \delta'$  is within  $\nu/2$  of the flat circular cone. We replace  $\delta_0$  by the minimum of  $\delta_0$  and  $\delta'$ . Fix  $d > 0$  and suppose that the result does not hold for these values of  $d, r_1$ , and  $\delta_0$  for any  $r_2$  with  $0 < r_2 < r_1$ . Then there are Alexandrov balls  $B_n = B(x_n, 1)$  of curvature  $\geq -1$  and area  $\geq a$  and points  $z_n \in B(x_n, 7/8)$  at distance at least  $d$  from  $\partial B_n$  for which the result does not hold for a constant  $r_{2,n}$  where  $r_{2,n} \rightarrow 0$  as  $n \rightarrow \infty$ . Taking a subsequence we can arrange that there is a Gromov-Hausdorff limit  $(B(x, 1), z)$  with  $z \in \overline{B(x, 7/8)}$ . By Corollary 10.6 we know that  $d(z, \partial B(x, 1))$  is at least  $d$ . Hence, for any sequence  $\lambda_n \rightarrow \infty$  the balls  $\lambda_n(B(x, 1), z)$  converge to the tangent cone of  $B(x, 1)$  at  $z$ . Thus, there is  $r(z)$  with  $0 < r(z) \leq r_1$  such that  $B(x, 1)$  is interior  $\mu_0$ -good at  $z$  on scale  $r(z)$ . Since  $\mu_0 \leq \mu'(\mu, a'(a))$ , it follows that if the angle of the tangent cone is  $< 2\pi - \delta_0$  then, by Proposition 10.12, the ball  $(1/r(z))B(z, r(z))$  has a  $(\mu, s_0)$ -good annular region. Thus, under this assumption on the limiting cone angle, Case 3 holds for all  $n$  sufficiently large, and we have a contradiction. If the limiting cone angle is  $\geq 2\pi - \delta$ , it follows from the choice of  $\delta_0$  and the fact that  $\mu_0 \leq \nu/2$  that  $B$  is interior  $\nu$ -flat at  $z$  on scale  $r(z)$  and hence interior  $\mu$ -flat, at all scales  $\leq r(z)$ . This implies that for all  $n$  sufficiently large  $B(x_n, 1)$  is interior  $\mu$ -flat at  $z_n$  on all scales  $\leq r(z)$ , which implies that Case 4 holds for all  $n$  sufficiently large. Once again that is a contradiction. This completes the proof of the result.  $\square$

#### 10.4.1 Geodesics approximating the boundary

Proposition 10.22 refers to points in the boundary and points whose distance from the boundary is at least  $d$ . To understand the points not covered by this result, it turns out that near the flat part of the boundary it is better to take neighborhoods centered around geodesics near the boundary rather than balls centered around boundary points. Here, we follow [33] closely.

**Definition 10.23.** Fix a 2-dimensional Alexandrov ball  $X$  with curvature  $\geq -1$ . Suppose that  $\gamma$  is an oriented geodesic in  $X$  with initial point  $e_-$  and final point  $e_+$  and of length  $\ell = \ell(\gamma)$ . We define

$$f_\gamma = \frac{1}{2}(d(e_-, \cdot) - d(e_+, \cdot)) \quad \text{and} \quad h_\gamma = d(\gamma, \cdot).$$

These are 1-Lipschitz functions. Further, for any  $\xi > 0$  we define

$$\nu_\xi(\gamma) = f_\gamma^{-1}([- \ell/4, \ell/4]) \cap h_\gamma^{-1}([0, \xi\ell]),$$

and

$$\bar{\nu}_\xi(\gamma) = f_\gamma^{-1}([- \ell/4, \ell/4]) \cap h_\gamma^{-1}([0, \xi\ell]).$$

We denote by  $\nu_\xi^0(\gamma) = \nu_\xi(\gamma) \setminus \overline{\nu_{\xi^2}(\gamma)}$ . The *ends* of  $\nu_\xi(\gamma)$  are their intersections with  $f_\gamma^{-1}(\pm\ell/4)$ , and the *side* of  $\overline{\nu_\xi(\gamma)}$  is its intersection with  $h_\gamma^{-1}(\xi\ell)$ . For any  $-\ell/4 \leq a < b \leq \ell/4$  we set

$$\nu_{\xi,[a,b]}(\gamma) = f_\gamma^{-1}([a, b]) \cap h_\gamma^{-1}([0, \xi\ell])$$

and we denote by  $\bar{\nu}_{\xi,[a,b]}(\gamma)$  its closure. The boundary of  $\bar{\nu}_{\xi,[a,b]}(\gamma)$  is made up of the side, given by  $h_\gamma^{-1}(\xi\ell)$ , and the two ends, given by  $f_\gamma^{-1}(a)$  and  $f_\gamma^{-1}(b)$ . We say that  $\xi\ell$  is the *width* of the neighborhood and  $(b - a)\ell$  is its *length*. The level set  $f_\gamma^{-1}(0)$  is the *center line* of  $\nu_\xi(\gamma)$ . See FIG. 3.

**Lemma 10.24.** Fix  $\xi > 0$  sufficiently small. Then there is  $0 < \alpha_0 = \alpha_0(\xi) \leq 10^{-3}$  such that for all  $\mu > 0$  sufficiently small the following hold. Suppose that  $X = B(x, 1)$  is a 2-dimensional Alexandrov ball of curvature  $\geq -\alpha_0^2$  with  $X$  being boundary  $\mu$ -flat near  $x$  on all scales  $\leq 1$ . Suppose that  $\gamma$  is a geodesic of length  $\ell \geq 1/100$  with endpoints  $e_-, e_+$  in  $\partial X \cap B(x, 15/16)$ . Then the following hold:

1. There is an arc  $A$  in  $B(x, 15/16)$  with endpoints  $e_\pm$ . The arc  $A$  and  $\gamma$  are within  $\xi^2\ell/100$  of each other in the Hausdorff distance in  $X$ .
2. For each  $y \in \bar{\nu}_\xi(\gamma)$  the comparison angle  $\tilde{\angle} e_- y e_+$  is greater than  $\pi - 6\xi$ .
3. For each point  $y \in \nu_\xi(\gamma) \setminus \nu_{\xi^2}(\gamma)$  there are points  $z, w \in B(x, 1)$  at distance at least  $1/8$  from  $y$  such that for any geodesic  $\mu$  from  $\gamma$  to  $y$ , denoting the point  $\mu \cap \gamma$  by  $a$ , we have

$$(a) \quad \tilde{\angle} a y z > \pi/2 - \xi^2,$$

$$(b) \quad \tilde{\angle} z y w > \pi/2 - \xi^2,$$

$$(c) \quad \tilde{\angle} a y w > \pi - \xi^2,$$

$$(d) \quad \tilde{\angle} e_- y z > \pi - 5\xi.$$

4. For any level set  $L$  of  $f_\gamma$  in  $\bar{\nu}_\xi(\gamma)$  and for any  $c \in [\xi^2, \xi]$  the distance from  $L \cap \gamma$  to any point of  $L \cap h_\gamma^{-1}([0, c\ell])$  is less than  $(1 + 2\xi)c\ell$ .

*Proof.* Direct computation shows that the result holds for  $\xi > 0$  sufficiently small for  $X$  being a ball of radius 1 in  $\mathbb{R} \times [0, \infty)$  centered about a boundary point when the comparison angles are measured in curvature 0. By taking  $\alpha_0 > 0$  sufficiently small we can arrange that if a triangle of side lengths  $\leq 2$  has Euclidean comparison angle  $\beta \geq \pi/4$  then the ratio of the comparison angle in curvature  $-\alpha_0^2$  to the flat comparison angle is arbitrarily close to 1. The result is then immediate by fixing  $\xi$  and taking limits as  $\alpha_0$  and  $\mu$  tend to zero.  $\square$

**Definition 10.25.** Fix  $\xi > 0$  sufficiently small so that the previous lemma holds. Let  $X = B(x, 1)$  be a 2-dimensional Alexandrov ball with curvature  $\geq -1$  and let  $s$  be given with  $0 < s \leq \alpha_0(\xi)$ . Suppose that we have a geodesic  $\gamma$  with endpoints in  $\partial X$ . If  $\nu_\xi(\gamma)$  satisfies the six conclusions in Lemma 10.24, then we say that  $\nu_\xi(\gamma)$  is a  $\xi$ -box and we call  $\nu_{\xi^2}(\gamma)$  the *core* of the  $\xi$ -box. For any  $\mu > 0$  we say that a geodesic  $\gamma \subset X$  is a  $\mu$ -approximation to  $\partial X$  on scale  $s$  if  $\gamma$  is a geodesic of length at least  $s/100$  and if there is a point  $y \in B(x, 15/16)$  near which  $X$  is boundary  $\mu$ -flat on scales  $\leq s$  with  $\gamma \subset B(y, s/3) \subset B(x, 15/16)$  and with the endpoints of  $\gamma$  contained in  $\partial X$ . The point  $y$  is a *control point* for  $\gamma$ .

Notice that if  $\mu$  is less than a positive constant depending only on  $\xi$  then for any  $\mu$ -approximation  $\gamma$  to the boundary of an Alexandrov ball of curvature  $\geq -1$  on scale  $s \leq \alpha_0(\xi)$  the regions  $\nu_\xi(\gamma)$  is  $\xi$ -box. The point is that  $s^{-1}B(y, s)$  is an Alexandrov ball of curvature  $\geq -\alpha_0^2(\xi)$ .

#### 10.4.2 Intersections of $\xi$ -boxes

We need to know how two approximations to the boundary in a single ball are related, see FIG. 4.

**Lemma 10.26.** *Given  $\xi > 0$  the following holds for all  $0 < \mu$  sufficiently small. Suppose that  $B(\bar{x}, 1)$  is a 2-dimensional Alexandrov ball that is boundary  $\mu$ -flat at  $\bar{x}$  on all scales  $\leq 1$  and that  $\bar{\gamma}_1, \bar{\gamma}_2$  are geodesics in  $B(\bar{x}, 7/8)$  with lengths  $\ell(\bar{\gamma}_i)$  between (0.24) and (0.26) and each geodesic with endpoints in  $\partial B(\bar{x}, 1)$ . Suppose that there are points in  $x_1 \in \nu(\bar{\gamma}_1)$  and  $x_2 \in \nu(\bar{\gamma}_2)$  with  $d(x_1, x_2) < (0.01)$ . Then there are arcs  $A \subset \partial B(\bar{x}, 1)$ ,  $\alpha_1 \subset \bar{\gamma}_1$  and  $\alpha_2 \subset \bar{\gamma}_2$ , with the following properties*

1. *The endpoints of  $A$  are the two middle endpoints (measured along  $\partial B(\bar{x}, 1)$ ) of the union of the endpoints of  $\bar{\gamma}_1$  and those of  $\bar{\gamma}_2$ . If the both endpoints of  $A$  are endpoints of  $\bar{\gamma}_1$  then  $\alpha_1 = \bar{\gamma}_1$ ; similarly if the two endpoints of  $A$  are those of  $\bar{\gamma}_2$ . Otherwise, each  $\alpha_i$  shares exactly one endpoint with  $\bar{\gamma}_i$ .*
2. *For  $i = 1, 2$  we have  $d(A, \alpha_i) < \xi^2/100$  and  $d(\alpha_1, \alpha_2) < \xi^2/100$ .*

*Proof.* Suppose that the result does not hold for some  $\xi > 0$ . Then there is a sequence  $\mu_n \rightarrow 0$  and for each  $n$  a counter example for  $\mu_n$  consisting of  $B(\bar{x}_n, 1)$  and geodesics  $\bar{\gamma}_{n,1}, \bar{\gamma}_{n,2} \subset B(\bar{x}_n, 7/8)$ . Passing to a subsequence we can assume that the  $B(\bar{x}_n, 1)$  converge to  $B(\bar{x}_\infty, 1)$  and the  $\bar{\gamma}_{n,i}$  converge to  $\bar{\gamma}_{\infty,i}$  with endpoints in  $\partial B(\bar{x}_\infty, 1)$ . Since the  $\mu_n \rightarrow 0$ , it follows that  $B(\bar{x}_\infty, 1)$  is a sub-ball of half-space, and

$\bar{\gamma}_{\infty,i}$  are sub-geodesics of the boundary. We set  $A_\infty = \alpha_{i,\infty} = \bar{\gamma}_{\infty,1} \cap \bar{\gamma}_{\infty,2}$ . This arc is the limit of the arcs  $A_n$  on the boundary between the middle (as measured along  $\partial B(\bar{x}_n, 1)$ ) two endpoints. Clearly, there are arcs  $\alpha_{n,i} \subset \gamma_{n,i}$  sharing endpoints with the  $\bar{\gamma}_{n,i}$  as indicated, converging to  $A_\infty$ . Thus, the conclusion of the lemma holds for all  $n$  sufficiently large, which is a contradiction and establishes the lemma.  $\square$

The same argument as in the previous proof can be used to show the following result which allows us to compare the way that neighborhoods around two geodesic approximations to the boundary meet, see FIG. 4.

**Corollary 10.27.** *For all  $\xi > 0$  sufficiently small, the following hold for all  $\mu > 0$  sufficiently small. Let  $X = B(x, 1)$  be a 2-dimensional Alexandrov ball of curvature  $\geq -1$ . Suppose that  $X$  is boundary  $\mu$ -flat near  $x$  on all scales  $\leq 1$ . Suppose that we have geodesics  $\gamma_1$  and  $\gamma_2$  as in the previous lemma. Fix a direction along  $\partial X \cap B(x, 15/16)$  and let endpoints of  $\gamma_i$ , denoted  $e_\pm(\gamma_i)$ , be chosen so that in the given direction along  $\partial X$  we have  $e_-(\gamma_i) < e_+(\gamma_i)$  for  $i = 1, 2$ . Then the following hold:*

1. *For any point  $y \in \bar{\nu}_\xi(\gamma_1) \cap \bar{\nu}_\xi(\gamma_2)$ , the comparison angles satisfy:*

$$\tilde{Z}e_-(\gamma_1)ye_+(\gamma_2) > \pi - 10\xi$$

and

$$\tilde{Z}e_-(\gamma_2)ye_+(\gamma_1) > \pi - 10\xi.$$

2. *Suppose that a level set  $L \subset \bar{\nu}_\xi(\gamma_2)$  for  $f_{\gamma_2}$  meets  $\nu_\xi(\gamma_1)$ . Then for any  $y_1, y_2 \in L \cap \nu_\xi(\gamma_1)$  we have*

$$|f_{\gamma_1}(y_1) - f_{\gamma_2}(y_2)| < \xi^2 \ell(\gamma_1).$$

#### 10.4.3 Intersection of $\xi$ -boxes and boundary $\mu$ -good balls

We must also compare flat regions near the boundary with balls around boundary points, see FIG. 5.

**Lemma 10.28.** *Given  $\xi > 0$  sufficiently small, the following hold for all  $0 < \mu$  sufficiently small and given  $a' > 0$ , with  $s_1 = s_1(a')$  as in Proposition 10.18. Suppose that  $X = B(x, 1)$  is a 2-dimensional Alexandrov ball of curvature  $\geq -1$  and area  $\geq a'$  that is boundary  $\mu''_0(\mu, a')$ -good near  $x$  on scale 1. Suppose that  $\gamma \subset X$  is a geodesic of length at most  $s_1/2$  contained in  $A = B(x, 15/16) \setminus B(x, 1/64)$  with endpoints  $e_\pm$  in the same component of  $(B(x, 15/16) \setminus B(x, 1/64)) \cap \partial X$ . We orient  $\gamma$  so that  $e_-$  separates  $e_+$  from  $\partial B(x, 1/16)$  along  $\partial X$ . Then:*

1. *For any  $y \in \nu_\xi(\gamma)$  the comparison angle  $\tilde{Z}xye_+$  is greater than  $\pi - \xi$ .*
2. *For any level set  $L$  of  $d(x, \cdot)$  that meets  $\bar{\nu}_{\xi, [-(.24)\ell, (.24)\ell]}$ , the intersection  $L \cap \bar{\nu}_\xi(\gamma)$  is an interval with one endpoint in  $\partial X$  and the other in the side of  $\bar{\nu}_\xi(\gamma)$ .*
3. *The function  $f_\gamma$  varies by at most  $8\xi\ell(\gamma)$  on  $L \cap \bar{\nu}_\xi(\gamma)$ .*

*Proof.* Let  $\gamma$  be a geodesic in  $A = B(x, 15/16) \setminus B(x, 1/64)$  with endpoints  $z, z'$  in the same component of  $\partial A$ , with  $z'$  farther from  $x$  than  $z$ , and let  $y$  be a point in the interior of  $\gamma$ . Choose a point  $w \in \partial X$  be a point at distance  $1/16$  from  $z'$  and farther from  $x$  than  $z'$ . Then, given  $\xi$ , for all  $\mu$  sufficiently small we have  $\tilde{\angle}xyz' > \pi - \xi$ . This is clear by taking limits as  $\mu \rightarrow 0$  since any such limit is a flat cone in  $\mathbb{R}^2$ . Fix geodesics  $\beta_1$  from  $y$  to  $w$  and  $\beta_2$  from  $x$  to  $z'$ . Then  $\beta_1 \cap \beta_2$  is a point  $u$ . We have

$$\tilde{\angle}xyw \leq \tilde{\angle}xyu \leq \tilde{\angle}xyu + \tilde{\angle}uyz' \leq \tilde{\angle}xyz',$$

showing that for  $\mu$  sufficiently small we have  $\tilde{\angle}xyz' > \pi - \xi$ .

Since any level set of  $f_\gamma$  contained in  $\nu_\xi(\gamma)$  has diameter less than  $2\xi\ell$ , we see that  $d(x, \cdot)$  varies by at most  $2\xi\ell$  on any such level set. Hence, if  $L$  is a level set for  $f_\gamma$  contained in  $\nu_{\xi, [-0.24\ell, 0.24\ell]}(\gamma)$ , the values of the restriction of  $d(x, \cdot)$  to  $L$  lie strictly between the values of the restriction of  $d(x, \cdot)$  to either end of  $\nu_\xi(\gamma)$ . Since  $d(x, \cdot)$  is regular on  $A$ , it follows that the level sets of this function contained in  $A$  are intervals with end points in the boundary. The functions  $d(x, \cdot)$  and  $h_\gamma$  are Lipschitz coordinates on  $\nu_\xi^0(\gamma)$ , so that any level set of  $d(x, \cdot)$  that meets  $\nu_{\xi, [-0.24\ell, 0.24\ell]}(\gamma)$  crosses each level set of  $h_\gamma$  in  $\nu_\xi^0$  exactly once. It now follows that the intersection of any such level set of  $d(x, \cdot)$  with  $\nu_\xi(\gamma)$  is an interval with one endpoint in  $\partial A$  and the other in the side of  $\nu_\xi(\gamma)$ .

Lastly, let  $a, b \in \nu_{\xi, [-0.24\ell, 0.24\ell]}(\gamma)$  be two points in a level set for  $d(x, \cdot)$ . Let  $a', b' \in \gamma$  be points on the same level sets for  $f_\gamma$  as  $a$  and  $b$ , respectively. Then,  $|d(x, a) - d(x, a')|$  and  $|d(x, b) - d(x, b')|$  are both  $< 2\xi\ell$  so that  $|d(x, a') - d(x, b')| < 4\xi\ell$ . On the other hand, it follows from the comparison angle inequality that for points  $a, b \in \gamma \cap \nu_\xi(\gamma)$  we see that  $|d(x, a) - d(x, b)| > d(a, b)/2$ . This implies that the distance along  $\gamma$  from  $a'$  to  $b'$  is at most  $8\xi\ell$ , and hence that  $|f_\gamma(a') - f_\gamma(b')| < 8\xi\ell$ , completing the proof of the third statement.  $\square$

**Upper bound for  $\xi$ .** At this point we choose  $0 < \xi_0 \leq 10^{-6}$  sufficiently small such that the three previous results hold for all  $\xi < \xi_0$ . Then for any  $\xi < \xi_0$  we choose  $\alpha_0 = \alpha_0(\xi) \leq 10^{-3}$ . These values are fixed from now on.

Now we give an analogue of Proposition 10.18 using  $\epsilon$ -solid cylinder neighborhoods.

**Proposition 10.29.** *For any  $\xi$  with  $0 < \xi < \xi_0$ , let  $\mu > 0$  be sufficiently small and let  $a' > 0$  be a positive constant. For  $i = 1, 2$ , let  $s_i = s_i(a')$  be as in Proposition 10.18. Then the following holds for all  $\mu''$  less than  $\mu_0''(\mu, a')$ . Suppose that  $B = B(x, 1)$  is an Alexandrov ball of curvature  $\geq -1$  and area  $\geq a'$  that is boundary  $\mu''$ -good near  $x$  on scale 1. Then  $A = B(x, 7/8) \setminus B(x, 1/64)$  is contained in the union of the open set,  $U_0$ , of points at which  $B$  is interior  $\mu$ -flat on all scales  $\leq \min(s_2(a'), \xi^2 s_1/100)$  and the open set,  $U_1 \subset A$ , of points within  $\xi^2 s_1/100$  of  $\partial B(x, 1)$ . Furthermore, for any  $y \in \partial A$  the ball  $B(x, 1)$  is boundary  $\mu$ -flat near  $y$  on all scales  $\leq s_1$ .*

*Proof.* Given  $\xi, a'$  and  $\mu$  fix Let  $B = B(x, 1)$  be as in the statement of this proposition for some  $\mu'' \leq \mu_0''(\xi, \mu, a')$ , and denote  $B(x, 7/8) \setminus B(x, 1/64)$  by  $A(x)$ . Then

according to Proposition 10.18 the ball  $B$  is boundary  $\mu$ -flat on all scales  $\leq s_1$  near every  $y \in A(x) \cap \partial B$  and for every  $y \in A(x) \cap \text{int } B$  the ball  $B$  is interior  $\mu$ -flat at  $y$  on all scales  $\leq \min(\tilde{s}_2(a'), d(y))$  where  $d(y)$  is the distance from  $y$  to  $\partial B$ . Now we set  $d = \xi^2 s_1 / 100$ . Then every point of  $A$  either has the property that  $B$  is interior  $\mu$ -flat at this point on all scales  $\leq \min(s_2(a'), d)$  or it is within  $d$  of  $\partial B$ .  $\square$

Now we are ready to reformulate Proposition 10.22 using the approximations to the boundary.

**Theorem 10.30.** *For every  $0 < \xi < \xi_0$ , and fixing  $a > 0$ , then there is a positive constant  $\mu_1(a, \xi)$  such that for every  $0 < \mu \leq \mu_1(a, \xi)$  Lemma 10.24, Lemma 10.26, Corollary 10.27, Lemma 10.28, and Proposition 10.29 hold. Furthermore, setting  $r_0 = \min(\alpha_0(\xi), 10^{-6})$ , there are positive constants  $\delta_0, r_1, r_2, s_0, s_1, s_2$  depending on  $\xi, \mu$  and  $a$  with  $r_2 < r_1 < r_0$ , such that for any 2-dimensional Alexandrov ball  $B = B(x, 1)$  of curvature  $\geq -1$  and area  $\geq a$  and any  $y \in B(x, 7/8)$  one of the following two cases holds.*

1. *The distance from  $y$  to  $\partial B$  is at least  $\xi^2 r_1 s_1 / 100$  and one of the following two holds:*
  - (a)  *$B$  is interior  $\mu$ -good at  $y$  of angle  $\leq 2\pi - \delta_0$  on some scale  $r = r(y)$  with  $r_2 \leq r(y) \leq r_1$  and  $(1/r)B(y, r)$  has a  $(\mu, s_0)$ -good annular region.*
  - (b)  *$B$  is interior  $\mu$ -flat at  $y$  on all scales  $\leq r_2$ .*
2. *There is a point  $z \in \partial B$  with  $d(y, z) < \xi^2 r_1 s_1 / 100$  and one of the following two holds:*
  - (a)  *$B$  is boundary  $\mu$ -flat at  $z$  on all scales  $\leq r_1 s_1$ . In this case there is a  $\mu$ -approximation  $\gamma$  to the boundary on scale  $r_1 s_1$ , with the length of  $\gamma$  being  $r_1 s_1 / 4$ , such that  $y \in \nu_{\xi^2/2}(\gamma)$ . Furthermore, given any  $b$  with  $-r_1 s_1 / 16 \leq b \leq r_1 s_1 / 16$  we can choose  $\mu$ -approximation  $\gamma$  of length  $r_1 s_1 / 4$  so that  $f_\gamma(y) = b$ .*
  - (b) *The ball  $B$  is boundary  $\mu$ -good near  $z$  of angle  $\leq \pi - \delta_0$  on some scale  $r = r(z)$  with  $r_1 \leq r \leq r_0$  and  $(1/r)(B(z, r))$  has a  $(\mu, s_1, s_2)$ -good collar.*

*Proof.* Given  $0 < \xi < \xi_0$ ,  $a > 0$ , we fix  $0 < \mu$  sufficiently small so that Lemma 10.24, Lemma 10.26, Corollary 10.27, Lemma 10.28, and Proposition 10.29 hold. Now we set  $r_1, r_2, \delta_0$  equal to the constants by the same name in Proposition 10.22 for these values of  $a, \mu, r_0$ . Also, we take  $s_0, s_1$  as in that proposition. Next, we set  $d = \xi^2 r_1 s_1 / 100$ . Now let  $s_2$  be the minimum of  $d$  and  $s_2(a')$ . Fix  $y \in B(x, 7/8)$ . If  $d(y, \partial B) \geq d$ , then by Proposition 10.22, Case 1 of this result holds for  $y$ . If  $d(y, \partial B) < d$ , then let  $z \in \partial B$  be a point with  $d(y, z) < d$ . Then  $z \in B(x, 15/16)$ , and by Proposition 10.22 either Case 2(b) holds or  $B$  is boundary  $\mu$ -flat at  $z$  on all scales  $\leq r_1$  and *a fortiori* on all scales  $\leq r_1 s_1$ . Suppose that the latter holds. Orient  $\partial B$  near  $z$  and let  $\gamma_+$  and  $\gamma_-$  be geodesics of length  $r_1 s_1 / 4$  with endpoints in  $\partial B$ , consistently oriented, so that  $e_-(\gamma_+) = z = e_+(\gamma_-)$ . Then these geodesics are

contained in  $B(z, r_1 s_1/3)$  and hence are  $\mu$ -approximations to the boundary. Furthermore,  $f_{\gamma_+}(z) = r_1 s_1/4$  and  $f_{\gamma_-}(z) = -r_1 s_1/4$ . Hence,  $f_{\gamma_+}(y) > b$  and  $f_{\gamma_-}(y) < b$ . As we deform a geodesic  $\gamma$  keeping its length  $r_1 s_1/4$  and keeping its endpoints in  $\partial B$  from  $\gamma_+$  to  $\gamma_-$ , the geodesic remains in  $B(z, r_1 s_1/3)$  and consequently remains a  $\mu$ -approximation to the boundary. Also, the value of  $f_\gamma(y)$  varies continuously. Thus, one of the geodesics  $\gamma$  with these properties between  $\gamma_+$  and  $\gamma_-$  is such that  $f_\gamma(y) = b$ . Since  $d(y, z) < d$  and since the distance between  $\gamma$  and the arc of  $\partial B$  with the same endpoints is at most  $\xi^2 r_1 s_1/100$ , we see that  $h_\gamma(y) < \xi^2 r_1 s_1/50$ . It follows that  $y \in \nu_{\xi^2/2}(\gamma)$ , so that Case 2(a) holds for  $y$ .  $\square$

### 10.5 Transition between the 2- and 1-dimensional part

We need to understand the passage between the regions of  $M_n$  close to 1- and to 2-dimensional Alexandrov balls. A non-compact 1-dimensional Alexandrov ball  $B(x, 1)$  is either an open interval of length 2 or is a half-open interval of length  $\ell$  with  $1 \leq \ell \leq 2$ .

**Lemma 10.31.** *The following hold for all  $\beta > 0$  and for all  $a > 0$  less than a positive constant  $a_2(\beta)$ . Let  $B(x, 1)$  be a 2-dimensional Alexandrov ball of curvature  $\geq -1$  and suppose that there is a point  $y \in B(x, 24/25)$  with the area of  $B(y, 1/100)$  being at most  $a$ . Then  $B(x, 1)$  is within  $\beta$  in the Gromov-Hausdorff distance of 1-dimensional Alexandrov ball  $J$ .*

*Proof.* Fixing  $\beta > 0$  suppose that the result does not hold for any  $a > 0$ . Then there is a sequence  $a_k \rightarrow 0$  and a sequence  $B(x_k, 1)$  of 2-dimensional Alexandrov balls of curvature  $\geq -1$  and points  $y_k \in B(x_k, 24/25)$  with the area of  $B(y_k, 1/100)$  equal to  $a_k$  for which the result does not hold. Passing to a subsequence we can extract a limit  $\bar{B}$  with the  $y_k$  converging to  $\bar{y} \in \bar{B}$ . Because of the area condition, the neighborhood  $B(\bar{y}, 1/100)$  must be 1-dimensional, and hence  $\bar{B}$  is a 1-dimensional Alexandrov ball.  $\square$

If we choose  $\beta > 0$  sufficiently small, then it follows that  $d(x, \cdot)$  is regular on  $B(x, 7/8) \setminus B(x, 1/8)$  and for all  $t \in (1/8, 7/8)$  each connected component of the level set of  $\{y | d(x, y) = t\}$  is either a simple closed curve or a closed interval.

## 11 3-dimensional analogues

Now we discuss the structure of balls in a 3-dimensional Riemannian manifold that are close to the various 1- and 2-dimensional balls that we have been discussing. Since we shall need the results for 3-dimensional balls near 2-dimensional Alexandrov balls in our study of 3-dimensional balls near 1-dimensional balls, we start with the 2-dimensional case. Recall that for any  $x \in M_n$  we denote by  $g'_n(x)$  the rescaled metric  $\rho_n^{-2}(x)g_n$ . Throughout this section we consider  $B_{\lambda^2 g_n}(x, 1)$  where  $x \in M_n$  and  $\lambda \geq \rho_n(x)^{-1}$ . Of course, the sectional curvatures of these balls are bounded below by  $-1$ . Any time we refer to such  $B_{\lambda^2 g_n}(x, 1)$ , unless we explicitly state the contrary, we are implicitly assuming that it is disjoint from the boundary.



Let us describe the nature of regions in  $M_n$  near the four different types of regions in 2-dimensional Alexandrov balls that we listed in the last section. Here  $\epsilon > 0$ ,  $\delta > 0$ , and  $\mu > 0$  are fixed sufficiently small, and the statements below hold for all  $n$  sufficiently large.

1. If  $B_{\lambda^2 g_n}(x, 1)$  is close in the Gromov-Hausdorff sense to a 2-dimensional Alexandrov ball  $B(\bar{x}, 1)$  that is interior  $\mu$ -flat at  $\bar{x}$ , then there is a neighborhood of  $x$  in  $M_n$  on which the metric  $g_n$  is, after rescaling,  $C^N$ -close (for some sufficiently large  $N$ ) to a product of a circle of length 1 and a 2-dimensional Euclidean ball  $B(0, \epsilon^{-1})$ . These regions are called *almost  $S^1$ -product regions*.
2. If  $B_{\lambda^2 g_n}(x, 1)$  is close in the Gromov-Hausdorff sense to a 2-dimensional Alexandrov ball  $B(\bar{x}, 1)$  that is interior  $\mu$ -good at  $x$  of angle  $\leq 2\pi - \delta$  on scale  $r$ , then there is a neighborhood  $V$  containing  $B_{\lambda^2 g_n}(x, 3r/4)$  in  $M_n$  that is an open solid torus. Furthermore, the complement of a compact, unknotted sub-torus  $S$  of  $V$  is covered by almost  $S^1$ -product neighborhoods as in 1). The circle factors in these almost product regions are isotopic in  $V$  into  $\partial S$  and are homotopically non-trivial in  $V$ .
3. If  $B_{\lambda^2 g_n}(x, 1)$  is close in the Gromov-Hausdorff sense to a 2-dimensional Alexandrov ball  $B(\bar{x}, 1)$  that is boundary  $\mu$ -flat near  $\bar{x}$ , then there is a neighborhood of  $x$  in  $M_n$  that is diffeomorphic to a product  $\text{int } D^2 \times I$ . The complement of a compact subset of the form  $\overline{D}' \times I$ , for  $\overline{D}'$  a compact sub-disk of  $D^2$ , is covered by almost  $S^1$ -product neighborhoods of Type 1 above, and the circles in these product neighborhoods which are outside of  $\overline{D}' \times I$  are isotopic in  $D^2 \times I \setminus \overline{D}' \times I$  to the boundary of the  $D^2$ -factors.
4. If  $B_{\lambda^2 g_n}(x, 1)$  is close in the Gromov-Hausdorff sense to a 2-dimensional Alexandrov ball  $B(\bar{x}, 1)$  that is boundary  $\mu$ -good at  $\bar{x}$  on scale  $r$  of angle  $\leq \pi - \delta$  then  $B_{\lambda^2 g_n}(x, 3r/4)$  is diffeomorphic to a 3-ball. Furthermore, each metric sphere  $S_{\lambda^2 g_n}(x, t)$ , for  $1/4 \leq t \leq 7/8$  is contained in the union of two disjoint neighborhoods of type 3) and an open subset of points of type 1).

Refined versions of all these statements will be established in this section.

### 11.1 Generic interior points of 2-dimensional Alexandrov spaces

We begin with a description of the 3-dimensional part of a Riemannian 3-manifold  $M$  that is near the ‘generic’ part of a 2-dimensional Alexandrov ball of curvature  $\geq -1$ .

**Lemma 11.1.** *The following hold for all  $\epsilon > 0$ , for all  $\mu > 0$  less than a positive constant  $\mu_2(\epsilon)$ , and, given  $0 < s_0 \leq 1/2$ , for all  $\hat{\epsilon} > 0$  less than a positive constant  $\hat{\epsilon}_0(\epsilon, s_0)$ . Suppose that the ball  $B_{\lambda^2 g_n}(x, 1)$  is within  $\hat{\epsilon}$  of a 2-dimensional Alexandrov ball  $B = B(\bar{x}, 1)$  of curvature  $\geq -1$  that is interior  $\mu$ -flat at  $\bar{x}$  on all scales  $\leq s_0$ . Then there exist a smooth embedding  $\varphi: S^1 \times B(0, \epsilon^{-1}) \rightarrow M_n$  with  $x \in \varphi(S^1 \times \{0\})$*

and a constant  $\lambda' > \epsilon^{-1}\lambda$  such that the metric  $\varphi^*((\lambda')^2 g_n)$  is within  $\epsilon$  in the  $C^{[1/\epsilon]}$ -topology to the product of the metric of length 1 on the circle and the restriction of the standard Euclidean metric to  $B(0, \epsilon^{-1})$ .

*Proof.* Let us first show that it suffices to prove the first conclusion for  $s_0 = 1/2$ . For, suppose that we have established the conclusion in this special case with constants  $\mu_2(\epsilon)$  and  $\hat{\epsilon}_0(\epsilon, 1/2)$ , and let us consider the statement for another value  $0 < s_0 \leq 1/2$ . Suppose for some  $\mu < \mu_2(\epsilon)$  and  $\hat{\epsilon} < 2s_0\hat{\epsilon}_0(\epsilon, 1/2)$ , the ball  $B_{\lambda^2 g_n}(x, 1)$  is within  $\hat{\epsilon}$  of  $B(\bar{x}, 1)$ , the latter being interior  $\mu$ -flat  $\bar{x}$  on all scales  $\leq s_0$ . Then  $B_{(\lambda^2/4s_0^2)g_n}(x, 1)$  is within  $\hat{\epsilon}/(2s_0)$  of  $\frac{1}{2s_0}B(\bar{x}, 2s_0)$ , and the latter is  $\mu$ -flat at  $\bar{x}$  on all scales  $\leq 1/2$ . Since, by construction,  $\hat{\epsilon}/(2s_0) < \hat{\epsilon}_0(\epsilon, 1/2)$ , the result for  $s_0 = 1/2$  implies the existence of a constant  $(\lambda')^2$  as required. (Of course,  $(\lambda') > (\lambda/2s_0)$  since  $B_{(\lambda^2/4s_0^2)g_n}(x, 1)$  is close to a 2-dimensional ball whereas  $B_{(\lambda')^2 g_n}(x, 1)$  has 3-dimensional volume bounded away from zero.)

Thus, we can now assume that  $s_0 = 1/2$ . Fix  $\epsilon > 0$  and suppose that the first conclusion does not hold for this constant. Then there are sequences  $\mu_k \rightarrow 0$  and  $\hat{\epsilon}_k > 0$  both tending to zero as  $k \rightarrow \infty$  such that for each  $k$  there is an index  $n(k)$  and a point  $x_{n(k)} \in M_{n(k)}$  and constants  $\lambda_k \geq \rho_{n(k)}^{-1}(x_{n(k)})$  so that the ball  $B_{n(k)} = B_{\lambda_k^2 g_{n(k)}}(x_{n(k)}, 1)$  is within  $\hat{\epsilon}_k$  of a 2-dimensional Alexandrov ball  $B_k = B(\bar{x}_k, 1)$  that is interior  $\mu_k$ -flat at  $\bar{x}_k$  on all scales  $\leq 1/2$ , yet no  $x_{n(k)}$  satisfies the first conclusion of the lemma. The fact that the  $B_{\lambda_k^2 g_{n(k)}}(x_{n(k)}, 1)$  converge to a 2-dimensional ball implies that the volumes  $v_k$  of these balls go to zero.

Since  $\mu_k \rightarrow 0$  and  $\hat{\epsilon}_k \rightarrow 0$ , it follows from Lemma 10.10 that for each  $\delta > 0$  there is  $s > 0$  such that for all  $k$  sufficiently large  $B_{n(k)}$  has a  $(2, \delta)$ -strainer of size  $s$ . Now we let  $\omega$  be the volume of the unit ball in  $\mathbb{R}^3$  and we rescale  $B_{n(k)}$  by a constant  $\alpha_k$  such that the volume of the unit ball about  $x_{n(k)}$  in the rescaled ball is  $\omega/2$ . This is possible since  $B_{n(k)}$  is a Riemannian 3-manifold and since the volumes of the  $B_{n(k)}$  tend to zero. It follows from the latter fact and Bishop-Gromov comparison that the  $\alpha_k \rightarrow \infty$ . Hence, for every  $R < \infty$  and every  $\delta > 0$ , for all  $k$  sufficiently large, there is a  $(2, \delta)$ -strainer of size  $R$  centered at  $x_{n(k)}$  in  $\alpha_k B_{n(k)}$ . After passing to a subsequence there is a limit,  $(X, x)$ , of the  $\alpha_k B_{n(k)}$ . Since we arranged that the volumes of the unit balls in the sequence are constant, by Proposition 9.46 the limit  $X$  is a smooth, complete, non-compact manifold of non-negative curvature and without boundary, and (after passing to a further subsequence) the convergence is a smooth. The existence of the  $(2, \delta_k)$ -strainers of size going to infinity in the sequence implies that there is an isometric copy of  $\mathbb{R}^2$  in  $X$  through  $x$ . Hence, by Corollary 9.17,  $X$  splits as a product of  $\mathbb{R}^2$  with a complete, connected 1-manifold without boundary. This 1-manifold cannot be  $\mathbb{R}^1$  because the volume of the unit ball in  $X$  is one-half the volume of the unit ball in Euclidean space. Thus,  $X$  is the product of a circle with  $\mathbb{R}^2$ . Rescaling again by a fixed constant, we can make the limit the product of the circle of length 1 with  $\mathbb{R}^2$ . The conclusion of the lemma then holds for all  $k$  sufficiently large by taking limits. This is a contradiction and proves the existence of the map  $\varphi$  as required.

Now let us compare  $\lambda'$  and  $\lambda$ . Under  $(\lambda')^2 g_n$  the volume of the  $S^1$ -product neighborhood is at least  $\pi\epsilon^{-2}/2$  whereas its volume under  $\lambda^2 g_n$  goes to zero with  $\hat{\epsilon}$ .

Thus,  $\lambda'/\lambda \rightarrow \infty$  as  $\hat{\epsilon} \rightarrow 0$ . □

**Definition 11.2.** Any time we have an embedding  $\varphi: S^1 \times B(0, \epsilon^{-1}) \rightarrow M$  with  $x \in \varphi(S^1 \times \{0\})$  that satisfies the conclusion of the previous lemma, we say that the image of  $\varphi$  is an  $S^1$ -product neighborhood with  $\epsilon$ -control. The point  $x$  is said to be the center of the neighborhood, and the neighborhood is said to be centered at  $x$ . The horizontal spaces of an  $S^1$ -product neighborhood are the subspaces  $\varphi(\{\theta\} \times B(0, \epsilon^{-1}))$  for  $\theta \in S^1$ .

We need a semi-local version of this result. First a definition.

**Definition 11.3.** Recall that given a point  $y$  in an Alexandrov space  $B$  and given a compact subset  $A$  of  $B$  disjoint from  $y$  we denote by  $A' \subset S_y$  the compact subset of the tangent sphere of  $B$  at  $y$  consisting of the tangent directions at  $y$  to all minimal length geodesics from  $y$  to  $A$ . Given four compact sets  $A_1, A_2, B_1, B_2$  disjoint from  $y$  we say that  $\{A'_1, A'_2, B'_1, B'_2\} \subset S_y$  form a  $(2, \delta)$ -strainer if the following hold:

1.  $d(A'_i, B'_i) > \pi - \delta$  for  $i = 1, 2$ ,
2.  $d(A'_1, A'_2) > \pi/2 - \delta$ ,
3.  $d(B'_1, B'_2) > \pi/2 - \delta$ , and
4.  $d(A'_i, B'_j) > \pi/2 - \delta$  for all  $i \neq j$ ,

where  $d$  denotes the distance function on  $S_y$ .

**Proposition 11.4.** For every  $\epsilon' > 0$  sufficiently small there is a positive constant  $\epsilon_0(\epsilon')$  such that for all  $0 < \epsilon \leq \epsilon_0(\epsilon')$  the following hold. For all  $\delta > 0$  sufficiently small, and, given  $d > 0$  and a length  $r > 0$  with  $0 < d, r \leq 1/2$ , there is  $\hat{\epsilon}(\epsilon', \epsilon, \delta, d, r) > 0$  such that the following hold for all  $\hat{\epsilon} < \hat{\epsilon}(\epsilon', \epsilon, \delta, d, r)$ . Suppose that  $B_{\lambda^2 g_n}(x, 1)$  is within  $\hat{\epsilon}$  of a 2-dimensional Alexandrov ball  $B(x, 1)$  of curvature  $\geq -1$ . Suppose that  $A_1, A_2, B_1$  are compact subsets of  $B(x, 1)$ . Let  $F = (f_1, f_2): B(x, 1) \rightarrow \mathbb{R}^2$  where  $f_1 = \frac{1}{2}(d(A_1, \cdot) - d(B_1, \cdot))$  and  $f_2 = d(A_2, \cdot)$ . Let  $D = F^{-1}(R)$  where  $R$  is the rectangle  $[a, a'] \times [c, c']$  with side-lengths,  $a' - a$  and  $c' - c$ , each at least  $r$ , and suppose that each of  $A_1, A_2, B_1$  are at distance at least  $d$  from  $D$ . Suppose also that for each  $z \in D$  there is a point  $b(z)$  at distance at least  $d$  from  $z$  such that the subsets  $A'_1, A'_2, B'_1, b(z)'$  form a  $(2, \delta)$ -strainer in the tangent sphere  $S_z$ . Suppose that  $\tilde{A}_1, \tilde{A}_2, \tilde{B}_1$  are compact subsets of  $B_{\lambda^2 g_n}(x, 1)$  within  $\hat{\epsilon}$  of  $A_1, A_2, B_1$ , respectively, and let  $\tilde{F}: B_{\lambda^2 g_n}(x, 1) \rightarrow \mathbb{R}^2$  be the map given by  $\tilde{F}$  where  $\tilde{F} = (\tilde{f}_1, \tilde{f}_2)$  with  $\tilde{f}_1 = \frac{1}{2}(d(\tilde{A}_1, \cdot) - d(\tilde{B}_1, \cdot))$  and  $\tilde{f}_2 = d(\tilde{A}_2, \cdot)$ . Set  $\tilde{D} = \tilde{F}^{-1}(R)$ . Then:

1. For every  $\tilde{z} \in \tilde{D}$  there is an  $S^1$ -product neighborhood with  $\epsilon$ -control,  $\varphi: S^1 \times B(0, \epsilon^{-1}) \rightarrow B_{\lambda^2 g_n}(x, 1)$  centered at  $\tilde{z}$ .
2. The map  $\tilde{F}: \tilde{D} \rightarrow R$  is a topological  $S^1$ -fibration.

3. For any  $S^1$ -product neighborhood with  $\epsilon$ -control  $\varphi: S^1 \times B(0, \epsilon^{-1}) \rightarrow B_{\lambda^2 g_n}(x, 1)$ , any fiber of  $\tilde{F}|_{\tilde{D}}$  through any point of  $\varphi(S^1 \times B(0, \epsilon^{-1}/2))$  is contained in  $S^1 \times B(0, \epsilon^{-1})$  and is a circle that is within  $\epsilon'$  of orthogonal to the horizontal subspaces<sup>9</sup> of the  $S^1$ -product structure and meets each horizontal subspace in a single point.

*Proof.* Fix  $\epsilon' > 0$  and  $\epsilon > 0$ . Eventually we will put conditions on the size of  $\epsilon$ , but for now it is simply fixed. Suppose that we have constants and balls satisfying the hypothesis of the proposition. Given  $\mu > 0$ , according to Lemma 10.10, if  $\delta > 0$  is sufficiently small there is  $d' > 0$  depending on  $d$  and  $\delta$  such that  $B(x, 1)$  is interior  $\mu$ -flat at every point of  $D$  on all scales  $\leq d'$ . Thus, provided that  $\delta$  is sufficiently small and that  $\hat{\epsilon}$  is sufficiently small (given  $d'$  and  $\epsilon$ ), it follows from Lemma 11.1 that every point of  $\tilde{R}$  is the center of an  $S^1$ -product neighborhood with  $\epsilon$ -control.

Provided that  $\delta$  is sufficiently small, and given  $\delta$  and  $d$ , provided that  $\hat{\epsilon}$  is sufficiently small, it follows from Theorem 12.7 of [3] the fibers of  $\tilde{F}|_{\tilde{D}}$  are compact, connected 1-manifolds with boundary in the boundary of  $B_{\lambda^2 g_n}(x, 1)$ . Since this ball is disjoint from the boundary, this implies that the fibers of  $\tilde{F}|_{\tilde{D}}$  are circles. Furthermore, by Theorem 11.14 of [3] given  $q \in \tilde{D}$  and a sequence  $q_k \in \tilde{F}^{-1}(\tilde{F}(q))$  converging to  $q$ , any limit  $\tau$  in the tangent sphere  $S_q$  of the directions of any subsequence of secant geodesics  $qq_k$  satisfies  $\tilde{f}'_i(\tau) = 0$  for  $i = 1, 2$  (see also, Lemma 9.45).

Now suppose that we have a sequence  $\epsilon_k \rightarrow 0$ , the other constants (also indexed by  $k$ ) sufficiently small for each  $k$  so that the results of the previous two paragraphs hold, and examples indexed by  $k$  satisfying the hypothesis for  $\epsilon_k$  and the other constants. The following holds at any point  $\tilde{z}_k \in \tilde{D}_k$ . Let  $\varphi_k: S^1 \times B(0, \epsilon) \rightarrow B_{\lambda^2 g_n}(x, 1)$  be an  $S^1$ -product structure with  $\epsilon$ -control with the property that  $\tilde{z}_k \in \varphi_k(S^1 \times B(0, \epsilon^{-1}/2))$ . Fix geodesics  $\gamma_{1,k}, \gamma_{2,k}, \gamma_{3,k}, \gamma_{4,k}$  from  $\tilde{z}_k$  to  $A_{1,k}, A_{2,k}, B_{1,k}, b(\tilde{z}_k)$ , respectively. Passing to a subsequence and rescaling the metric on the  $S^1$ -product neighborhoods gives a sequence converging to  $S^1 \times \mathbb{R}^2$  with the  $\tilde{z}_k$  converging to the central point  $p = (1, 0)$ . The pre-image under  $\varphi_k$  of these 4 geodesics converge as  $k \rightarrow \infty$  to 4 horizontal straight lines  $L_1, L_2, L_3, L_4$  in  $S^1 \times \mathbb{R}^2$ . By Proposition 9.23 (or more precisely by Addendum 9.24) we can choose the Euclidean coordinates on the  $\mathbb{R}^2$ -factor of the limit so that the  $L_1, L_3$  are the negative and positive  $x$ -axis and  $L_2$  and  $L_4$  are the positive and negative  $y$ -axis. The standard contradiction argument shows that given  $\epsilon' > 0$  provided that  $\epsilon > 0$  sufficiently small, the subspace of the tangent sphere  $S_{\tilde{z}}$  to at any point  $\tilde{z} \in \tilde{D}$  that is the intersection of the zero loci  $f'_1$  and  $f'_2$  in the tangent sphere at  $\tilde{z}$  consists of two points within  $\epsilon'$  of the tangent directions to any  $S^1$ -factor in an  $S^1$ -product structure with  $\epsilon$ -control. This means that, given  $\epsilon'$ , provided that  $\epsilon > 0$  is sufficiently small, all limiting directions of secant lines as in Item 3 are within  $\epsilon'$  of orthogonal to the horizontal plane in any  $S^1$ -product structure with  $\epsilon$ -control.

The last thing to see is that, provided that  $\epsilon > 0$  is sufficiently small, the fibers of  $\tilde{F}$  meet each horizontal plane at most once. But, given what we established in the previous paragraph, that is clear from Lemma 9.45.  $\square$

<sup>9</sup>This means that fixing any  $q$  in the neighborhood the limit as  $q' \in F^{-1}(F(q))$  approaches  $q$  of the geodesic in the  $S^1$ -product structure from  $q$  to  $q'$  is within  $\epsilon'$  of the  $S^1$ -direction.

**Remark 11.5.** The argument in the next to the last paragraph of the proof can be enhanced allowing us to use the local  $S^1$ -product structures with  $\epsilon$ -control to establish that  $\tilde{F}$  is a fibration with fibers that are circles close to the fibers of the local  $S^1$ -product structures. This allows one to avoid the reference to [3].

**Addendum 11.6.** We formulated this proposition for three fixed compact sets  $A_1, A_2, B_1$ , a rectangle  $R$  defined by coordinate functions  $f_1 = \frac{1}{2}(d(A_1, \cdot) - d(B_1, \cdot))$  and  $f_2 = d(A_2, \cdot)$  and a fourth point  $b(z)$ , depending on  $z \in R$ , forming  $(2, \delta)$ -strainers. But it can equally well be formulated with two fixed compact sets  $A_1, A_2$ , a rectangle defined by coordinate functions  $f_1 = d(A_1, \cdot)$  and  $f_2 = d(A_2, \cdot)$ , and points  $b_1(z), b_2(z)$  depending on  $z \in R$  so that for each  $z \in R$  the  $A_1, A_2, b_1(z), b_2(z)$  form a  $(2, \delta)$ -strainer at  $z$ . Details are left to the reader.

**Corollary 11.7.** *There is a universal constant  $C < \infty$  such that under the hypotheses of Lemma 11.1 the diameters of the circle factors of the  $S^1$ -product structure in the metric  $\lambda^2 g_n$  are bounded above  $C\hat{\epsilon}$ .*

*Proof.* Suppose that  $B_{\lambda^2 g_n}(x, 1)$  is within  $\hat{\epsilon}$  of  $B(\bar{x}, 1)$  which is interior  $\mu$ -flat at scale  $r_2$  at  $\bar{y} \in B(\bar{x}, 7/8)$ . Let  $a_1, a_2, b_1, b_2$  be a  $(2, \delta)$  strainer of size  $r_2$  for  $\bar{y}$ . (Here,  $\delta$  depends on  $\mu$  and goes to zero as  $\mu$  does.) Suppose that  $y \in B_{\lambda^2 g_n}(x, 1)$  is within  $\hat{\epsilon}$  of  $\bar{y}$  and  $\tilde{a}_1, \tilde{a}_2, \tilde{b}_1, \tilde{b}_2$  are within  $\hat{\epsilon}$  of  $a_1, a_2, b_1, b_2$  and hence these latter four points form a  $(2, \delta')$ -strainer at every point of a ball  $B(y, r)$  for some  $r > 0$  depending only on  $r_2$ . (Here,  $\delta'$  approaches  $\delta$  as  $\hat{\epsilon}$  and goes to zero.) Now let  $\varphi: S^1 \times B(0, \epsilon^{-1}) \cong U \subset B(x, 1)$  be an  $S^1$ -product neighborhood centered at  $y$ . By the last statement in Lemma 11.1, provided that  $\hat{\epsilon}$  is sufficiently small, this neighborhood is contained in  $B(y, r)$ . We have the map  $F = (d(\tilde{a}_1, \cdot), d(\tilde{a}_2, \cdot)): U \rightarrow \mathbb{R}^2$ . Let  $p, q$  be points of the fiber  $F^{-1}(F(y))$  through  $y$ . Then for  $i = 1, 2$  we have  $d(\tilde{a}_i, p) = d(\tilde{a}_i, q)$ . Let  $\bar{p}, \bar{q} \in B(\bar{x}, 1)$  be within  $\hat{\epsilon}$  of  $p$  and  $q$ . It follows that for  $i = 1, 2$  we have  $|d(a_i, \bar{p}) - d(a_i, \bar{q})| < 4\hat{\epsilon}$ . This means that under the map  $\bar{F} = (d(a_1, \cdot), d(a_2, \cdot))$  we have  $|\bar{F}(\bar{p}) - \bar{F}(\bar{q})| < 8\hat{\epsilon}$ . Since for  $\mu$  and  $\hat{\epsilon}$  sufficiently small,  $\bar{F}$  is a 2 almost isometry, we see that  $d(\bar{p}, \bar{q}) < 16\hat{\epsilon}$ , and hence  $d(p, q) < 20\hat{\epsilon}$ .

This shows that the diameter of the fibers of  $F$  are bounded above by  $20\hat{\epsilon}$ . It follows from the previous proposition that the diameter of a fiber of  $F$  through the central point of a  $S^1$ -product neighborhood is within a factor of 2 of the diameter of that central fiber. This completes the proof of the corollary.  $\square$

Next, we establish a truly global result obtained by piecing together the  $S^1$ -product structures to form a global  $S^1$ -fibration.

## 11.2 The global $S^1$ -fibration

**Proposition 11.8.** *For all  $\epsilon' > 0$  sufficiently small the following holds for all  $\epsilon > 0$  less than a positive constant  $\epsilon_1(\epsilon')$ . Let  $(M, g)$  be a Riemannian manifold. Suppose that  $K \subset M$  is a compact subset and each  $x \in K$  is the center of an  $S^1$ -product neighborhood with  $\epsilon$ -control. Then there is a finite collection  $\{\varphi_i: S^1 \times B(0, \epsilon^{-1})\}$  of  $S^1$ -product structures with  $\epsilon$ -control, constants  $T_i < \epsilon^{-1}$ , and embeddings  $\psi_i: S^1 \times$*

$B(0, T_i) \rightarrow S^1 \times B(0, \epsilon^{-1})$  that are within  $\epsilon'$  of the inclusion in the  $C^{[1/\epsilon']}$ -topology with the following properties:

1.  $K$  is contained in

$$V = \cup_i \varphi_i \circ \psi_i(S^1 \times B(0, T_i)).$$

2. There is an  $S^1$ -fibration structure on  $V$  whose restriction to each  $\varphi_i \circ \psi_i(S^1 \times B(0, T_i))$  agrees with the fibration structure induced by the product structure.

Define a circle action on each  $\varphi_i \circ \psi_i(S^1 \times B(0, T_i))$  as follows. For any  $p_1, p_2$  in the same fiber  $F$ , let  $\ell(p_1, p_2)$  be the length of the arc on  $F$  from  $p_1$  to  $p_2$  where the arc moves in the direction of the orientation on the  $S^1$ -factor and the length is measured using  $g$ . Similarly, let  $\ell(F)$  denote the length of  $F$  in  $g$ . Then  $\theta \cdot p_1 = p_2$  when  $\theta = 2\pi\ell(p_1, p_2)/\ell(F)$ . Pulling this local action back via  $(\varphi_i \circ \psi_i)^{-1}$  gives an action of  $S^1$  on  $S^1 \times B(0, T_i)$  that is within  $\epsilon'$  in the  $C^{[1/\epsilon']}$ -topology of the standard action coming from the product structure.

The proof of this proposition takes up this entire subsection. For  $\epsilon > 0$  sufficiently small, we set  $N = [1/\epsilon]$ . Recall that an  $S^1$ -product neighborhood  $U \subset M$  is the image  $\varphi(S^1 \times B(0, \epsilon^{-1}))$  with the property that there is  $\lambda_U > 0$  such that  $\varphi^*(\lambda_U^2 g)$  is within  $\epsilon$  in the  $C^N$ -topology of  $g_{\text{std}}$ , the product of the Riemannian metric of length 1 on  $S^1$  and the usual Euclidean metric on the ball  $B(0, \epsilon^{-1})$  in the plane.

### 11.2.1 Comparing the standard metrics on the overlap

The first thing to do is to show that on the overlap of  $S^1$ -product neighborhoods the standard metrics are close.

**Claim 11.9.** *Given  $\epsilon' > 0$  there is  $\epsilon > 0$  such that the following holds. Suppose that  $U_1 = \varphi_1(S^1 \times B(0, \epsilon^{-1}))$  and  $U_2 = \varphi_2(S^1 \times B(0, \epsilon^{-1}))$  are  $S^1$ -product neighborhoods with  $\epsilon$ -control in a Riemannian 3-manifold  $(M, g)$ . Suppose that there is a point*

$$x \in \varphi_1(S^1 \times B(0, \epsilon^{-1}/2)) \cap \varphi_2(S^1 \times B(0, \epsilon^{-1}/2)).$$

*Then for  $i = 1, 2$  the circle factor  $F_i$  through  $x$  in the product structure on  $U_i$  is within  $\epsilon'$  of vertical in the product structure of  $U_{3-i}$ . The length of this fiber is between  $1 - \epsilon'$  and  $1 + \epsilon'$  times the length of any circle factor in the product structure of  $U_{3-i}$  as is the ratio  $\lambda_{U_1}/\lambda_{U_2}$ . The homotopy class of  $F_i$  generates  $\pi_1(U_{3-i})$ . [All lengths are measured using  $g$ .]*

*Proof.* Without loss of generality we can assume that  $\lambda_{U_2} \geq \lambda_{U_1}$ . Let  $\zeta$  be the  $g$ -shortest homotopically non-trivial loop through  $x$  in  $U_2$ . Its  $g$ -length is close to  $\lambda_{U_2}^{-1}$ . Hence, it is contained in  $U_1$  and its length with respect to the product metric  $g_{\text{std}}$  on  $U_1$  is close to  $(\lambda_{U_1}/\lambda_{U_2}) \leq 1$ . Let us suppose that it is homotopically trivial in  $U_1$ . Then it bounds a disk contained in the  $g$ -neighborhood of size  $2\lambda_{U_2}^{-1}$  of  $x$ . This disk is then contained in  $U_2$ , which is a contradiction. It follows that  $\zeta$  is a homotopically non-trivial loop in  $U_1$  through  $x$ . Since its length in the metric  $g_{\text{std}}$  on  $U_1$  is close  $\lambda_{U_1}/\lambda_{U_2} \leq 1$ , the loop  $\zeta$  generates the fundamental group of  $U_1$ . It follows that  $\lambda_{U_1}/\lambda_{U_2}$  must be close to one. The errors in these estimates go to zero as  $\epsilon$  tends to zero.  $\square$

**Corollary 11.10.** *We continue with the notation of the previous claim. Given  $\epsilon' > 0$ , if  $\epsilon > 0$  is sufficiently small then the restrictions of  $(\varphi_1^{-1})^*g_{\text{std}}$  and  $(\varphi_2^{-1})^*g_{\text{std}}$  to  $\varphi_1(S^1 \times B(0, \epsilon^{-1}/2)) \cap \varphi_2(S^1 \times B(0, \epsilon^{-1}/2))$  are within  $\epsilon'$  in the  $C^N$ -topology.*

### 11.2.2 Bounding the intersections

Now we turn to constructing a finite cover with a uniformly bounded number of neighborhoods meeting any given neighborhood.

**Claim 11.11.** *Fix  $R < \infty$  and  $\epsilon' > 0$ . Then for all  $\epsilon > 0$  sufficiently small (in particular  $\epsilon^{-1} > R + 1$ ), there is a finite collection of  $S^1$ -product neighborhoods with  $\epsilon$ -control*

$$\varphi_1(S^1 \times B(0, \epsilon^{-1})), \dots, \varphi_T(S^1 \times B(0, \epsilon^{-1}))$$

*such that the union of the images  $U'_i = \varphi_i(S^1 \times B(0, R))$  cover  $K$ , and the  $\varphi_i(S^1 \times B(0, R/3))$  are disjoint. Furthermore for every  $i, j$ , the Riemannian metrics  $(\varphi_i^{-1})^*g_{\text{std}}$  and  $(\varphi_j^{-1})^*g_{\text{std}}$  are within  $\epsilon'$  in the  $C^N$ -topology for Riemannian metrics on*

$$\varphi_i(S^1 \times B(0, \epsilon^{-1}/2)) \cap \varphi_j(S^1 \times B(0, \epsilon^{-1}/2)).$$

*Proof.* Fix  $\epsilon > 0$  sufficiently small. If  $\varphi_i(S^1 \times B(0, R/3)) \cap \varphi_j(S^1 \times B(0, R/3)) \neq \emptyset$ , then, by the previous result, the standard metrics on the two images almost agree, and in particular, their union is contained in  $\varphi_i(S^1 \times B(0, R))$ . Take a collection  $\{\widehat{U}_i = \varphi_i(S^1 \times B(0, \epsilon^{-1}))\}$  of  $S^1$ -product neighborhoods with  $\epsilon$ -control centered at points of  $K$ , maximal with respect to the property that the  $\varphi_i(S^1 \times B(0, R/3))$  are disjoint. Then the  $U'_i = \varphi_i(S^1 \times B(0, R))$  cover  $K$ . If we have chosen  $\epsilon > 0$  sufficiently small, the last statement follows from the previous result.  $\square$

**Claim 11.12.** *Given  $R > 4$ , there is an integer  $C = C(R)$  such that following holds for all  $\epsilon > 0$  sufficiently small. Let  $(M, g)$  be a Riemannian 3-manifold. Suppose that we have a collection  $\{\widehat{U}_i = \varphi_i(S^1 \times B(0, \epsilon^{-1}))\}_i$  of  $S^1$ -product neighborhoods with  $\epsilon$ -control. Let  $U_i$  be the image of  $\varphi_i(S^1 \times B(0, R + 1))$ . Suppose also that  $\varphi_i(S^1 \times B(0, R/3)) \cap \varphi_j(S^1 \times B(0, R/3)) = \emptyset$  for all  $i \neq j$ . Then for each  $i$  the number of  $j$  for which  $U_i \cap U_j \neq \emptyset$  is at most  $C - 1$ .*

*Proof.* This is immediate from volume comparison and the fact that the standard metrics almost agree on the overlaps of the  $U_i$ .  $\square$

For  $R < \epsilon^{-1}$  we define a *reduced  $S^1$ -product structure with  $\epsilon$ -control of size  $R$*  to be an embedding  $\varphi: S^1 \times B(0, R) \rightarrow M$  with the property that there is  $\lambda > 0$  such that  $\varphi^*\lambda^2g$  is within  $\epsilon$  in the  $C^N$ -topology to the standard product metric  $g_{\text{std}}$  on this product.

Fix  $4 < R < \epsilon^{-1}$  and a covering  $\{U_a\}_{a \in A}$  of  $K$  as in Claim 11.12. It follows directly from Claim 11.12 that we can divide the open sets  $\{U_a\}$  into  $C$  groups  $\mathcal{U}_1, \dots, \mathcal{U}_C$  with the following properties:

1. Each  $\mathcal{U}_i$  is the union of a finite number of the  $U_a$ , denoted  $U_{i,1}, \dots, U_{i,j_0(i)}$  that are pairwise disjoint in  $M$ .

2. Each  $U_a$  in the original collection occurs as exactly one of the  $U_{i,j}$ , so that in particular, setting  $\mathcal{U}'_i$  equal to the images  $\varphi_{i,j}(S^1 \times B(0, R))$  for  $1 \leq j \leq j_0(i)$ , the union  $\cup_{i=1}^C \mathcal{U}'_i$  covers  $K$ .

**Definition 11.13.** For each  $0 \leq D \leq 1$  we define  $\mathcal{U}_i^{[D]}$  to be the union of the images  $\varphi_{i,j}(S^1 \times B(0, R + 1 - D))$ . Notice that  $\mathcal{U}'_i = \mathcal{U}_i^{[1]}$ .

### 11.2.3 The Gluing

Suppose that we have an open subset  $W \subset M$  that is the union of restrictions of  $S^1$ -product neighborhoods with  $\alpha$ -control to subsets  $U_i = \varphi_i(S^1 \times B(0, R'))$  for some  $R \leq R' \leq R + 1$ , and suppose that the circle fibrations of the various  $U_i$  are compatible so that they define a circle fibration on  $W$ . Suppose also that we have a reduced  $S^1$ -product structure with  $\epsilon$  control  $\varphi: S^1 \times B(0, R + 2) \rightarrow M$ . Let  $U = \varphi(S^1 \times B(0, R + 1))$ . Assuming that  $\alpha$  and  $\epsilon$  are sufficiently small, let us define a map from the saturation,  $\text{Sat}_W(U \cap W)$ , of  $U \cap W$  under the  $S^1$ -fibration on  $W$  to  $S^1 \times B(0, R + 2)$ . For  $\alpha$  and  $\epsilon$  sufficiently small  $\text{Sat}_W(U \cap W)$  is contained in  $\varphi(S^1 \times B(0, R + 2))$ . Suppose that  $p$  is a point of  $\text{Sat}_W(U \cap W)$ , say  $p = \varphi(\theta, x)$ . Let  $F_p$  be the fiber of the fibration structure on  $W$  through  $p$ . For each  $q \in F_p$  we have  $(\theta(q), x(q))$  defined by  $\varphi^{-1}(q) = (\theta(q), x(q))$ . We form

$$\hat{x}(p) = \frac{1}{\ell(F_p)} \int_{F_p} x(q) d\mu_{F_p},$$

where  $d\mu_{F_p}$  is the measure induced by the restriction of the Riemannian metric of  $M$  to  $F_p$  and  $\ell(F_p)$  is the length of this circle in  $M$ , and define the map

$$\psi(p) = (\theta(p), \hat{x}(p)).$$

The following is obvious from the definitions

**Claim 11.14.** *If  $F$  is an orbit of the  $S^1$ -fibration on  $W$  passing through a point of  $U$ , then  $\hat{x}: F \rightarrow B(0, R + 2)$  is constant.*

**Corollary 11.15.** *Given  $\epsilon_1 > 0$ , then for all  $\alpha, \epsilon > 0$  sufficiently small, the map  $\hat{x}: \text{Sat}_W(U \cap W) \rightarrow B(0, R + 2)$  is within  $\epsilon_1$  in the  $C^{N+1}$ -topology of the restriction to  $\text{Sat}_W(U \cap W) \subset U$  of the composition of  $\varphi^{-1}$  followed by the projection in product structure to  $B(0, R + 2)$ .*

*Proof.* It follows immediately from Corollary 11.10 that the fibers of the  $S^1$ -fibration on  $\text{Sat}_W(U \cap W)$  induced from the fibration on  $W$  are geodesics in a metric that is  $C^N$ -close to the metric  $g_{\text{std}}$  on  $U$ . From this we see that the map  $p \mapsto \hat{x}(p)$  is  $C^{N+1}$ -close to the composition of  $\varphi^{-1}$  with the projection to  $B(0, R + 2)$  with the same error estimate.  $\square$

It follows from Corollary 11.15 that given  $\epsilon_1 > 0$ , there is a constant  $\alpha_0(\epsilon_1) > 0$  such that if  $\alpha$  and  $\epsilon$  are less than  $\alpha_0(\epsilon_1)$ , then we can define a map  $\psi: \text{Sat}_W(U \cap W) \rightarrow S^1 \times B(0, R + 2)$  by sending  $p = \varphi(\theta, x)$  to  $\psi(p) = (\theta(p), \hat{x}(p))$ . Again invoking Corollary 11.15, we see that:



**Corollary 11.16.** *Provided that  $\alpha$  and  $\epsilon$  are less than  $\alpha_0(\epsilon_1)$ , the composition*

$$\text{Sat}_W(U \cap W) \xrightarrow{\psi} S^1 \times B(0, R+2) \xrightarrow{\varphi} \varphi(S^1 \times B(0, R+2))$$

*is within  $\epsilon_1$  in the  $C^{N+1}$ -topology of the inclusion of  $\text{Sat}_W(U \cap W) \subset \varphi(S^1 \times B(0, R+2))$ .*

Let  $\beta: [0, R'] \rightarrow [0, 1]$  be a weakly monotone function that is identically 1 near  $R'$  and with  $\beta^{-1}(0) = [0, R' - 1/C]$ . We define  $\beta_i: U_i \rightarrow [0, 1]$  by  $\beta_i(\varphi_i(\theta, x)) = \beta(|x|)$ , and extend  $\beta_i$  to all of  $M$  by defining it to be identically 1 on  $M \setminus U_i$ . For all  $i$  such that  $U_i \cap U \neq \emptyset$ , the gradients of the  $\beta_i$  with respect to  $\lambda_U^2 g$  are bounded independent of  $i$ . (Recall that  $\lambda_U^2 g$  is the multiple of  $g$  which is close to the standard product metric  $g_{\text{std}}$  on  $U$ .) We set  $\hat{\beta}: M \rightarrow [0, 1]$  equal to the product over the  $i$  of the  $\beta_i$ . This function is identically 1 in the complement of  $W$  and the restriction to  $U$  of  $\hat{\beta}$  has a gradient with respect to  $g_{\text{std}}$  that is bounded depending only on  $C$ . Define  $\Psi: U \rightarrow S^1 \times B(0, R+2)$  by

$$\Psi(p) = \hat{\beta}(p)\varphi^{-1}(p) + (1 - \hat{\beta}(p))\psi(p),$$

where we use the local linear structure on  $S^1 \times B(0, \epsilon^{-1})$  to form the linear combination.

**Claim 11.17.** *Given  $\epsilon_1$  there is  $\alpha_1 = \alpha_1(\epsilon_1) > 0$  such that if  $\alpha$  and  $\epsilon$  are less than  $\alpha_1$ , then  $\Psi$  is within  $\epsilon_1$  of  $\varphi^{-1}$  in the  $C^{N+1}$ -topology using the metrics  $\lambda_U^2 g$  on the domain and  $g_{\text{std}}$  on the range.*

*Proof.* This follows immediately from Corollary 11.16.  $\square$

We set  $W' \subset W$  equal to  $\beta^{-1}(0)$ . The following is immediate from the definitions and Claim 11.17.

**Claim 11.18.** *Fix  $0 < \epsilon_1 \ll 1/C$  and  $\alpha_1 = \alpha_1(\epsilon_1)$  from Claim 11.17. Fix  $0 < \epsilon, \alpha < \epsilon_1$ . With these conditions on the parameters we have:  $W'$  is the union of  $\varphi_i(S^1 \times B(0, R''))$  where  $R'' = R' - 1/C$ . In particular,  $W'$  is saturated under the  $S^1$ -fibration structure on  $W$ . The image of  $\Psi$  contains  $S^1 \times B(0, R+1 - 1/C)$ . Setting  $\varphi': S^1 \times B(0, R+1 - 1/C) \rightarrow M$  equal to the restriction of the inverse of  $\Psi$ , we have*

1.  $\varphi'$  is a reduced  $S^1$ -product neighborhood with  $\epsilon'$ -control of size  $R+1 - 1/C$ .
2. If  $\varphi'(\theta, x) \subset W'$ , then  $\varphi'(S^1 \times \{x\})$  is a fiber of the  $S^1$ -fibration on  $W'$ , so that the  $S^1$ -fibration structure on  $U'$  coming from the  $S^1$ -product structure and the given  $S^1$ -fibration structure on  $W'$  are compatible on the overlap  $U' \cap W'$  and hence together define an  $S^1$ -fibration structure on  $W' \cup \varphi'(S^1 \times B(0, R+1 - 1/C))$ .
3. For any  $T \leq R+1$ , the image  $\varphi'(S^1 \times B(0, T))$  contains  $\varphi(S^1 \times B(0, T - 1/C))$ .

We denote the image  $\varphi'(S^1 \times B(0, R+1-1/C))$  by  $U'$ . The claim shows that, at the expense of shrinking  $W$  to  $W'$  and at the expense of deforming  $\varphi$  slightly in the  $C^N$ -topology to a reduced  $S^1$ -product structure with  $\epsilon'$ -control,  $\varphi': S^1 \times B(0, R+1-1/C) \rightarrow M$ , we can make the  $S^1$ -fibrations compatible on the overlap, so that together they define an  $S^1$ -fibration on the union  $W' \cup U'$ . One more remark is in order. If we have not a single reduced  $S^1$ -product neighborhood with  $\epsilon$ -control  $U$ , but rather a collection of them  $U_{i_0,j}$ ,  $1 \leq j \leq j_0(i_0)$ , whose images are disjoint, then we can perform this operation simultaneously on all of them, so as to deform them all to  $S^1$ -product neighborhoods with  $\epsilon_1$ -control compatible with the circle fibration on  $W'$ .

Now we are ready to apply this gluing argument by induction to the  $\mathcal{U}_1, \dots, \mathcal{U}_C$ . We begin with  $\mathcal{U}_1$ . In the inductive step, deforming and gluing in  $\mathcal{U}_{i_0}$ , we cut down the  $S^1$ -product neighborhoods in the neighborhoods that make up the previous  $\mathcal{U}_i$  by  $1/C$ . The deformation of the maps  $\varphi_{i_0,j}$  produces a reduced  $S^1$ -product neighborhood with  $\epsilon_1$ -control where the amount of the deformation and  $\epsilon_1$  depend only on the control we have at the previous step. Thus, we can iterate this construction  $C$  times keeping a fixed control,  $\epsilon'$ , on all the  $S^1$ -product neighborhoods and a given control on the size of the deformations, provided only that we arrange that the original control,  $\epsilon$ , is sufficiently small given  $C$ ,  $\epsilon'$ , and the desired control on all deformations.

It follows from the second conclusion of Claim 11.18 that the  $S^1$ -fibrations induced by the product structures on the deformed  $\mathcal{U}_i$  are compatible and hence define a global  $S^1$ -fibration on the union. It follows from the third conclusion of Claim 11.18 that the union of the deformed  $S^1$ -product neighborhoods contains  $K$ . All the estimates stated in Proposition 11.8 are immediate from the construction. This completes the proof of Proposition 11.8.

### 11.3 Balls centered at points of $\partial M_n$

The results about the generic behavior over interior points of the base are enough to establish what the neighborhoods of the boundary of the  $M_n$  look like.

**Proposition 11.19.** *Fix  $\hat{\epsilon} > 0$ . For all  $n$  sufficiently large, for any point  $x \in \partial M_n$  the ball  $B_{g'_n(x)}(x, 1)$  is within  $\hat{\epsilon}$  of the interval of length 1, and  $x$  is within  $\hat{\epsilon}$  of the endpoint of  $J$ .*

*Proof.* Suppose that the result is not true. Then after passing to a subsequence (in  $n$ ) we can suppose that for each  $n$  we have  $x_n \in \partial M_n$  for which the result does not hold. Let  $T_n$  be the component of  $\partial M_n$  containing  $x_n$  and let  $C_n$  be the topologically trivial collar containing the neighborhood of size 1 of  $T_n$ . Since  $\partial M_n$  is convex and  $\rho_n \leq \text{diam } M_n/2$ , the balls  $B_{g'_n(x_n)}(x_n, 1)$  are Alexandrov balls. Because the curvatures on the topologically trivial collar which includes the neighborhood of size 1 about  $\partial M_n$ , are bounded above by  $-3/16$ , it follows that  $\rho_n(x_n) \leq \sqrt{16/3}$ . Hence,  $B_n = B_{g'_n(x_n)}(x_n, 1/4)$  is contained in  $C_n$ . After passing to a subsequence, we shall show that the  $B_{g'_n(x_n)}(x_n, 1/4) \subset C_n$  converge to the interval. Assuming this, it follows that the  $B_{g'_n(x_n)}(x_n, 1)$  also converge to an interval.

We have already remarked that because of the convexity of  $\partial M_n$ , the  $B_n$  are Alexandrov balls. Passing to a subsequence, there is a Gromov-Hausdorff limit  $J$  which is an Alexandrov ball of curvature  $\geq -1$  of diameter  $1/4$  centered at  $\bar{x} = \lim x_n$ . Because of the volume collapsing condition on the  $M_n$ , it follows that  $J$  is either of dimension 1 or 2. We rule out the possibility that  $\dim J = 2$ . Suppose to the contrary that the dimension of  $J$  is 2. Let  $\epsilon_n$  be the Gromov-Hausdorff distance from  $B_n$  to  $J$ . Fix  $\epsilon' > 0$  a universally small constant. We can suppose that  $\epsilon$  is sufficiently small so that Proposition 11.4 holds for the given values of  $\epsilon'$  and  $\epsilon$ . Fix  $\ell_0$  as in Proposition 9.36. Fix  $\delta > 0$  sufficiently small so that Proposition 11.4 holds. Then there is a point  $\bar{y}$  of  $J$  within distance  $\ell_0/2$  of  $\bar{x}$  that has a  $(2, \delta/2)$ -strainer  $\{a_1, a_2, b_1, b_2\}$  of some size  $d > 0$ . Then for  $0 < r < \ell_0/2$ , with  $r$  sufficiently small (depending on  $d$  and  $\delta$ ), the same set of four points form a  $(2, \delta)$ -strainer centered at any point of  $B(\bar{y}, r)$  of size  $d/2$ . We set  $F = (f_1, f_2): B(\bar{y}, 1) \rightarrow \mathbb{R}^2$  by defining  $f_1 = \frac{1}{2}(d(a_1, \cdot) - d(b_1, \cdot))$  and  $f_2 = d(a_2, \cdot)$ . Thus, there is a rectangle  $R$  in  $\mathbb{R}^2$  with side lengths  $r'$ , depending only on  $r, \delta$ , and  $d$  such  $D = F^{-1}(R)$  is contained in  $B(\bar{y}, r)$ . For all  $n$  sufficiently large  $\epsilon_n < \hat{\epsilon}(\epsilon', \epsilon, \delta, d/2, r')$  from Proposition 11.4 and also less than  $\ell_0/2$ . We lift the  $(2, \delta)$  strainer to four points  $\{\tilde{a}_1, \tilde{a}_2, \tilde{b}_1, \tilde{b}_2\}$  in  $B_{g'_n(x_n)}(x_n, 1/4)$  and define  $\tilde{F} = (\tilde{f}_1, \tilde{f}_2)$  with  $\tilde{f}_1 = \frac{1}{2}(d(\tilde{a}_1, \cdot) - d(\tilde{b}_1, \cdot))$  and  $\tilde{f}_2 = d(\tilde{a}_2, \cdot)$  and define  $\tilde{D} = \tilde{F}^{-1}(R)$ . According to Proposition 11.4, for all  $n$  sufficiently large  $\tilde{F}: \tilde{D} \rightarrow R$  is a locally trivial  $S^1$ -fibration. In particular,  $\pi_1(\tilde{D}) \cong \mathbb{Z}$ . Of course, for any point  $y_n \in B_{g'_n(x_n)}(x_n, 1/4)$  within  $\epsilon_n$  of  $\bar{y}$  the ball  $U_n = B_{\lambda^2 g_n}(y_n, r'/2)$  is contained in  $\tilde{D}$ . Consequently, the image,  $\Gamma_n$ , of the homomorphism  $\pi_1(U_n) \rightarrow \pi_1(C_n, x_n)$  induced by the inclusion mapping is either trivial or infinite cyclic and the quotient  $\pi_1(C_n, x_n)/\Gamma_n$  contains an infinite cyclic factor.

Since  $\rho(x_n) < 3$ , the diameter of  $T_n$  in the metric  $\rho_n^{-2}(x_n)g_n$  is at most  $3Kw_n$ , it follows that  $\pi_1(C_n, x_n)$  is generated by elements represented by loops based at  $x_n$  of length at most  $6Kw_n$ . In particular, there is a loop based at  $x_n$  of length at most  $6Kw_n$  whose image in  $\pi_1(C_n, x_n)/\Gamma_n$  is of infinite order. Since  $w_n \rightarrow 0$  as  $n \rightarrow \infty$ , this contradicts Proposition 9.36. This completes the proof that the Gromov-Hausdorff limit,  $J$ , is an interval.

Now let us show that the boundaries  $T_n$  converge to the endpoint of  $J$ . Since  $w_n \rightarrow 0$  as  $n \rightarrow \infty$ ,  $T_n$  converges to some point  $y \in J$ . Suppose to the contrary that  $y \in \text{int } J$ . Fix  $z \neq y$  in  $J$  and let  $z_n$  be an approximating point in  $C_n$ . Then, for all  $n$  sufficiently large, the distance function from  $z_n$  is regular in a neighborhood of  $T_n$ . According to Section 13 of [3] this implies that there is a neighborhood of  $T_n$  in  $C_n$  that is topologically a locally trivial fibration over an open interval. This is absurd since  $T_n$  is a boundary component of a manifold.

Since  $B_{g'_n(x_n)}(x_n, 1)$  has diameter 1, it follows that  $J$  is isometric to  $[0, 1]$ .  $\square$

#### 11.4 The interior cone points

**Proposition 11.20.** *For any  $\epsilon' > 0$  sufficiently small, for all  $\epsilon > 0$  less than a positive constant  $\epsilon_2(\epsilon')$  and for any  $a > 0$ , the following holds for all  $\mu > 0$  less than a positive constant  $\mu_3(\epsilon, a)$ , for any  $0 < r_2 \leq r_1 \leq 1$ , and for all  $\hat{\epsilon} > 0$  less than a positive constant  $\hat{\epsilon}_1(\epsilon, a, r_1, r_2)$ . Suppose that, for some  $n$ , there are a point*

$x \in M_n$  and a constant  $\lambda \geq \rho^{-1}(x)$  with the property that the ball  $B_{\lambda^2 g_n}(x, 1)$  is within  $\hat{\epsilon}$  of a 2-dimensional Alexandrov ball  $B(\bar{x}, 1)$  of curvature  $\geq -1$  and of area  $\geq a$  that is interior  $\mu$ -good at  $\bar{x}$  on scale  $r'$ , where  $r_2 \leq r' \leq r_1$ . Every point of  $U = B_{\lambda^2 g_n}(x, 3r'/4) \setminus B_{\lambda^2 g_n}(x, r'/4)$  is the center of an  $S^1$ -product neighborhood with  $\epsilon$ -control.

There is an open subset  $\tilde{U}$  of  $U$  containing  $B_{\lambda^2 g_n}(x, r'/2) \setminus B_{\lambda^2 g_n}(x, 3r'/8)$  with  $\tilde{U}$  being the total space of an  $S^1$ -fibration with fibers making angle within  $\epsilon'$  of  $\pi/2$  with the horizontal spaces of the  $S^1$ -product neighborhoods with  $\epsilon$ -control at every point of  $\tilde{U}$  and with the fibers isotopic to the  $S^1$ -factors by a small isotopy. Furthermore, given any such  $\tilde{U}$  and  $S^1$ -fibration there is a 2-torus in  $\tilde{U}$  that is invariant under the  $S^1$ -fibration structure and is contained in  $B_{\lambda^2 g_n}(x, r'/2)$ . This 2-torus is the boundary of a solid torus in  $B_{\lambda^2 g_n}(x, r'/2)$ .

*Proof.* First let us show that it suffices to prove the result when  $r_2 = r_1 = 1$ . For suppose that for every  $\epsilon' > 0$  sufficiently small and  $\epsilon > 0$  less than  $\epsilon_2(\epsilon')$  and  $a > 0$  we have positive constants  $\mu'_3(\epsilon, a)$  and  $\hat{\epsilon}'_1(\epsilon, a)$  so that the proposition holds for  $r_2 = r_1 = 1$ . Let  $a'$  be the positive constant associated to  $a$  by Lemma 10.8. Suppose  $\mu < \mu_3(\epsilon, a')$  and  $\hat{\epsilon} < r_2 \hat{\epsilon}'_1(\epsilon, a')$ . Given balls  $B_{\lambda^2 g_n}(x, 1)$  and  $B(\bar{x}, 1)$  as in the statement for these values of  $\mu$  and  $\hat{\epsilon}$  and  $a$ , and some  $r'$  with  $r_2 \leq r' \leq r_1$ . Then  $(1/r')B(\bar{x}, r')$  is interior  $\mu$ -good at scale 1 at  $\bar{x}$  of area  $\geq a'$ . On the other hand  $B_{(1/r')^2 \lambda^2 g_n}(x, 1)$  is within  $(1/r')\hat{\epsilon} < \hat{\epsilon}'_1(\epsilon, a')$  of  $(1/r')B(\bar{x}, r')$ . By our assumption that the result holds in the special case when  $r_2 = r_1$ , we see that the conclusion holds for  $B_{(1/r')^2 \lambda^2 g_n}(x, r')$  with  $r'$  replaced by 1. Hence, by rescaling we see that it holds for  $B_{\lambda^2 g_n}(x, 1)$  with the given value of  $r'$ .

This allows us to assume, as we shall, that  $r_2 = r_1 = 1$ . Suppose that there are sequences  $\mu_k \rightarrow 0$  and  $\hat{\epsilon}_k \rightarrow 0$  as  $k \rightarrow \infty$  and balls  $B_{\lambda_k^2 g_{n(k)}}(x_{n(k)}, 1)$  within  $\hat{\epsilon}_k$  of standard 2-dimensional balls  $B(\bar{x}_k, 1)$  of area  $\geq a$  that are interior  $\mu_k$ -good at  $\bar{x}_k$  on scale 1 and yet the conclusion of the proposition does not hold for any  $k$ . Passing to a subsequence, we can suppose that the  $B(\bar{x}_k, 1)$  converge to a 2-dimensional ball  $B(\bar{x}_\infty, 1)$  of curvature  $\geq -1$ . Because the  $\mu_k \rightarrow 0$ , it follows that  $B(\bar{x}_\infty, 1)$  is a circular cone of some cone angle  $\theta \leq 2\pi$ , which is bounded away from zero because  $a$  is greater than zero. Since the  $\hat{\epsilon}_k \rightarrow 0$ , the  $B_{\lambda_k^2 g_{n(k)}}(x_{n(k)}, 1)$  also converge to  $B(\bar{x}_\infty, 1)$ .

Let us first consider the case when  $\theta = 2\pi$  so that  $B(\bar{x}_\infty, 1)$  is isometric to a ball in  $\mathbb{R}^2$ . Then, for any  $\delta/2 > 0$  for all  $k$  sufficiently large there is a  $(2, \delta/2)$ -strainer  $\{\tilde{a}_1, \tilde{a}_2, \tilde{b}_1, \tilde{b}_2\}$  for  $x_{n(k)}$  of size  $1/2$  and there is  $d' > 0$  depending on  $\delta$  so that the same set of four points is a  $(2, \delta)$ -strainer at any point of  $B_{\lambda_k^2 g_{n(k)}}(x_{n(k)}, d')$ . Without loss of generality we can assume that  $d' \ll r'$ . It follows from Proposition 11.4 that for all  $k$  sufficiently large, there are  $0 < s < d'$  depending on  $d'$  and  $\delta$  and a closed subset  $W_{n(k)}$  containing  $B_{\lambda_k^2 g_{n(k)}}(x_{n(k)}, s)$  and contained in  $B_{\lambda_k^2 g_{n(k)}}(x_{n(k)}, d')$  such that the function  $\tilde{F} = (\tilde{f}_1, \tilde{f}_2)$  where  $\tilde{f}_1 = \frac{1}{2}(d(\tilde{a}_1, \cdot) - d(\tilde{b}_1, \cdot))$  and  $\tilde{f}_2 = d(\tilde{a}_2, \cdot)$  defines a projection mapping from  $W_{n(k)}$  to a closed rectangle in the plane which is the projection mapping of a fibration of  $W_{n(k)}$  by circles. Furthermore, by Lemma 11.1, for all  $k$  sufficiently large, there is an  $S^1$ -product neighborhood  $V$  with  $\epsilon$  control centered at  $x_{n(k)}$ . Also, according to Proposition 11.4

the circle of the fibration structure on  $W_{n(k)}$  passing through  $x_{n(k)}$  is almost orthogonal to the horizontal spaces of the  $S^1$ -product structure centered at that point and this circle is isotopic in  $V$  to the  $S^1$ -factor. This means that the closure of  $V$  is a solid torus contained in  $W_{n(k)}$  whose core is isotopic to the fiber of the fibration structure on  $W_{n(k)}$ . It follows that the inclusion of  $V \subset W_{n(k)}$  induces an isomorphism on fundamental groups, both groups being isomorphic to  $\mathbb{Z}$ . Also, it follows that  $W_{n(k)} \setminus V$  is homeomorphic to  $T^2 \times I$ . We have inclusions  $V \subset B_{\lambda_k^2 g_{n(k)}}(x_{n(k)}, s) \subset W_{n(k)} \subset B_{\lambda_k^2 g_{n(k)}}(x_{n(k)}, d')$ . For all  $k$  sufficiently large, the distance function from  $x_{n(k)}$  is regular on  $B_{\lambda_k^2 g_{n(k)}}(x_{n(k)}, 7/8) \setminus B_{\lambda_k^2 g_{n(k)}}(x_{n(k)}, s/2)$ , and consequently, the inclusion of the smaller ball into the larger induces an isomorphism on the fundamental group. It then follows from the sequence of inclusions that the fundamental group of  $B_{\lambda_k^2 g_{n(k)}}(x_{n(k)}, s)$  is isomorphic to  $\mathbb{Z}$  and hence the metric sphere  $S_{\lambda_k^2 g_{n(k)}}(x_{n(k)}, s)$  is a 2-torus. This 2-torus is contained in  $W_{n(k)} \setminus V$  and separates the two boundary components of this region. Since we have already seen that the difference  $W_{n(k)} \setminus V$  is homeomorphic to a product  $T^2 \times I$ , it follows that  $S_{\lambda_k^2 g_{n(k)}}(x_{n(k)}, s)$  is isotopic in  $W_{n(k)}$  to the boundary of  $W_{n(k)}$  and that  $B_{\lambda_k^2 g_{n(k)}}(x_{n(k)}, s)$  is a solid torus. Consequently, since the distance function from  $x_{n(k)}$  is regular on the pre-image of  $[s, 7/8]$  it follows that  $B_{\lambda_k^2 g_{n(k)}}(x_{n(k)}, a)$  is a solid torus for every  $a \in [s, 7/8]$ . It is immediate from Proposition 11.8 that there is an open subset  $\tilde{U}$  containing  $B_{\lambda^2 g_n}(x, r'/2) \setminus B_{\lambda^2 g_n}(x, 3r'/8)$  with  $\tilde{U}$  being the total space of an  $S^1$ -fibration with fibers making angle within  $\epsilon'$  of  $\pi/2$  with the horizontal spaces of the  $S^1$ -product neighborhoods with  $\epsilon$ -control at every point of  $\tilde{U}$  and with the fibers isotopic to the  $S^1$ -factors by a small isotopy. Furthermore, given any such  $\tilde{U}$  with such an  $S^1$ -fibration there is a compact sub-fibration  $X$  contained in it that separates the metric spheres  $S_{\lambda^2 g_n}(x, r'/2)$  and  $S_{\lambda^2 g_n}(x, r'/4)$ . One of the boundary components  $\partial_0 X$  of  $X$  must also separate these spheres. Of course,  $\partial_0 X$  is a 2-torus. Since the region between the metric spheres is homeomorphic to  $T^2 \times I$ , it follows that  $\partial_0 X$  is parallel to each and hence bounds an unknotted solid torus in  $B_{\lambda^2 g_n}(x, r'/2)$ . This contradiction proves the result in the case when the limiting 2-dimensional cone has cone angle  $\theta = 2\pi$ .

Now suppose that limiting the cone angle  $\theta$  is strictly less than  $2\pi$ . According to Proposition 9.49 the following holds for all  $k$  sufficiently large. There is  $x'_{n(k)} \in M_{n(k)}$  such that  $d_{\lambda_k^2 g_{n(k)}}(x_{n(k)}, x'_{n(k)}) \rightarrow 0$  as  $k \rightarrow \infty$  such that for each  $k$  sufficiently large, one of the following two alternatives holds:

1. the distance function from  $x'_{n(k)}$  has no critical points on  $B_{\lambda_k^2 g_{n(k)}}(x'_{n(k)}, 1/2) \setminus \{x'_{n(k)}\}$ , or
2. there is  $\zeta_k \rightarrow 0$  such that the distance function from  $x'_{n(k)}$  has no critical points in  $B_{\lambda_k^2 g_{n(k)}}(x'_{n(k)}, 1/2) \setminus \overline{B_{\lambda_k^2 g_{n(k)}}}(x'_{n(k)}, \zeta_k)$  and has a critical point at distance  $\zeta_k$  from  $x'_{n(k)}$ .

In Case 1 the level sets of the distance function are 2-spheres and the metric balls are topological 3-balls. Let us suppose that Case 2 holds. According to Proposition 9.49 after passing to a subsequence the rescaled balls  $\zeta_k^{-1} B_{\lambda_k^2 g_{n(k)}}(x'_{n(k)}, 1/2)$  converge

in the Gromov-Hausdorff topology to a complete 3-dimensional Alexandrov space of curvature  $\geq 0$ . By Proposition 9.46 the limit is actually a smooth, orientable Riemannian manifold of curvature  $\geq 0$  and the convergence is  $C^\infty$ . Thus, the limit has a soul which is either a point, a circle, or a compact surface of non-negative curvature.

**Claim 11.21.** *The soul is not a surface.*

*Proof.* If the soul is a surface, then either the limiting 3-manifold or its double covering is a Riemannian product of a surface with  $\mathbb{R}$ . The limit cannot be the product of a surface with  $\mathbb{R}$ , for if it were, by rescaling we see that the limit of the  $B_{\lambda_k^2 g_{n(k)}}(x_{n(k)}, 1)$  is one-dimensional. If the limiting 3-manifold is a non-orientable  $\mathbb{R}$ -bundle over that surface. It would then follow that given any  $\beta > 0$  there is  $R < \infty$  such that for all  $k$  sufficiently large any triangle  $ax'_{n(k)}b$  with  $|ax'_{n(k)}| = |bx'_{n(k)}| = R$  has comparison angle less than  $\beta$  at  $x'_{n(k)}$ . On the other hand, because the limit of the  $B_{\lambda_k^2 g_{n(k)}}(x'_{n(k)}, 1)$  is 2-dimensional, there is  $\beta_0 > 0$  such that for all  $k$  sufficiently large there are geodesics from  $x'_{n(k)}$  to points at a fixed positive distance that make a comparison angle at  $x'_{n(k)}$  which is least  $\beta_0$ . This contradicts the fact that comparison angles are lower continuous under limits.  $\square$

This shows if Case 2 holds then the soul of the limiting manifold is either a circle or a point, and hence the level sets  $d(x'_{n(k)}, \cdot)^{-1}(b)$  are either 2-tori or 2-spheres for every  $b$  with  $\zeta_k < b \leq 1/2$  and these bound either solid tori or 3-balls in the metric ball. In Case 1, the level sets are topological 2-spheres and they bound 3-balls in the metric ball.

Next, we shall show that in either case, provided that  $\epsilon > 0$  is sufficiently small, the level sets of the distance function from  $x'_{n(k)}$  must be 2-tori. Fix  $0 < \epsilon < \epsilon_1(\epsilon')$  such that Proposition 11.8 holds for these values of  $\epsilon$  and  $\epsilon'$ . Consider the annular region  $A_k = d(x'_{n(k)}, \cdot)^{-1}([1/4, 3/4])$ . This is a compact subset and if  $k$  is sufficiently large, then every point of this compact set is within  $\hat{\epsilon}$  of a point of  $B(\bar{x}_k, 1)$  at which  $B(\bar{x}_k, 1)$  is interior  $\mu$ -flat of some fixed scale  $s$ , depending only on  $a$ . Provided that  $\mu$  is sufficiently small, and having taking  $\hat{\epsilon}$  sufficiently small, depending on  $\mu$  and  $a$ , by Lemma 11.1 every point of  $A_k$  is the center of an  $S^1$ -product neighborhood with  $\epsilon$ -control and by Proposition 11.8 there is an open subset  $U_{n(k)} \subset M_{n(k)}$  containing  $A_k$  that is the total space of a circle fibration where the fibers of the fibration make angle at most  $\epsilon'$  with the horizontal spaces of the  $S^1$ -product neighborhoods with  $\epsilon$ -control at every point of  $A_k$ . Of course, there is a compact subsurface  $\Sigma_k$  contained in the base of the fibration with the property that the pre-image,  $W_k$ , of  $\Sigma_k$  contains  $A_k$ . Each component of  $\partial W_k$  is a torus. For every  $b \in (1/4, 1/2)$  the level set  $d(x'_{n(k)}, \cdot)^{-1}(b)$  separates two boundary components of  $W_k$ . Since a 2-sphere in the total space of a circle bundle cannot separate boundary components of that circle bundle, it follows that the level sets  $d(x'_{n(k)}, \cdot)^{-1}(b)$  are 2-tori.

This implies that for all  $k$  sufficiently large, Case 2 holds, and the soul of the limiting 3-manifold is a circle. Thus, for every  $k$  sufficiently large, for every  $0 < b \leq 1/2$  the pre-image  $d(x'_{n(k)}, \cdot)^{-1}([0, b])$  is a solid torus, denoted  $T_b$ . We fix  $b = 3/8$ . Of

course, provided that  $k$  is sufficiently large  $B(x_{n(k)}, 1/4) \subset T_b \subset B(x_{n(k)}, 1/2)$ . This gives a contradiction and completes the proof of the claims in the first paragraph of the statement.

Suppose that we have an open set  $\tilde{U}$  with an  $S^1$ -fibration as given in the second paragraph of the statement. Since the fibers of  $\tilde{U}$  are small, there is a saturated open subset  $V \subset \tilde{U}$  that contains the metric sphere at distance  $3/8$  from  $x$  and contained in  $A = B(x, 1/2) \setminus \overline{B(x, 1/4)}$ . A slightly smaller compact saturated subset  $V'$  also contains this metric sphere. The boundary components of  $V'$  are tori contained in  $A$ . Since  $V'$  separates the metric spheres at distance  $1/4$  and  $1/2$ , so does at least one of the boundary components of  $V'$ . This boundary component is then parallel to the metric sphere at distance  $1/4$  from  $x$  and hence bounds a solid torus in  $B(x, 1/2)$ .  $\square$

There is a further result that is not actually necessary for what follows but which makes the picture clearer and also simplifies somewhat several of the arguments.

**Proposition 11.22.** *Under the notation and hypothesis of the previous proposition, possibly after making the positive constants  $\mu_3(\epsilon, a)$  and  $\hat{\epsilon}_1(\epsilon, a, r_1, r_2)$  smaller, the  $S^1$ -factors in the local  $S^1$ -product structures with  $\epsilon$ -control contained in  $B_{\lambda^2 g_n}(x, 3r'/4) \setminus B_{\lambda^2 g_n}(x, r'/4)$  are homotopically non-trivial in  $B_{\lambda^2 g_n}(x, 3r'/4)$ .*

*Proof.* Let us suppose that the result does not hold. The previous argument shows that we may as well assume that  $r_2 = r_1$  and consider sequences  $\mu_k, \hat{\epsilon}_k$  tending to 0 and a sequence of counter examples  $B_k$  within  $\hat{\epsilon}_k$  of 2-dimensional balls  $B(\bar{x}_k, 1)$  which are interior  $\mu_k$ -flat on scale 1. The limit is a circular cone with cone angle  $\theta \leq 2\pi$ . The fundamental group  $\Gamma_k$  of  $B_k$  based at  $x_{n_k}$  is infinite cyclic and the shortest homotopically non-trivial loop through  $x_{n_k}$  has a length that tends to zero as  $k \rightarrow \infty$ . Thus, for any  $\epsilon > 0$  the number of elements in  $\pi_1(B_k, x_{n_k})$  represented by loops based at  $x_{n_k}$  of length  $< \epsilon$  goes to infinity as  $k \rightarrow \infty$ . Fix  $0 < d < \ell_0$ , where  $\ell_0$  is the constant from Proposition 9.36. Since the circular cone is interior flat at any point at distance  $d$  from the cone point on a scale depending only on  $d$  and the area of the cone is  $\geq a$ , the argument in the proof of the previous result shows that the following hold for all  $k$  sufficiently large. For any  $y \in Z$  with  $d(y, z) = d$  and for any  $y_{n_k} \in B_k$  within  $\hat{\epsilon}_k$  of  $y$ , the ball  $B(y_{n_k}, d/2)$  is contained in the total space  $V_k$  of an  $S^1$ -fibration over a disk with fibers isotopic to the  $S^1$ -factors in an  $S^1$ -product structure. The image of the fundamental group of  $V_k$  in  $\pi_1(B_k, x_{n_k})$  is then contained in the cyclic subgroup generated by a fiber of the  $S^1$ -product structures (all such fibers in all such  $S^1$ -product structures in  $B(x_{n_k}, 7r/8)$  are homotopic). But our assumption is that these fibers are homotopically trivial. This would imply that the image of the fundamental group of  $V_k$  is trivial. This contradicts Proposition 9.36.  $\square$

The topological import of this result about the fundamental group is the following:

**Corollary 11.23.** *Under the notation and hypotheses of the second paragraph of Proposition 11.20, the  $S^1$ -fibration structure on  $\tilde{U}$  extends to a Seifert fibration over  $\tilde{U} \cup B_{\lambda^2 g_n}(x, r'/2)$  with at most one singular fiber.*

**Definition 11.24.**  $B_{\lambda^2 g_n}(x, r'/4)$  satisfying the conclusions of Propositions 11.20 and 11.22 and Corollary 11.23 is an  $\epsilon'$ -solid torus neighborhood near a 2-dimensional interior cone point, or an  $\epsilon'$ -solid torus neighborhood for short.

**Remark 11.25.** In fact, a strengthening of this argument (see Theorem 0.2 and the material in Section 4 of [32]) proves that the order of the exceptional fiber is bounded above by  $2\pi/\alpha$  where  $\alpha$  is the cone angle of the nearby interior  $\mu$ -good ball at its central point. We shall not make use of this result.

### 11.5 Near almost flat boundary points

Now let us turn to the parts of the  $M_n$  close to flat boundary points of a 2-dimensional Alexandrov ball. First we need a result that tells us that as we pass from one 3-dimensional ball to another the points close to boundary points of close 2-dimensional balls don't change too much.

**Lemma 11.26.** *Given  $0 < d < (0.1)$  and  $a > 0$ , there is a positive constant  $\hat{\epsilon}'_0(d, a)$  such that the following hold for all  $0 < \hat{\epsilon} \leq \hat{\epsilon}'_0(d, a)$ . Suppose that  $x, x', y \in M_n$  and  $B_{\lambda^2 g_n}(x, 1)$  and  $B_{(\lambda')^2 g_n}(x', 1)$  are within  $\hat{\epsilon}$  in the Gromov-Hausdorff distance of 2-dimensional Alexandrov balls  $B(\bar{x}, 1)$  and  $B(\bar{x}', 1)$ , respectively, of curvature  $\geq -1$  and area  $\geq a$ . Suppose that  $y \in B_{\lambda^2 g_n}(x, 1/3) \cap B_{(\lambda')^2 g_n}(x', 1/3)$ . Suppose that  $(1/2)^{-1} \leq \lambda/\lambda' \leq 2$ . Suppose that, viewing  $y$  as a point of  $B_{\lambda^2 g_n}(x, 1)$ , it is within  $\hat{\epsilon}$  of a point  $\bar{y} \in \partial B(\bar{x}, 1)$ . Then there is  $z \in B_{(\lambda')^2 g_n}(x', 1)$  with  $d_{(\lambda')^2 g_n}(y, z) < d$  and with  $z$  being within  $\hat{\epsilon}$  of a point  $\bar{z} \in \partial B(\bar{x}', 1)$ .*

*Proof.* We set  $R = \lambda/\lambda'$ . Let us first consider the case when  $1 \leq R \leq 2$ . Then  $R \cdot B_{(\lambda')^2 g_n}(y, R^{-1}/2) = B_{\lambda^2 g_n}(y, 1/2)$ . Let  $\bar{y}' \in B(\bar{x}', 1)$  be a point within  $\hat{\epsilon}$  of  $y$ , when the latter is viewed as a point of  $B_{(\lambda')^2 g_n}(x', 1)$ . We consider the balls  $R \cdot B(\bar{y}', R^{-1}/2)$  and  $B(\bar{y}, 1/2)$ . By Lemma 9.22 the first is within  $4R\hat{\epsilon}$  of  $R \cdot B_{(\lambda')^2 g_n}(y, R^{-1}/2)$ , and the second is within  $4\hat{\epsilon}$  of  $B_{\lambda^2 g_n}(y, 1/2)$ . Since these latter two balls are equal, we see that  $R \cdot B(\bar{y}', R^{-1}/2)$  and  $B(\bar{y}, 1/2)$  are within  $(4R + 4)\hat{\epsilon}$  of each other in the Gromov-Hausdorff distance. Also, it is clear that for each of these two balls the curvature is bounded below by  $-1$  and the area is each bounded below by a positive constant depending only on  $a$ . Lastly, by construction  $\bar{y} \in \partial B(\bar{y}, 1/2)$ . By Lemma 10.7, if, given  $d$  and  $a$ ,  $\hat{\epsilon}$  is sufficiently small, then  $\bar{y}'$  is within distance  $d/2$  of a point  $\bar{w} \in \partial[R \cdot B(\bar{y}', R^{-1}/2)]$ . Let  $w \in B_{(\lambda')^2 g_n}(x', 1)$  be within distance  $\hat{\epsilon}$  of  $\bar{w}$ . By the triangle inequality,  $d_{(\lambda')^2 g_n}(y, w) < d/2 + (1 + R)\hat{\epsilon}$ , and the right-hand side is less than  $d$  if  $\hat{\epsilon}$  is sufficiently small. This establishes the result when  $R \geq 1$ . The other case is symmetric.  $\square$

Now we are ready to study the local structure of points near to 2-dimensional boundary points, see FIG. 6.

**Proposition 11.27.** *Fix  $\epsilon' > 0$  and  $0 < \epsilon < \epsilon_0(\epsilon')$ . Then there is a positive constant  $\xi_1(\epsilon) \leq \xi_0$  such that for every  $0 < \xi < \xi_1(\epsilon)$  there is a positive constant  $\bar{\mu}(\xi)$  such that for any  $0 < \mu < \bar{\mu}(\xi)$  and for any  $0 < s_1 \leq \alpha_0(\xi)$ , where  $\alpha_0(\xi)$  is the constant defined just before Proposition 10.29, and for all  $\hat{\epsilon} > 0$  less than a positive constant*



$\hat{e}_2(\epsilon, \xi, s_1)$ , the following hold. Suppose that, for some  $n$ , there are a point  $x \in M_n$  and a constant  $\lambda \geq \rho^{-1}(x)$  with the property that  $B_{\lambda^2 g_n}(x, 1)$  is within  $\hat{e}$  of a 2-dimensional Alexandrov ball  $X = B(\bar{x}, 1)$  of curvature  $\geq -1$ . Suppose that  $\gamma$  is a  $\mu$ -approximation to  $\partial X \cap B(\bar{x}, 3/4)$  on scale  $s_1$ . Suppose that  $\tilde{\gamma}$  is a geodesic in  $B_{\lambda^2 g_n}(x, 1)$  whose endpoints are within  $\hat{e}$  of those of  $\gamma$ . Then:

1. The  $\xi$ -box  $\bar{\nu}_\xi(\tilde{\gamma})$  is homeomorphic to  $D^2 \times [0, 1]$  where the disks in this (topological) product structure are the level sets of  $f_{\tilde{\gamma}}$ . The complement of its core, denoted  $\nu_\xi^0(\tilde{\gamma})$ , is homeomorphic to  $S^1 \times (0, 1) \times [0, 1]$  where each circle factor is the intersection of a level set of  $f_{\tilde{\gamma}}$  with a level set of  $h_{\tilde{\gamma}}$ . (These intersections are called level circles.)
2. Each point of  $\nu_\xi^0(\tilde{\gamma})$  is the center of an  $S^1$ -product neighborhood with  $\epsilon$ -control.
3. For any  $q \in \nu_\xi^0(\tilde{\gamma})$  and any  $S^1$ -product neighborhood with  $\epsilon$ -control containing  $q$ , the angle (in the sense given in the footnote to Proposition 11.4) at  $q$  between the level circle  $S(q) = F^{-1}(F(q))$  through  $q$  and the horizontal space of the  $S^1$ -product neighborhood is within  $\epsilon'$  of  $\pi/2$ . Furthermore, if  $q$  is contained in  $\varphi(S^1 \times B(0, \epsilon^{-1}/2))$ , then  $S(q)$  is isotopic in the  $S^1$ -product neighborhood to an  $S^1$ -factor.

*Proof.* We fix positive constants  $\epsilon'$ ,  $\epsilon < \epsilon_0(\epsilon')$ ,  $\xi < \xi_0$ , and  $s_1 \leq \alpha_0(\xi)$ . Rescaling has no effect on  $\xi$  nor on  $\mu$  and scales  $\hat{e}$  linearly. Thus, as before, we can assume that  $s_1$  is fixed throughout the argument. We denote the endpoints of  $\gamma$  by  $e_\pm$  and those of  $\tilde{\gamma}$  by  $\tilde{e}_\pm$ . We work with the metric  $\lambda^2 g_n$ , so that in particular,  $\ell(\tilde{\gamma})$  means the length of  $\tilde{\gamma}$  with respect to this metric.

If the endpoints of  $\tilde{\gamma}$  are sufficiently close to those of  $\gamma$ , then  $\tilde{\gamma}$  is close to a geodesic in  $B(\bar{x}, 1)$  with the same endpoints as  $\gamma$ . This geodesic is also a  $\mu$ -approximation to  $\partial X$  and we can simply replace  $\gamma$  by this geodesic. This allows us to assume the following: for any fixed  $\beta > 0$  we can choose  $\hat{e} > 0$  sufficiently small so that  $\tilde{\gamma}$  is within  $\beta$  of  $\gamma$ .

Provided that  $\xi > 0$  is sufficiently small,  $\mu$  is sufficiently small, and  $\hat{e}$  is sufficiently small, it follows from Lemma 10.24 that  $f_{\tilde{\gamma}}$  is regular on  $\bar{\nu}_\xi(\tilde{\gamma})$ . Hence, each level set  $L$  of  $f_{\tilde{\gamma}}$  is a Lipschitz surface and these level surfaces foliate  $\bar{\nu}_\xi(\tilde{\gamma})$ . It also follows from this lemma that for any  $y \in \nu_\xi^0(\tilde{\gamma})$  there is a point  $z$  such that the set  $\{((e_+)', (e_-)', \gamma', z')\}$  is a  $(2, 10\xi)$ -strainer of size  $\xi^2$  at  $y$ . Hence, by Proposition 11.4 provided that  $\xi$  is sufficiently small given  $\epsilon'$  and  $\epsilon$  and provided that  $\hat{e}$  is sufficiently small given  $\epsilon', \epsilon, \xi, s_1$ , we have:

1. every point of  $\nu_\xi^0(\tilde{\gamma})$  is the center of an  $S^1$ -product structure with  $\epsilon$ -control, and
2. the map  $F = (f_{\tilde{\gamma}}, h_{\tilde{\gamma}})$  determines a fibration of  $\nu_\xi^0(\tilde{\gamma})$  with fibers that are circles almost  $\epsilon'$ -orthogonal to the horizontal spaces of the  $S^1$ -product structures at the various points. Hence, for  $\xi$  and  $\mu$  sufficiently small, depending only on  $\epsilon$ , and for  $\hat{e}$  sufficiently small,  $\nu_\xi^0(\tilde{\gamma})$  is homeomorphic to  $S^1 \times (0, 1) \times [0, 1]$  where the circle-factors are the level circles of  $F$ .

This establishes the second and third parts of the proposition. We turn to the first part. We shall show that provided that  $\xi$  is sufficiently small given  $\epsilon$ , provided that  $\mu$  is sufficiently small, and provided that  $\hat{\epsilon}$  is sufficiently small given  $\xi, \epsilon$ , the level sets of  $f_{\tilde{\gamma}}$  are homeomorphic to disks. From the immediately preceding discussion, it follows that the boundary of any level surface for  $f_{\tilde{\gamma}}$  is a single circle. Since the level sets of  $f_{\tilde{\gamma}}$  are connected, to show these level sets are homeomorphic to disks it suffices to show that they have virtually abelian fundamental groups and are orientable. The level sets are orientable since  $M_n$  is and since they are the level sets of a regular Lipschitz function so that there is a neighborhood of the level set in  $M_n$  that is homeomorphic to the product of the level set with  $I$ . Thus, the first part of the result is completed by showing the following:

**Claim 11.28.** *For  $\xi$  sufficiently small, for  $\mu$  sufficiently small, given  $\xi$ , and for  $\hat{\epsilon} > 0$  sufficiently small, the fundamental groups of the level sets of  $f_{\tilde{\gamma}}$  are virtually abelian.*

*Proof.* We suppose that the claim does not hold. Then there are sequences of  $\xi_k \rightarrow 0$ ,  $\mu_k$  tending to zero sufficiently small so that Lemma 10.24 holds for  $\xi_k$ , and  $\hat{\epsilon}_k \rightarrow 0$  and counter-examples  $\nu_{\xi_k}(\tilde{\gamma}_k)$  with the fundamental groups of the level sets of  $f_{\tilde{\gamma}_k}$  not virtually abelian. Take as base points  $p_k$  the midpoints of  $\tilde{\gamma}_k$ . Notice that for all  $k$  sufficiently large, since the length  $\ell(\gamma_k)$  of the boundary approximating geodesic  $\gamma_k$  in the 2-dimensional ball is at least  $s_1/50$ , by Part 5 of Lemma 10.24, we have that  $B(p_k, \xi s_1/200)$  is contained  $\nu_{\xi}(\tilde{\gamma}_k)$ . Also, notice that the map  $\pi_1(L_k, p_k) \rightarrow \pi_1(\nu_{\xi}(\tilde{\gamma}_k), p_k)$  induced by the inclusion is an isomorphism. We denote by  $\ell' = \ell_0 \xi s_1/200$ , where  $\ell_0$  is the constant from Proposition 9.36.

The above argument shows that given  $0 < t \leq \xi$ , for every  $k$  sufficiently large  $\nu_{\xi_k}(\tilde{\gamma}_k) \setminus \nu_{t\xi}(\tilde{\gamma}_k)$  is homeomorphic to  $S^1 \times I \times [t\xi\ell(\gamma_k), \xi\ell(\gamma_k)]$  where the circle factors are the level circles of  $F_k = (f_{\tilde{\gamma}_k}, h_{\tilde{\gamma}_k})$ . It follows that for all  $k$  sufficiently large there is a point  $y_k \in B(p_k, \ell')$  and an open set  $U_k$  which is the total space of a  $S^1$ -bundle over a disk with  $B(y_k, \ell'/4) \subset U_k \subset B(y_k, \ell'/2)$ , and hence  $\pi_1(U_k, y_k) \cong \mathbb{Z}$ . It also follows that for all  $k$  sufficiently large  $L_k \cap (\nu_{\xi_k}(\tilde{\gamma}_k) \setminus \nu_{t\xi}(\tilde{\gamma}_k))$  is an annulus and hence, denoting  $L_k(t\xi)$  by the intersection  $L_k(t\xi) = L_k \cap \nu_{t\xi}(\tilde{\gamma}_k)$ , the inclusion map induces an isomorphism of fundamental groups  $\pi_1(L_k(t\xi), p_k) \xrightarrow{\cong} \pi_1(\nu_{\xi}(\tilde{\gamma}_k), p_k)$ . Hence, the inclusion induces a injection  $\pi_1(L_k(t\xi), p_k) \rightarrow \pi_1(B(p_k, \xi s_1/200), p_k)$ .

It follows from Part 4 of Lemma 10.24 that the diameter of  $L_k(t\xi)$  is at most  $4s_1(1 + 2\xi)t\xi$ , and hence  $\pi_1(L_k(t\xi), p_k)$  is generated by elements  $\{c_1, \dots, c_{r(k)}\}$  represented by loops of length at most  $8s_1(1 + 2\xi)t\xi$  based at  $p_k$ . If  $\pi_1(L_k(t\xi), p_k) = \pi_1(L_k, p_k)$  is not virtually abelian then, since it is the fundamental group of a non-compact surface, it is a free group of rank at least two. Hence, at least one of the  $c_i$ , let's call it  $c_1$ , has the property that no power of the image of  $c_1$  in  $\pi_1(B(p_k, \xi s_1/200), p_k)$  is contained in the image of  $\pi_1(U_k, y_k) \rightarrow \pi_1(B(p_k, \xi s_1/200), p_k)$ . If  $t$  is sufficiently small, then contradicts Proposition 9.36, as we see by rescaling by  $200/s_1\xi$ .  $\square$

The claim establishes the first part of the proposition and hence completes the proof of the proposition.  $\square$

The above arguments show that in fact the  $S^1$ -product neighborhoods in this result can be chosen in the following way.

**Addendum 11.29.** Under the hypothesis and notation of the previous proposition, possibly after making the positive constants  $\bar{\mu}(\xi)$  and  $\hat{\epsilon}_2(\epsilon, \xi, s_1)$  smaller the following holds. For any point  $x \in \nu_\xi^0(\tilde{\gamma})$  and any the  $S^1$ -product structure with  $\epsilon$ -control centered at  $x$ ,  $\varphi: S^1 \times B(0, \epsilon^{-1}) \rightarrow M_n$ , the Euclidean coordinates on  $\mathbb{R}^2$  can be chosen so that the following hold for any point  $q \in \varphi(S^1 \times B(0, \epsilon^{-1}/2))$ :

1. For any geodesic  $\zeta$  from  $\tilde{\gamma}$  to  $q$ ,  $\varphi^{-1}$  of intersection of  $\zeta$  with the  $S^1$ -product neighborhood is within  $\epsilon'$  of the straight line starting at  $q$  in the negative  $y$ -direction in the horizontal  $B(0, \epsilon^{-1})$ .
2. For any geodesics  $\zeta_\pm$  from  $e_\pm(\tilde{\gamma})$  to  $q$ ,  $\varphi^{-1}$  of the intersections of  $\zeta_\pm$  with the  $S^1$ -product neighborhood are within  $\epsilon'$  of horizontal straight lines in  $B(0, \epsilon^{-1})$  starting at  $q$  in the  $x_\pm$ -directions.

**Definition 11.30.** We call any neighborhood  $\nu_\xi(\tilde{\gamma})$  for which there is a geodesic  $\gamma$  in a 2-dimensional standard ball satisfying the hypotheses of Proposition 11.27 (and hence  $\nu_\xi(\tilde{\gamma})$  satisfies the conclusions of the last two results) an  $\epsilon$ -solid cylinder neighborhood at scale  $s_1$  near a flat boundary, or simply an  $\epsilon$ -solid cylinder neighborhood at scale  $s_1$  for short.

**Lemma 11.31.** For any  $0 < \xi < \xi_0$  and any  $0 < s_1 \leq \alpha_0(\xi)$  there are positive constants  $\hat{\epsilon}_3(\epsilon, \xi, s_1) \leq \hat{\epsilon}(\epsilon, \xi, s_1)$  and  $\mu_4(\xi) \leq \bar{\mu}(\xi)$  such that following hold for  $0 < \mu < \mu_4(\xi)$  and  $0 < \hat{\epsilon} < \hat{\epsilon}_3(\epsilon, \xi, s_1)$ . With notation and assumptions as in the previous proposition, let  $\tilde{e}_\pm$  be the endpoints of  $\tilde{\gamma}$ .

1. For any  $y \in \nu_\xi(\tilde{\gamma})$ , we have  $\tilde{Z}\tilde{e}_-y\tilde{e}_+ > \pi - 8\xi$ .
2. For each  $y \in \nu_\xi^0(\tilde{\gamma})$  there are points  $z, w$  at distance  $\ell(\tilde{\gamma})/8$  from  $y$  such that for any minimal length geodesic  $\alpha$  from  $\tilde{\gamma}$  to  $y$ , denoting by  $a$  the intersection  $\alpha \cap \tilde{\gamma}$  we have that  $\tilde{Z}ayz, \tilde{Z}zyw$  are each greater than  $\pi/2 - 2\xi^2$  and  $\tilde{Z}ayw > \pi - 2\xi^2$ . Lastly,  $\tilde{Z}\tilde{e}_-yz > \pi - 6\xi$ .
3. For any  $c \in [\xi^2, \xi]$  and for any level surface  $L$  of  $f_{\tilde{\gamma}}$  the distance from any point of  $L \cap h_{\tilde{\gamma}}^{-1}(c \cdot \ell(\tilde{\gamma}))$  to  $L \cap \tilde{\gamma}$  is at most  $(1 + 4\xi)c \cdot \ell(\tilde{\gamma})$ .
4.  $\nu_{\xi^2}(\tilde{\gamma})$  contains  $B(y, \xi^2 \ell(\tilde{\gamma})/10)$  about the intersection of the center of  $\nu_\xi(\tilde{\gamma})$  with  $\tilde{\gamma}$ .
5. The geodesic  $\tilde{\gamma}$  is within  $\xi^2 \ell(\tilde{\gamma})/100$  of the arc on  $\partial B(\bar{x}, 1)$  with the same endpoints as  $\gamma$ .

(Here all distances and  $\ell(\tilde{\gamma})$  are measured with respect to  $\lambda^2 g_n$ .)

*Proof.* The first four items are a direct consequence of Lemma 10.24 and a standard limiting argument. Let us consider the last statement. If it is false then we have a sequence  $\mu_n \rightarrow 0$  and for each  $n$  a sequence  $\hat{\epsilon}_{n,m}$  tending to zero as  $m \rightarrow \infty$  and

counter-examples  $\nu_{n,m}$  with generating geodesics  $\tilde{\gamma}_{n,m}$  whose endpoints are within  $\hat{\epsilon}_{n,m}$  of those of  $\gamma_{n,m}$  which is a  $\mu_n$ -approximation to  $\partial B(\bar{y}_{n,m}, s_1(n))$  on scale  $s_1(n)$ . This means that  $B(\bar{x}_{n,m}, 1)$  is boundary  $\mu_n$ -flat near  $\bar{y}_{n,m} \in \partial B(\bar{x}_{n,m}, 1/3)$  on all scales  $\leq s_1(n)$  and  $\gamma_{n,m} \subset B(\bar{y}_{n,m}, s_1(n))$  has endpoints in  $\partial B(\bar{y}_{n,m}, 7s_1(n)/8)$ . For each  $n$ , since  $\hat{\epsilon}_{n,m} \rightarrow 0$  as  $m \rightarrow \infty$ , passing to a subsequence in  $m$  we can assume that the  $B(\bar{y}_{n,m}, s_1(n))$  converge to  $B(\bar{y}_{n,\infty}, s_1(n))$  and that the  $\tilde{\gamma}_{n,m}$  converge to a geodesic  $\gamma_{n,\infty}$  in  $B(\bar{y}_{n,\infty}, s_1(n))$  of length at least  $s_1(n)/100$ . By Corollary 10.6 the endpoints of  $\gamma_{n,\infty}$  are contained in  $\partial B(\bar{y}_{n,\infty}, s_1(n))$  and indeed are the limits of the endpoints of the  $\gamma_{n,m}$ . Now we consider  $(1/s_1(n))B(\bar{y}_{n,\infty}, s_1(n))$ . Since the  $\mu_n$  tend to zero, these unit balls converge to the unit ball in half-space centered around a boundary point, and passing to a subsequence in  $n$  we can assume that the  $\gamma_{n,\infty}$  converge to a geodesic  $\gamma_{\infty,\infty}$ . Since the limit is a ball in flat half-space it follows that  $\gamma_{\infty,\infty}$  is contained in the boundary. Similarly, the arcs  $(1/s_1(n))\alpha_{n,\infty}$  in  $(1/s_1(n))\partial B(\bar{y}_{n,\infty}, s_1(n))$  converge to  $\gamma_{\infty,\infty}$ . Thus, for each  $n$  we can choose  $m(n)$  such that both the  $\tilde{\gamma}_{n,m(n)}$  and the  $\alpha_{n,m(n)}$  converge to  $\gamma_{\infty,\infty}$ . Since the length of  $\tilde{\gamma}_{n,m(n)}$  is at least  $s_1(n)/100$ . This shows that the 5<sup>th</sup> condition holds for all  $(n, m(n))$  for all  $n$  sufficiently large, which is a contradiction.  $\square$

We shall also need smooth vector fields well-adapted to  $\nu_\xi(\tilde{\gamma})$ .

**Proposition 11.32.** *Again with the notation and assumptions of Proposition 11.27 there is a smooth unit vector field  $\chi$  on  $\nu_\xi(\tilde{\gamma})$  such that, setting  $d_\pm$  equal to the distance function from the endpoints  $\tilde{e}_\pm$  of  $\tilde{\gamma}$ , we have  $d'_-(\chi) > 1 - 36\xi$ ,  $d'_+(\chi) < -1 + 44\xi$ . Furthermore, on  $\nu_\xi(\tilde{\gamma}) \setminus \nu_{2\xi^2}(\tilde{\gamma})$  we have  $|h'_\chi(\chi)| < 11\xi^2$ . Since  $\xi < 10^{-3}$ , for any points  $p, q$  on a flow line of the flow generated by  $\chi$ , with  $p \in \nu_{3\xi/4}(\tilde{\gamma}) \setminus \nu_{2\xi^2}(\tilde{\gamma})$ , we have*

$$\left| \frac{h_{\tilde{\gamma}}(p) - h_{\tilde{\gamma}}(q)}{f_{\tilde{\gamma}}(p) - f_{\tilde{\gamma}}(q)} \right| < 12\xi^2.$$

*In particular, any maximal flow line of  $\chi$  that meets  $h_{\tilde{\gamma}}^{-1}[0, 3\xi/4]$  is a closed interval with endpoints in the ends of  $\nu_\xi(\tilde{\gamma})$  and this interval meets each level set of  $f_{\tilde{\gamma}}$  in a single point.*

*Proof.* We consider the subset  $V$  of the unit tangent bundle of  $\nu_\xi^0(\tilde{\gamma})$  consisting of all unit tangent vectors  $\tau$  at points  $y$  for which the following hold:

1. The distance between  $\tilde{\gamma}'_y$  and  $\tau$  in  $S_y(M_n)$  is greater than  $\pi/2 - 2\xi^2$ ,
2. The distance between  $(d_-)'_y$  and  $\tau$  is greater than  $\pi - 6\xi$ .
3. There exists a point  $w$  at distance  $\ell(\tilde{\gamma})/8$  from  $y$  such that (i) the distance in the tangent sphere at  $y$ ,  $S_y$ , from  $\tilde{\gamma}'_y$  to  $w'_y$  is greater than  $\pi - 2\xi^2$  and (ii) the distance in  $S_y$  from  $w'_y$  to  $\tau$  is greater than  $\pi/2 - 2\xi^2$ .

By Lemma 9.5 the subset  $V$  of the unit tangent bundle of  $\nu_\xi^0(\tilde{\gamma})$  is open. By the previous proposition its image under the projection mapping is all of  $y \in \nu_\xi^0(\tilde{\gamma})$ . Because of the third item above, for any  $y \in \nu_\xi^0(\tilde{\gamma})$  there are antipodes  $n, s$  of the tangent sphere  $S_y(M_n)$  to  $M_n$  at  $y$  such that the directions of geodesics from  $y$  to

$\tilde{\gamma}$  lie in the ball of radius  $2\xi^2$  about  $n$  and the directions of geodesics from  $y$  to the point  $w$  lie in the ball of radius  $4\xi^2$  about  $s$ . It follows that the intersection of  $V$  with  $S_y(M_n)$  is contained in the collar  $C$  centered around the equator  $E$  determined by  $n$  and  $s$ , a collar that has width  $12\xi^2$ . Furthermore, there is a point  $e$  on  $E$  whose  $23\xi$ -neighborhood contains  $\tilde{e}'_-$ , so that all points of  $V \cap S_y(M_n)$  are contained in the  $13\xi$  ball, denoted  $D$ , about the antipode of  $e$ .

Unfortunately, the subsets  $V \cap S_y(M_n)$  are not convex. We remedy this defect by replacing  $V$  by  $\hat{V}$  which is obtained by taking fiber-wise (geodesic) convex hull of  $V$ . The latter is an open subset of the unit tangent sphere bundle with convex fibers. It is an easy exercise in 2-dimensional spherical trigonometry to show that, since  $\xi < \xi_0 \leq 10^{-3}$ , every tangent vector  $\tau'$  in  $\hat{V} \cap S_y(M_n)$  lies in  $\hat{C} \cap D$  where  $\hat{C}$  is the collar of width  $14\xi^2$  centered around  $E$ . Thus, every tangent vector  $\tau'$  in  $\hat{V} \cap S_y(M_n)$  satisfies the three conditions above with  $2\xi^2$  replaced by  $11\xi^2$  in Condition 1 and in Part (ii) of Condition 3 and  $6\xi$  replaced by  $36\xi$  in Condition 2. We then have that the distance between  $(\tilde{e}_+)'_y$  and  $\tau$  is less than  $44\xi$ .

It then follows that for every  $\tau \in \hat{V}$  we have  $-11\xi^2 < h'_{\tilde{\gamma}}(\tau) < 11\xi^2$  and  $f'_{\tilde{\gamma}}(\tau) > 1 - 40\xi$ . Since  $\hat{V}$  has non-empty convex fibers over every  $y \in \nu_{\xi}^0(\tilde{\gamma})$ , there is a smooth vector field  $\chi$  defined on all of  $\nu_{\xi}^0(\tilde{\gamma})$  lying in  $\hat{V}$ .

It follows immediately from these inequalities that if  $y, z$  lie on the same flow line of  $\chi$  then

$$\frac{|h_{\tilde{\gamma}}(z) - h_{\tilde{\gamma}}(y)|}{|f_{\tilde{\gamma}}(z) - f_{\tilde{\gamma}}(y)|} < \frac{11\xi^2}{1 - 40\xi} < 12\xi^2.$$

Since  $\xi < 10^{-3}$ , if  $\xi\ell(\tilde{\gamma})/2 \leq h_{\tilde{\gamma}}(y) \leq 3\ell(\tilde{\gamma})\xi/4$ , then for any point  $z$  on the flow line through  $y$  we have  $\xi\ell(\tilde{\gamma})/4 < h_{\tilde{\gamma}}(z) < 7\xi\ell(\tilde{\gamma})/8$ .

This defines a vector field as required on  $\nu_{\xi}^0(\tilde{\gamma})$ . On  $\nu_{2\xi^2}(\tilde{\gamma})$  there is a smooth unit vector field  $\chi'$  with the property that  $d'_-(\chi) > 1 - 36\xi$ ,  $d'_+(\chi) < -1 + 44\xi$ . Patching these together with a partition of unity completes the construction of the vector field as required.  $\square$

**Definition 11.33.** The metric  $\lambda^2 g_n$  that was used in the previous proposition is called the *metric used to define the neighborhood*  $\nu_{\xi}(\tilde{\gamma})$ . By  $\ell(\tilde{\gamma})$  we always mean the length of the geodesic  $\tilde{\gamma}$  with respect to the metric used to define the neighborhood. By a *spanning disk* in an  $\epsilon$ -solid cylinder we mean a 2-disk with boundary contained in the side of the solid cylinder that separates the ends of the solid cylinder.

### 11.5.1 Intersections of the $\nu_{\xi}(\tilde{\gamma})$

It is important to have control over the intersections of the various  $\epsilon$ -solid cylinder neighborhoods near a flat boundary, see FIG. 7.

**Lemma 11.34.** *Given  $0 < \xi < \xi_0$  there is a positive constant  $\mu_5(\xi)$  such that for every  $\mu < \mu_5(\xi)$  and, given  $0 < s_1 \leq \alpha_0(\xi)$ , there is a positive constant  $\hat{\epsilon}_4(\xi, \mu, s_1)$  such that the following hold for all  $\hat{\epsilon} < \hat{\epsilon}_4(\xi, \mu, s_1)$ . For some  $n$ , let  $B_1 = B_{g'_n(x_1)}(x_1, 1) \subset M_n$  and  $B_2 = B_{g'_n(x_2)}(x_2, 1) \subset M_n$  be within  $\hat{\epsilon}$  of 2-dimensional Alexandrov balls  $\bar{B}_1 = B(\bar{x}_1, 1)$  and  $\bar{B}_2 = B(\bar{x}_2, 1)$  be of curvature  $\geq -1$  and area  $a$ . Suppose that*

$\bar{y}_i \in \partial B(\bar{x}_i, 1)$  with  $d(\bar{x}_i, \bar{y}_i) < \xi^2 s_1/100$ . Suppose that  $\bar{B}_i$  is boundary  $\mu$ -flat at  $\bar{y}_i$  on scale  $s_1$  and let  $\bar{\gamma}_i \subset B(\bar{y}_i, s_1/3)$  be  $\mu$ -approximations to  $\partial \bar{B}_i$  of length  $s_1/4$ . Suppose that  $\tilde{\gamma}_i \subset B_i$  is a geodesic whose endpoints are within  $\hat{\epsilon}$  of those of  $\bar{\gamma}_i$ . Denote  $\nu_\xi(\tilde{\gamma}_i)$  by  $\nu_i$  and suppose that  $\nu_1 \cap \nu_2 \neq \emptyset$ . Then there are arcs  $\tilde{\alpha}_i \subset \tilde{\gamma}_i$  such that the following hold:

1. For  $i = 1, 2$  the length of  $\tilde{\alpha}_i$  is at least  $\min(\ell(\tilde{\gamma}_1), \ell(\tilde{\gamma}_2))/5$ .
2. Either for each  $i = 1, 2$ , the subgeodesic  $\tilde{\alpha}_i$  contains an endpoint of  $\tilde{\gamma}_i$  or one of the  $\tilde{\alpha}_i$  is equal to  $\tilde{\gamma}_i$ .
3. The geodesics  $\tilde{\alpha}_1$  and  $\tilde{\alpha}_2$  in  $M_n$  are within  $\xi^2 s_1/100$  of each other.

Furthermore,  $\tilde{\gamma}_2$  meets  $\nu_1$  and the intersection of  $\tilde{\gamma}_2$  with  $\nu_1$  is contained in  $\nu_{\xi^2}(\tilde{\gamma}_1)$  and  $\tilde{\gamma}_2$  meets each level set of  $f_{\tilde{\gamma}_1}$  in at most one point. In particular, for any  $c \geq \xi$  the intersection of  $\tilde{\gamma}_2$  with the boundary of  $\bar{\nu}_{c\xi}(\tilde{\gamma}_1)$  is contained in the ends of this neighborhood.

*Proof.* Suppose that the result does not hold for some  $\xi$  with  $0 < \xi < \xi_0$ . Then there is a sequence  $\mu_k \rightarrow 0$  and for each  $k$  a constant  $s_{1,k} \leq \alpha_0(\xi)$ , a sequence  $\hat{\epsilon}_{k,m} \rightarrow 0$  as  $m \rightarrow \infty$ , and counter-examples to the result for these constants. For every  $k$  and  $m$  and for  $i = 1, 2$  we denote the various constituents of counter-examples as follows. Let  $B_{k,m,i} = B_{g'_{n(k,m)}(x_{k,m,i})}(x_{k,m,i}, 1) \subset M_{n(k,m)}$  be the 3-dimensional balls and  $\bar{B}_{k,m,i} = B(\bar{x}_{k,m,i}, 1)$  be the 2-dimensional balls and  $\bar{y}_{k,m,i} \in \partial \bar{B}_{k,m,i}$  the control points and  $\bar{\gamma}_{k,m,i} \subset \bar{B}_{k,m,i}$  the geodesics of length  $s_{1,k}/4$ , and  $\tilde{\gamma}_{k,m,i} \subset B(\bar{y}_{k,m,i}, s_{1,k}/3) \subset B_{k,m,i}$  the geodesics whose endpoints are within  $\hat{\epsilon}_{k,m}$  of those of  $\bar{\gamma}_{k,m,i}$ .

We take points  $y_{k,m,i} \in B_{k,m,i}$  within  $\hat{\epsilon}_{k,m}$  of  $\bar{y}_{k,m,i}$ . First notice that  $\tilde{\gamma}_{k,m,i}$  and  $\nu_\xi(\tilde{\gamma}_{k,m,i})$  are both contained in

$$B_{g'_{n(k,m)}(x_{k,m,i})}(y_{k,m,i}, s_{1,k}/3) \subset B_{g'_{n(k,m)}(x_{k,m,i})}(x_{k,m,i}, 10^{-5}).$$

Since these  $\epsilon$ -solid cylinders intersect, we have

$$B_{g'_{n(k,m)}(x_{k,m,1})}(y_{k,m,1}, 10^{-5}) \cap B_{g'_{n(k,m)}(x_{k,m,2})}(y_{k,m,2}, 10^{-5}) \neq \emptyset.$$

By Lemma 6.1 this implies that  $g'_{n(k,m)}(x_{k,m,1}) = R_{k,m}^2 g'_{n(k,m)}(x_{k,m,2})$  for some  $R_{k,m}$  with  $(0.99) < R_{k,m} < 1.01$ .

The ball  $B_{g'_{n(k,m)}(x_{k,m,1})}(y_{k,m,1}, s_{1,k}/3)$  contains  $\nu_\xi(\tilde{\gamma}_{k,m,1})$  and hence contains a point of  $\nu_\xi(\tilde{\gamma}_{k,m,2})$ . The length of  $\tilde{\gamma}_{k,m,2}$  with respect to  $g'_{n(k,m)}(x_{k,m,2})$  is  $s_{1,k}/4$ , so its length with respect to  $g'_{n(k,m)}(x_{k,m,1})$  is  $R_{k,m}s_{1,k}/4$  which is between  $(0.24)s_{1,k}$  and  $(0.26)s_{1,k}$ . It follows that  $\tilde{\gamma}_{k,m,2} \subset B_{g'_{n(k,m)}(x_{k,m,1})}(y_{k,m,1}, (0.6)s_{1,k})$ . Similarly,

$$R_{k,m} B_{g'_{n(k,m)}(x_{k,m,2})}(y_{k,m,2}, s_{1,k}/3) \subset B_{g'_{n(k,m)}(x_{k,m,1})}(y_{k,m,1}, s_{1,k}).$$

It follows that  $R_{k,m} B(\bar{y}_{k,m,2}, s_{1,k}/3)$  is within  $6\hat{\epsilon}_{k,m}$  of a sub-ball of radius between  $(0.32)s_{1,k}$  and  $(0.34)s_{1,k}$  in  $B(\bar{y}_{k,m,1}, s_{1,k})$ . Passing to a subsequence (in  $m$ ) so that

limits  $B(\bar{y}_{k,\infty,1}, s_{1,k})$  and  $B(\bar{y}_{k,\infty,2}, s_{1,k})$  exist and so that the  $R_{k,m}$  have a limit  $R_{k,\infty}$ , we see that  $R_{k,\infty}B(\bar{y}_{k,\infty,2}, s_{1,k}/3)$  is identified with a sub-ball of  $B(\bar{y}_{k,\infty,1}, s_{1,k})$ , and this identification is the limit as  $m \rightarrow \infty$  of the inclusions

$$R_{k,m}B_{g'_{n(k,m)}}(x_{k,m,2})(y_{k,m,2}, s_{1,k}/3) \subset B_{g'_{n(k,m)}}(x_{k,m,1})(y_{k,m,1}, s_{1,k}).$$

We can also assume that the  $\bar{\gamma}_{k,m,i}$  converge to geodesics  $\bar{\gamma}_{k,\infty,i}$ . Passing to a subsequence we can arrange that the  $\tilde{\gamma}_{k,m,i}$  also converge to geodesics  $\bar{\gamma}_{k,\infty,i}$  with the same endpoints as the  $\bar{\gamma}_{k,\infty,i}$ . Now we scale by  $1/s_{1,k}$  and take a limit of the  $(s_{1,k})^{-1}B(\bar{y}_{k,\infty,1}, s_{1,k})$  as  $k \rightarrow \infty$ . Since the  $\mu_k \rightarrow 0$ , both the two sub-balls  $(s_{1,k})^{-1}B(\bar{y}_{k,\infty,1}, s_{1,k}/3)$  and  $s_{1,k}^{-1}RB(\bar{y}_{k,\infty,2}, s_{1,k}/3)$  of  $(s_{1,k})^{-1}B(\bar{y}_{k,\infty,1}, s_{1,k})$  converge to balls that are isometric to unit balls centered around boundary points of  $[0, \infty) \times \mathbb{R}$ . We can assume that the  $\bar{\gamma}_{k,\infty,i}$  converge to geodesics  $\bar{\gamma}_{\infty,\infty,i}$  and that the  $\tilde{\gamma}'_{k,\infty,i}$  converge to geodesics  $\bar{\gamma}'_{\infty,\infty,i}$ . These limiting geodesics are geodesics in the boundary so that  $\bar{\gamma}_{\infty,\infty,i} = \bar{\gamma}'_{\infty,\infty,i}$ . Furthermore, the intersection  $\gamma_{\infty,\infty,1} \cap \gamma_{\infty,\infty,2}$  is a sub-geodesic of each and either shares an endpoint with each of these geodesics or is equal to one of them. Notice also that  $\nu_\xi(\gamma_{\infty,\infty,1}) \cap \nu_\xi(\gamma_{\infty,\infty,2}) \neq \emptyset$ . Thus,  $\gamma_{\infty,\infty,1} \cap \gamma_{\infty,\infty,2}$  has length greater than (0.24). Now construct arcs  $\tilde{\alpha}_{k,m,i} \subset \tilde{\gamma}_{k,m,i}$  converging to the  $\alpha_i$  and with the property that any time an endpoint of  $\alpha_i$  is equal to an endpoint of  $\bar{\gamma}_{\infty,\infty,i}$  the corresponding endpoint of  $\alpha_{k,m,i}$  is equal to an endpoint of  $\tilde{\gamma}_{k,m,i}$ . For all  $k$  sufficiently large, and given  $k$  for all  $m$  sufficiently large these arcs have the required properties.

The last statement in the proposition follows directly from this. This is contradiction and establishes the result.  $\square$

We also need estimates about the vector fields from Lemma 11.32 and also about the distances between the sides of the neighborhoods.

**Lemma 11.35.** *There is a constant  $0 < \xi_2 \leq \xi_0$  such that for any  $0 < \xi < \xi_2$  and any  $0 < s_1 \leq \alpha_0(\xi)$ , with notation and under the assumptions as in Lemma 11.34, there are positive constants  $\mu_6(\xi)$  and  $\hat{e}_5(\xi, s_1)$  such that the following hold for  $0 < \mu < \mu_6(\xi)$  and  $0 < \hat{e} < \hat{e}_5(\xi, s_1)$ .*

1. *For a unit vector field  $\tilde{\tau}_1$  on  $\nu_\xi(\tilde{\gamma}_1)$  satisfying Corollary 11.32, at any point of  $\bar{\nu}_\xi(\tilde{\gamma}_1) \cap \bar{\nu}_\xi(\tilde{\gamma}_2)$  we have*

$$\|f'_{\tilde{\gamma}_2}(\tilde{\tau}_1)\| > 1 - 50\xi.$$

2. *For any constants  $c_1, c_2$  with  $2\xi \leq c_i \leq 3/4$  and with*

$$c_1\ell_1\rho_n(x_1) < (0.9)c_2\ell_2\rho_n(x_2)$$

*each level set of  $f_{\tilde{\gamma}_2}$  in  $\bar{\nu}_{c_2\xi}(\tilde{\gamma}_2)$  that meets  $\bar{\nu}_{\xi, [-0.24\ell_1, 0.24\ell_1]}(\tilde{\gamma}_1)$  meets  $\bar{\nu}_{c_1\xi}(\tilde{\gamma}_1)$  in a disk whose boundary is contained in the side of  $\bar{\nu}_{c_1\xi}(\tilde{\gamma}_1)$ , a disk that separates the ends of  $\bar{\nu}_{c_1\xi}(\tilde{\gamma}_1)$ .*

*Proof.* The proof of the previous result show that for  $\mu > 0$  and  $\hat{e} > 0$  sufficiently small, possibly after reversing the direction of  $\tilde{\gamma}_2$ , for every  $y \in \nu_\xi(\tilde{\gamma}_1) \cap \nu_\xi(\tilde{\gamma}_2)$  we

have  $\tilde{Z}e_-(\tilde{\gamma}_1)ye_+(\tilde{\gamma}_2) > \pi - 10\xi$  and  $\tilde{Z}e_-(\tilde{\gamma}_2)ye_+(\tilde{\gamma}_1) > \pi - 10\xi$ . The first statement is immediate from this and Proposition 11.32. It follows immediately from this that any level set of  $f_{\tilde{\gamma}_{n,2}}$  meets each flow line for  $\tilde{\gamma}_1$  in at most one point.

Now let us establish the second statement. Let  $y$  be a point in

$$\nu_\xi(\tilde{\gamma}_2) \cap \nu_{\xi, [-(.24)\ell_1, (.24)\ell_1]_1}(\tilde{\gamma}_1),$$

and consider the level surface  $L$  for  $f_{\tilde{\gamma}_2}$  through  $y$ . It follows from the limiting argument in Lemma 11.34 that, given any  $\nu > 0$ , for all  $\xi > 0$ ,  $\mu > 0$  sufficiently small, and  $\hat{\epsilon} > 0$  sufficiently small given  $s_1$ , the variation of  $f_{\tilde{\gamma}_1}$  on  $L \cap \nu_\xi(\tilde{\gamma}_1)$  is less than  $\nu\xi\ell_1$ . Thus, choosing  $\xi_2, \mu_6(\xi)$ , and  $\hat{\epsilon}_5(\xi, s_1)$  sufficiently small, this implies that  $L$  does not meet the ends of  $\bar{\nu}_\xi(\tilde{\gamma}_1)$ . Thus, under the given assumptions on  $c_1$  and  $c_2$  we see that  $L \cap \left(h_{\tilde{\gamma}_2}^{-1}([0, c_2\xi])\right)$  crosses the side of  $\bar{\nu}_{c_1\xi}(\tilde{\gamma}_1)$ .

Let us consider the intersection of  $L$  with

$$U = \nu_{c_1\xi}(\tilde{\gamma}_1) \setminus \nu_{\xi^2}(\tilde{\gamma}_1).$$

On  $U$  the functions  $f_{\tilde{\gamma}_2}$  and  $h_{\tilde{\gamma}_1}$  satisfy Lemma 9.45 and hence the intersection of the level sets of these functions are circles that are almost orthogonal to the horizontal spaces in  $S^1$ -product neighborhoods with  $\epsilon$ -control, circles that meet each of these horizontal spaces in a single point. This means that  $L \cap U$  is homeomorphic to  $S^1 \times (0, 1)$  and is foliated by circles which are the intersections of  $L$  with level sets of  $h_{\tilde{\gamma}_1}$ . Since  $L$  is a disk it follows that each of these circles bounds a disk in  $L$ , and thus,  $L \cap \nu_{c_1\xi}(\tilde{\gamma}_1)$  is also a disk. Clearly, since this disk is transverse to the flow lines of the vector field and meets each flow line in at most one point, it separates the ends of  $\nu_{c_1\xi}(\tilde{\gamma}_1)$ .  $\square$

**Addendum 11.36.** In the previous two lemmas, we assumed the metrics were  $g'_n(x_1)$  and  $g'_n(x_2)$ . The reason for this was that if the  $B_{g'_n(x_i)}(x_i, s_1)$  have non-trivial intersection then these metrics are within a multiplicative factor of 2 of each other. We also have analogous results when we use the same metric,  $\lambda^2 g_n$ , in two balls. The proofs are identical, since this time the metrics agree rather than differing by at most a factor of 7.

## 11.6 Boundary points of angle $\leq \pi - \delta$

**Proposition 11.37.** *For any  $0 < \xi < \xi_0$  sufficiently small and  $a > 0$ , there is a positive constant  $\mu_7(\xi, a)$  such that for all  $\mu < \mu_7(\xi, a)$ , setting  $r_0 = r_0(\xi)$  and  $r_1 = r_1(\xi, a, \mu)$  as in Theorem 10.30, there is a positive constant  $\hat{\epsilon}_6(\xi, a, \mu)$  such that for all  $\hat{\epsilon} < \hat{\epsilon}_6(\xi, a, \mu)$  the following hold. Suppose that for some  $n$  there is a point  $x \in M_n$  with the property that  $B_{\lambda^2 g_n}(x, 1)$  is within  $\hat{\epsilon}$  of a 2-dimensional Alexandrov ball  $X = B(\bar{x}, 1)$  of curvature  $\geq -1$  and of area  $\geq a$  with the property that there is  $\bar{z} \in \partial B(\bar{x}, 1)$  satisfying Condition 2(b) in Theorem 10.30 on scale  $r$ , where  $r_1 \leq r \leq r_0$  and  $d(\bar{x}, \bar{z}) < \xi^2 r_1 / 100$  and a point  $z \in B_{\lambda^2 g_n}(x, 1)$  within  $\hat{\epsilon}$  of  $\bar{z}$ . Then  $B(z, 7r/8)$  is a topological 3-ball and the distance function,  $d(z, \cdot)$ , is regular on  $B_{\lambda^2 g_n}(z, 7r/8) \setminus B_{\lambda^2 g_n}(z, r/8)$ .*



*Proof.* The constants  $\xi$  and  $a > 0$  are given. Let  $a' > 0$  be the constant associated to  $a$  by Lemma 10.8. Suppose that there is a sequence  $\mu_k \rightarrow 0$  and for each  $k$  a sequence  $\hat{\epsilon}_{k,\ell} \rightarrow 0$  for which the result does not hold for the given values of  $\xi, a$  and for  $r_{1,k} = r_1(\xi, a, \mu_k)$  being the constant from Theorem 10.30. This implies that for each  $k, \ell$  there is a counter-example  $B_{\lambda_{k,\ell}^2 g_{n(k,\ell)}}(x_{k,\ell}, 1)$  with these values of the constants. The balls  $B_{\lambda_{k,\ell}^2 g_{n(k,\ell)}}(x_{k,\ell}, 1)$  that are within  $\hat{\epsilon}_{k,\ell}$  of 2-dimensional Alexandrov balls  $B(\bar{x}_{k,\ell}, 1)$  of curvature  $\geq -1$  and of area  $\geq a$ , and there are points  $\bar{z}_{k,\ell} \in \partial B(\bar{x}_{k,\ell}, 1)$  near which  $B(\bar{x}_{k,\ell}, 1)$  are boundary  $\mu_k$ -good near on scale  $r(\bar{z}_{k,\ell})$  where  $r_{1,k} \leq r(\bar{z}_{k,\ell}) \leq r_0$  and  $\bar{x}_{k,\ell} \in B(\bar{z}_{k,\ell}, \xi^2 r_{1,k}/4)$ , and points  $z_{k,\ell}$  are within  $\hat{\epsilon}_{k,\ell}$  of  $\bar{z}_{k,\ell}$ . Clearly,  $x_{k,\ell} \in B_{\lambda_{k,\ell}^2 g_{n(k,\ell)}}(z_{k,\ell}, \xi^2 r_{1,k}/2)$ . The ball  $B_{r(\bar{z}_{k,\ell})^{-2} \lambda_{k,\ell}^2 g_{n(k,\ell)}}(z_{k,\ell}, 1)$  is within  $r(\bar{z}_{k,\ell})^{-1} \hat{\epsilon}_{k,\ell}$  of the unit ball  $r(\bar{z}_{k,\ell})^{-1} B(\bar{z}_{k,\ell}, r(\bar{z}_{k,\ell}))$ , and the latter is boundary  $\mu_k$ -good near  $\bar{z}_{k,\ell}$  on scale 1. Fixing  $k$  and, after passing to a subsequence in  $\ell$ , the  $r^{-1}(\bar{z}_{k,\ell})$  converges to a 2-dimensional ball  $B(\bar{z}_{k,\infty}, 1)$  of area  $\geq a'$ , a ball that is  $2\mu_k$ -good at  $\bar{z}_{k,\infty}$  on scale 1.

Since the  $\mu_k \rightarrow 0$ , passing to a subsequence in  $k$ , the balls  $B(\bar{z}_{k,\infty}, 1)$  converge to a flat cone  $C$  of radius 1 in  $\mathbb{R}^2$  of angle  $\leq \pi$ . The area of  $C$  is bounded below by  $a'$ . For each  $k$  we can choose  $\ell(k)$  such that  $\hat{\epsilon}_{k,\ell(k)}/r_{1,k}$  tends to zero, and *a fortiori*  $\hat{\epsilon}_{k,\ell(k)}/r(\bar{z}_{k,\ell(k)})$  tends to zero. Then the  $B_{r(\bar{z}_{k,\ell(k)})^{-2} \lambda_{k,\ell(k)}^2 g_{n(k,\ell(k))}}(z_{k,\ell(k)}, 1)$  also converge to  $C$ , with the  $z_{k,\ell(k)}$  converging to the cone point.

At this point in the argument we simplify the notation by re-indexing things so that  $(k, \ell(k))$  becomes  $k$  and by implicitly using the metric  $r(\bar{z}_k)^{-2} \lambda_k^2 g_{n(k)}$  on  $B(x_k, 1)$ . It follows that given any  $\zeta > 0$  for all  $k$  sufficiently large, the distance function  $d_k = d(z_k, \cdot)$  is regular on  $A_k = B(z_k, 15/16) \setminus B(z_k, \zeta)$ , and in particular this annular region is homeomorphic to a product with an interval with the slices of the product structure being the level sets of the distance function. We shall achieve a contradiction by showing that these level sets are 2-spheres and that the metric balls that they bound are homeomorphic to 3-balls.

Now we fix  $\epsilon' > 0$  sufficiently small and let  $\epsilon < \epsilon_1(\epsilon')$  as in Proposition 11.8. We also suppose that  $\xi < \min(\xi_0, \xi_1(\epsilon'))$  and we fix  $s_1$  as in Theorem 10.30. By passing to a subsequence we can also assume that for all  $k$  we have  $\mu_k < \bar{\mu}(\xi)$  and  $\hat{\epsilon}_k < \hat{\epsilon}_2(\epsilon, s_1, \xi)$ . We consider first the case when the cone angle at the cone point of  $C$  is  $\pi$ . In this case,  $C$  is isometric to a unit ball centered at a boundary point of  $\mathbb{R} \times [0, \infty)$ . Since  $\mu_k \rightarrow 0$  by Proposition 11.27, there is a constant  $\zeta > 0$  depending on  $\xi$  and  $s_1$  and less than  $\alpha_0(\xi)$ , such that for all  $k$  sufficiently large there neighborhood of  $z_k$  homeomorphic to  $D^2 \times I$  that contains  $B(z_k, \zeta)$  and is contained in  $B(z_k, 1/2)$ . The boundary of this neighborhood, which is a 2-sphere, separates the level set for  $d_k = d(z_k, \cdot)$  at distance  $\zeta$  from the level set at distance  $1/2$ . Since the region in between these level sets of  $d_k$  is a product, it follows that all the level sets of  $d_k$  are 2-spheres. Furthermore, since the level set at distance  $\zeta$  is a 2-sphere contained in a neighborhood of  $z_k$  homeomorphic to a 3-ball, this level set bounds a 3-ball in this neighborhood. It follows immediately that for all  $k$  sufficiently large, all the metric balls  $B(z_k, t)$  for  $\zeta \leq t \leq 15/16$  are homeomorphic to 3-balls. This is a contradiction, proving the result in the case when the cone angle of the limit,  $C$ , is  $\pi$ .

We now examine the case when the cone angle of  $C$  is strictly less than  $\pi$ . According to the Proposition 9.49 there is a sequence of points  $z'_k \in M_{n(k)}$  with  $d(z'_k, z_k) \rightarrow 0$  such that one of the two following cases holds:

1. the distance function from  $z'_k$  has no critical points in  $B(z'_k, 1/2) \setminus \{z'_k\}$ , or
2. there is a sequence  $\zeta_k \rightarrow 0$  such that the distance function from  $z'_k$  has no critical points at distances between  $\zeta_k$  and  $1/2$  and has a critical point at distance  $\zeta_k$ .

In the first case, all the level sets for the distance function from  $z'_k$  at distance strictly between 0 and  $1/2$  are 2-spheres and the corresponding metric balls are homeomorphic to 3-balls. Since the level sets of  $d(z'_k, \cdot)$  in this range separate level sets of  $d(z_k, \cdot)$ , it follows that the level sets for  $d(z_k, \cdot)^{-1}(t)$  are 2-spheres bounding 3-balls for  $\zeta_k < t < 15/16$ . In the second case, according to Proposition 9.49 rescaling the metric by  $\zeta_k^{-2}$  we get a sequence of 3-manifolds with a subsequence converging to a 3-dimensional Alexandrov space of curvature  $\geq 0$ . By Proposition 9.46 the convergence is in fact a smooth convergence and the limit is a smooth complete 3-manifold of non-negative curvature.

**Claim 11.38.** *The level sets of the distance function  $d'_k = d(z'_k, \cdot)$  at distance between  $\zeta_k$  and  $15/16$  are topological 2-spheres.*

Let us assume this claim for a moment and complete the proof of the lemma. It follows from this claim that the end of the limiting manifold is homeomorphic to  $S^2 \times [0, \infty)$ . The limiting manifold has a soul which is a manifold of non-negative curvature. Because the neighborhood of infinity of the limit is diffeomorphic to  $S^2 \times [0, \infty)$ , the soul must be either a point or  $\mathbb{R}P^2$ . The second case is not possible, since in this case, by exactly the same argument as given in the proof of Claim 11.21, the original manifolds would converge to an interval not a 2-dimensional Alexandrov space of area  $\geq a$ . Since its soul is a point, the limiting manifold is diffeomorphic to  $\mathbb{R}^3$ . Thus, for all  $k$  sufficiently large the level sets of  $d'_k$  at distance  $2\zeta_k$  bound 3-balls, and hence for all  $k$  sufficiently large all level sets  $(d_k)^{-1}(t)$  for  $t \in [1/16, 15/16]$  are 2-spheres and the associated metric balls are 3-balls.

This shows that, modulo the claim, in all cases, for all  $k$  sufficiently large, the metric spheres at distance  $t$ , with  $1/16 \leq t \leq 15/16$ , from  $z_k$  are topological 2-spheres and the metric balls they bound are topological 3-balls. As noted before, this implies that for all  $t \in [1/16, 15/16]$  the level set  $d(z_k, \cdot)^{-1}(t)$  is also a 2-sphere and the associated metric ball is homeomorphic to a 3-ball.

It remains to prove the claim.

*Proof.* (of the claim) We continue to use the metric  $r(\bar{z}_k)^{-2} \lambda_k^2 g_{n(k)}$  on  $M_{n(k)}$ . We know that the  $B(z'_k, 15/16)$  converge to a cone  $C$  in  $\mathbb{R}^2$  of cone angle  $< \pi$  and that

$$d'_k: B(z'_k, 15/16) \setminus B(z'_k, 1/16) \rightarrow (1/16, 15/16)$$

is the projection mapping of a topologically locally trivial fibration. Let  $\gamma, \gamma'$  be arcs of length  $s_1$  on  $\partial C$  centered at the two boundary points of  $C$  at distance  $1/2$  from

the cone point. Set  $b^+ = 1/2 + (s_1/10)$  and  $b^- = 1/2 - s_1/10$ . By Proposition 11.27, for all  $k$  sufficiently large, for geodesics  $\tilde{\gamma}_k$  and  $\tilde{\gamma}'_k$  whose endpoints are within  $\hat{\epsilon}$  of those of  $\gamma$  and  $\gamma'$  respectively, the  $\epsilon$ -solid cylinder neighborhoods  $\nu_\xi(\tilde{\gamma}_k)$  and  $\nu_\xi(\tilde{\gamma}'_k)$  in  $M_{n(k)}$  satisfy the conclusions of that proposition. Let  $U_k$  be the intersection of

$$A(k) = \overline{B(z'_k, b^+)} \setminus B(z'_k, b^-)$$

with the complement of  $\nu_{\xi^2}(\tilde{\gamma}_k) \cup \nu_{\xi^2}(\tilde{\gamma}'_k)$ . Then by Proposition 11.1 for all  $k$  sufficiently large every point of  $U_k$  is the center of an  $S^1$ -product structure with  $\epsilon$ -control. Hence, by Proposition 11.8 this compact set sits inside a larger open subset that is the total space of an  $S^1$ -fibration with fibers within  $\epsilon'$  of orthogonal to the horizontal spaces of the  $S^1$ -product structures with  $\epsilon$ -control. This implies that there is an annulus in  $U_k$  with boundary contained in  $\nu_\xi(\tilde{\gamma}_k) \cup \nu_\xi(\tilde{\gamma}'_k)$  that separates  $(d'_k)^{-1}(b^+) \setminus (\nu_\xi(\tilde{\gamma}_k) \cup \nu_\xi(\tilde{\gamma}'_k))$  from  $(d'_k)^{-1}(b^-) \setminus (\nu_\xi(\tilde{\gamma}_k) \cup \nu_\xi(\tilde{\gamma}'_k))$ . Since (by Proposition 11.27) the boundary loops of this annulus are homotopically trivial in  $\nu_\xi(\tilde{\gamma}_k) \cup \nu_\xi(\tilde{\gamma}'_k)$ , it follows that there is a map of  $S^2$  into  $\overline{B(z'_k, b^+)} \setminus B(z'_k, b^-)$  that is homologically non-trivial in this region. The claim follows.  $\square$

We have now completed the proof of Proposition 11.37.  $\square$

This argument can be used to prove more, see FIG. 8.

**Corollary 11.39.** *Fix  $\epsilon' > 0$  sufficiently small and let  $\epsilon > 0$  be less than  $\epsilon_1(\epsilon')$  as in Proposition 11.8 and let  $\xi$  be a positive constant less than  $\xi_1(\epsilon)$  (recall that the latter is at most  $\xi_0$ ). For every  $a > 0$ , the following holds for all  $\mu$  less than a positive constant  $\mu_8(\epsilon, \xi, a)$ , for  $r_0, r_1$  and  $s_1$  as in Theorem 10.30 for these values of  $\xi, a, \mu$ , and for all  $\hat{\epsilon}$  less than a positive constant  $\hat{\epsilon}_7(\epsilon, \xi, a, \mu)$ . Suppose that  $x \in M_n$  has the property that  $B_{\lambda^2 g_n}(x, 1)$  is within  $\hat{\epsilon}$  of a 2-dimensional Alexandrov ball  $B(\bar{x}, 1)$  of curvature  $\geq -1$  and area  $\geq a$  and  $\bar{z} \in 7B(\bar{x}, 7/8)$  has the property that  $B(\bar{x}, 1)$  is boundary  $\mu$ -good near  $\bar{z} \in \partial B(\bar{x}, 1)$  on scale  $r$ , where  $r_1 \leq r \leq r_0$ . Then:*

1. *For any  $z \in B_{\lambda^2 g_n}(x, 1)$  within  $\hat{\epsilon}$  of  $\bar{z}$  and for any  $b \in (r/8, 7r/8)$  the level set  $L_b = d(z, \cdot)^{-1}(b)$  is a topologically locally flat 2-sphere and the metric ball that it bounds is a topological 3-ball.*
2. *There are two geodesics  $\gamma_1$  and  $\gamma_2$  of length  $r_1 s_1$  that are  $\mu$ -approximations to  $\partial B(\bar{x}, 1)$  on scale  $r_1 s_1$  whose mid-points are at distance  $b$  from  $\bar{z}$ . These arcs are within  $\xi^2 r_1 s_1 / 100$  of arcs on  $\partial B(\bar{x}, 1)$  with the same endpoints.*
3. *For any geodesics  $\tilde{\gamma}_i$  whose endpoints are within  $\hat{\epsilon}$  of those of  $\gamma_i$ , every point of  $L_b$  that is not the center of an  $S^1$ -product neighborhood with  $\epsilon$ -control is contained in union  $\nu_{\xi^2}(\tilde{\gamma}_1) \cup \nu_{\xi^2}(\tilde{\gamma}_2)$ , and these  $\epsilon$ -solid cylinder neighborhoods satisfy the conclusions of Proposition 11.27 and Proposition 11.32.*
4. *For any  $b'$  with  $|b - b'| < r_1 s_1 / 20$ , and for any point  $y \in L_{b'}$  within  $\hat{\epsilon}$  of a point  $\bar{y}$  within  $\xi^2 r_1 s_1 / 10$  of  $\partial B(\bar{x}, 1)$ , we have  $y \in \nu_{\xi^2}(\tilde{\gamma}_1) \cup \nu_{\xi^2}(\tilde{\gamma}_2)$ .*

5. The level set  $L_b$  meets each  $\bar{\nu}_\xi(\tilde{\gamma}_i)$  in a spanning 2-disk, and for any  $c \in [\xi, 1]$  the level set  $h_{\tilde{\gamma}_i}^{-1}(c\xi\ell(\tilde{\gamma}_i))$  crosses  $L_b$  topologically transversally and the intersection is a circle bounding the disk  $L_b \cap \bar{\nu}_{c\xi}(\tilde{\gamma}_i)$ .

*Proof.* The first item is included in the previous. The second item follows from Lemma 10.24 provided that  $\mu$  is sufficiently small given  $\xi$ . The third item was also established in the course of the proof of the previous proposition. Let us consider a point  $y$  as in the fourth item. Let  $\bar{y} \in B(\bar{x}, 1)$  be a point within  $\hat{\epsilon}$  of  $y$  and also within  $\xi^2 r_1 s_1 / 10$  of  $\partial B(\bar{x}, 1)$ . Then for one of  $i = 1, 2$  the point  $\bar{y}$  is within  $\xi^2 r_1 s_1 / 9$  of a point of  $\nu_{\xi, [-r_1 s_1 / 18, r_1 s_1 / 18]}(\gamma_i) \cap \gamma_i$ . Hence,  $y \in \nu_{\xi^2}(\tilde{\gamma}_i)$  provided that  $\hat{\epsilon}$  is sufficiently small given  $\xi, r_1, s_1$ . Now we establish the fifth item. Let  $f$  denote the distance function from  $z$ . Then, provided that  $\hat{\epsilon}$  is sufficiently small relative to  $s_1 r_1$  and  $\mu$  is sufficiently small, it follows from Proposition 10.29 and Lemma 10.28 that  $f$  takes values on  $L_b$  strictly less than the values on the end of  $\nu_{\xi^2}(\tilde{\gamma}_i)$  furthest from  $z$  and strictly greater than the values on the other end of  $\nu_{\xi^2}(\tilde{\gamma}_i)$ . The statement that the intersection of  $L_b \cap \bar{\nu}_{c\xi}(\tilde{\gamma}_i)$  is a circle for all  $c \in [\xi, 1]$  follows from Proposition 11.4 applied to the functions  $f$  and  $h_{\tilde{\gamma}_i}$ , again using Proposition 10.29 and Lemma 10.28. Lastly, to see that each of these circles bounds a disk in  $L_b$  we will show that  $L_b \cap \bar{\nu}_{\xi/2}(\tilde{\gamma}_i)$  is a disk. Since  $L_b \cap (\nu_\xi(\tilde{\gamma}_i) \setminus \bar{\nu}_{\xi/2}(\tilde{\gamma}_i))$  is a product region, it will then follow that  $L_b \cap \nu_\xi(\tilde{\gamma}_i)$  is an open disk and hence that  $L_b \cap h_{\tilde{\gamma}_i}^{-1}(c\xi\ell(\tilde{\gamma}_i))$  bounds  $L_b \cap \bar{\nu}_{c\xi}(\tilde{\gamma}_i)$  which is a 2-disk.

To see that  $\Delta_b = L_b \cap \bar{\nu}_{\xi/2}(\tilde{\gamma}_i)$  is a disk we flow this intersection using the vector field  $\chi$  as in Corollary 11.32 to the end of  $\nu_\xi(\tilde{\gamma})$ . According to this corollary any flow line through the boundary of  $\Delta_b$  remains in  $\nu_\xi(\tilde{\gamma})$  until it meets the end of the  $\xi$ -box closest to  $e_+$ . Using Proposition 10.29 and Lemma 10.28 we see that each flow line of the vector field crosses  $L_b$  at most once. Thus, flowing in this manner produces an embedding of  $\Delta_b$  into the end of the  $\xi$ -box. Since the latter is a disk, since  $\Delta_b$  is compact, and since  $\partial\Delta_b$  consists of a single circle, it follows that  $\Delta_b$  is also a disk.  $\square$

**Definition 11.40.** Any time we have  $z \in B_{\lambda^2 g_n}(x, 1)$  satisfying Proposition 11.37 and Corollary 11.39 with  $r = r(\bar{z})$ , we say that the ball  $B_{\lambda^2 g_n}(z, r/4)$  is a 3-ball near a 2-dimensional corner.

### 11.6.1 Intersection of 3-balls near 2-dimensional corners and $\xi$ -boxes

**Lemma 11.41.** Given  $0 < \xi < \xi_0$ ,  $a > 0$  and  $\mu < \mu_1(a, \xi)$  and given  $r_0, r_1, s_1, s_2$  at most the constants of the same names depending on  $\xi, \mu, a$  given in Theorem 10.30 with  $s_1 \leq 10^{-3}$ , there is a positive constant  $\hat{\epsilon}_8(\xi, a, \mu, r_0, r_1, s_1, s_2)$  such that the following hold for all  $\hat{\epsilon} < \hat{\epsilon}_8(\xi, a, \mu, r_0, r_1, s_1, s_2)$ . Suppose that for some  $r$  with  $r_1 \leq r \leq r_0$  the ball  $B_{g'_n(x)}(z, r/4)$  is a 3-ball near a 2-dimensional corner with the property that the associated 2-dimensional Alexandrov ball  $B(\bar{x}, 1)$  is boundary  $\mu$ -good on scale  $r$  with  $r_1 \leq r \leq r_0$  at a point  $\bar{z}$  and  $d(\bar{z}, \bar{x}) < \xi^2 r_1 / 100$  with  $z$  within  $\hat{\epsilon}$  of  $\bar{z}$  and with  $(1/r)B(\bar{z}, r)$  having a  $(\mu, s_1, s_2)$ -good collar region. Suppose that  $\nu_\xi(\tilde{\gamma})$  is an  $\epsilon$ -solid cylinder on scale  $r_1 s_1$ , meaning that there is a ball  $B_{g'_n(x')}(x', 1)$  within  $\hat{\epsilon}$  of a 2-dimensional Alexandrov ball  $B(\bar{x}', 1)$  with a point  $\bar{y}' \in \partial B(\bar{x}', 1)$  with  $d(\bar{y}', \bar{x}') <$

$\xi^2 r_1 s_1 / 100$  such that  $B(\bar{x}', 1)$  is boundary  $\mu$ -flat near  $\bar{y}$  on all scales  $\leq r_1 s_1$ , and furthermore, there is a geodesic  $\bar{\gamma}$  of length  $r_1 s_1 / 4$  contained in  $B(\bar{y}', r_1 s_1 / 3)$  with endpoints in  $\partial B(\bar{y}', r_1 s_1 / 3)$  with the property that the endpoints of  $\tilde{\gamma}$  are within  $\hat{\epsilon}$  of those of  $\bar{\gamma}$ . Suppose that  $\nu_\xi(\tilde{\gamma}) \cap B_{g'_n(x)}(z, 7r/8) \neq \emptyset$ . Then  $\tilde{\gamma} \subset B_{g'_{n_k}(x)}(z, r)$  and either:

(i)  $\nu_\xi(\tilde{\gamma}) \subset B_{g'_n(x)}(x, r/16)$  or,

(ii)  $\tilde{\gamma}$  is within  $\xi^2 r_1 s_1 / 50$  of an arc on  $\partial B(\bar{z}, r)$  and, there is an orientation for  $\tilde{\gamma}$  so that  $d(z, \cdot)$  has directional derivative at least  $1 - \xi^2/2$  at every point of  $\nu_\xi(\tilde{\gamma}) \cap \tilde{\gamma}$  in the positive direction along  $\tilde{\gamma}$ .

*Proof.* Suppose that the result does not hold for some  $\xi$  with  $0 < \xi < \xi_0$ ,  $\mu < \mu_1(a, \xi) \leq \bar{\mu}(\xi)$ . Then there is a sequence  $\hat{\epsilon}_k$  tending to zero as  $k \rightarrow \infty$  and counter-examples  $B_k = B_{g'_{n(k)}(x_k)}(x_k, 1)$  and points  $z_k \in B_k$  within  $\hat{\epsilon}_k$  of 2-dimensional Alexandrov balls  $\bar{B}_k = B(\bar{x}_k, 1)$  and points  $\bar{z}_k \in \partial B(\bar{x}_k, 1)$  as in the statement for a constant  $\hat{r}_k$  with  $r_1 \leq \hat{r}_k \leq r_0$ . Also, there are  $\tilde{\gamma}_k$  generating  $\nu_k = \nu_\xi(\tilde{\gamma}_k)$ . The  $\nu_\xi(\tilde{\gamma}_k)$  are contained in balls  $B'_k = B_{g'_{n(k)}(x'_k)}(x'_k, 1)$  within  $\hat{\epsilon}_k$  of a 2-dimensional Alexandrov balls  $\bar{B}'_k$  with points  $\bar{y}'_k \in \partial \bar{B}'_k$  and geodesics  $\bar{\gamma}_k$  as in the statement. Let  $y'_k \in B'_k$  be a point within  $\hat{\epsilon}_k$  of  $\bar{y}'_k$ . Notice that, since  $\hat{r}_k < 10^{-6}$ , by Lemma 6.1 we see that  $R_k^2 g'_{n(k)}(x'_k) = g'_{n(k)}(x_k)$  for a constant  $R_k$  satisfying  $(0.99) < R_k < (1.01)$ . Passing to a subsequence we can suppose that the  $\bar{B}_k$  converge an Alexandrov ball  $\bar{B}_\infty = B(\bar{x}_\infty, 1)$ , the  $\bar{z}_k$  converge to  $\bar{z}_\infty \in \partial \bar{B}_\infty$ , the  $\bar{B}'_k$  converge to  $\bar{B}'_\infty = B(\bar{x}'_\infty, 1)$ , the  $\bar{y}'_k$  converge to a point  $\bar{y}'_\infty \in \partial \bar{B}'_\infty$ , the geodesics  $\bar{\gamma}_k$  converge to a geodesic  $\bar{\gamma}_\infty$  of length  $r_1 s_1 / 4$  with endpoints in  $\partial B(\bar{y}'_\infty, r_1 s_1 / 3)$ , the  $R_k$  converge to a constant  $R$ , and the  $\hat{r}_k$  converge to  $\hat{r}_\infty$ . We have  $RB_{g'_{n(k)}(x'_k)}(y'_k, r_1 s_1) = B_{g'_{n(k)}(x_k)}(y'_k, Rr_1 s_1)$  and, as a result,  $RB_{g'_{n(k)}(x'_k)}(y'_k, r_1 s_1)$  is identified with a sub-ball of  $B_{g'_{n(k)}(x_k)}(z_k, \hat{r}_k)$ . Since  $\hat{\epsilon}_k \rightarrow 0$ , passing to the limit  $R\bar{B}(\bar{y}'_\infty, r_1 s_1)$  is identified with a sub-ball of  $B(\bar{z}_\infty, \hat{r}_\infty)$ . If  $d(\bar{y}'_\infty, \bar{z}_\infty) \leq \hat{r}_\infty / 32$ , then  $\tilde{\gamma}_k \subset B_{g'(x_k)}(z_k, \hat{r}_k / 16)$  for all  $k$  sufficiently large and the contradiction is established.

Thus, we can (and shall) assume that  $d(\bar{y}'_\infty, \bar{z}_\infty) > \hat{r}_\infty / 32$ . Since  $\mu < \mu_1(a, \xi)$ , according to Lemma 10.24 we have that  $\bar{\gamma}_\infty$  is within  $\xi^2 r_1 s_1 / 100$  of the arc on  $\partial B(\bar{y}'_\infty, r_1 s_1)$  with the same endpoints as  $\bar{\gamma}_\infty$ . Hence  $R\bar{\gamma}_\infty$  is within  $R\xi^2 r_1 s_1 / 100$  of the arc on  $\partial B(\bar{z}, \hat{r}_\infty) \subset \bar{B}_\infty$  with the same endpoints. The geodesics  $R\tilde{\gamma}_k$  in  $B_{g'_{n(k)}(x_k)}(z_k, \hat{r}_k)$  converge to  $R\bar{\gamma}_\infty \subset B(\bar{z}_\infty, \hat{r}_\infty)$ . Consequently, for all  $k$  sufficiently large  $R\tilde{\gamma}_k$  is within  $R\xi^2 r_1 s_1 / 100$  of the arc  $\bar{\alpha}_\infty$  on the boundary of  $B(\bar{z}_\infty, \hat{r}_\infty)$  with the same endpoints as  $R\bar{\gamma}_\infty$ . Let  $\bar{\alpha}_k$  be arcs on  $\partial B(\bar{z}_k, \hat{r}_k)$  converging to  $\bar{\alpha}_\infty$ . Then, for all  $k$  sufficiently large,  $\tilde{\gamma}_k$  is within  $\xi^2 r_1 s_1 / 50$  of  $\bar{\alpha}_k \subset \partial B(\bar{z}_k, \hat{r}_k)$ . Let  $e_+(\bar{\alpha}_k)$  be the endpoint of this  $\bar{\alpha}_k$  furthest from  $\bar{z}_k$ . For  $k$  sufficiently large consider any point  $w_k$  in  $\tilde{\gamma}_k \cap \nu_\xi(\tilde{\gamma}_k)$ , and let  $e_+(\tilde{\gamma}_k)$  be the endpoint of  $\tilde{\gamma}_k$  furthest from  $z_k$ . Let  $\bar{w}_k \in \bar{\gamma}_\infty$  be a closest point on  $\bar{\gamma}_\infty$  to  $w_k$ . Then, passing to a subsequence, we can assume that the  $\bar{w}_k$  converge to  $\bar{w}_\infty \in \bar{\gamma}_\infty$  at distance at least  $\hat{r}_\infty / 64$  from  $\bar{z}_\infty$ . Thus, as  $k$  goes to  $\infty$  the comparison angle  $\angle z_k w_k e_+(\tilde{\gamma}_k)$  converges to  $\angle \bar{z}_\infty \bar{w}_\infty e_+(\bar{\gamma}_\infty)$ . By Lemma 10.28, this latter comparison angle is greater than  $\pi - \xi$ . It now follows that

for all  $k$  sufficiently large, at any point of  $\tilde{\gamma}_k \cap \nu_\xi(\tilde{\gamma}_k)$  the directional derivative of  $d(\bar{z}_k, \cdot)$  in the tangent direction along  $\tilde{\gamma}_k$  pointing toward  $e_+(\tilde{\gamma}_k)$  is greater than  $1 - \xi^2/2$ . This proves that the conclusions of the lemma hold for all  $k$  sufficiently large, which is a contradiction, establishing the lemma.  $\square$

### 11.7 Balls near open intervals

Now we are ready to describe the parts of  $M_n$  close to 1-dimensional Alexandrov balls.

**Lemma 11.42.** *Given  $\epsilon' > 0$  the following holds for all  $\beta$  less than a positive constant  $\beta_1(\epsilon')$ . Suppose that  $B = B_{g'_n(x)}(x, 1)$  is within  $\beta$  of a 1-dimensional ball  $J$  and  $y \in B_{g'_n(x)}(x, 24/25)$  is within  $\beta$  of a point  $\bar{y}$  whose distance from every endpoint of  $J$  is at least  $1/25$ . For any  $z \in B$  with  $d_{g'_n(x)}(y, z) \geq 1/30$  let  $f = d_{g'_n(x)}(z, \cdot) - d_{g'_n(x)}(z, y)$  and set  $U = f^{-1}(-3/100, 3/100)$ . Then  $B_{g'_n(x)}(y, 1/50) \subset U$  and  $f|_U: U \rightarrow (-3/100, 3/100)$  is an  $\epsilon'$ -approximation for which the following hold:*

1.  *$f$  is the projection mapping of a topological product structure.*
2. *The fibers of  $p$  are homeomorphic to either 2-spheres or 2-tori.*
3. *There is a smooth unit vector field  $\chi$  on  $U$  such that for any (minimal) geodesic  $\gamma$  of length  $\geq 1/4000$ , measured in the metric  $g'_n(x)$ , ending at a point  $w \in U$ , the angle at  $w$  between  $\chi(w)$  and  $\gamma'(w)$  is within  $\epsilon'$  of either 0 or  $\pi$ .*
4. *Given  $w, w' \in B_{g'_n(x)}(y, 1/50)$  with  $d_{g'_n(x)}(w, w') \geq 1/4000$ , the connected component of the level surface of the distance function  $d_{g'_n(x)}(w, \cdot)$  through  $w'$  is contained in  $U$  and is isotopic in  $U$  to a fiber of  $p$ .*

*Proof.* It is easy to see that if  $\beta$  is sufficiently small, then  $f$  is an  $\epsilon'$ -approximation.

Fix  $z \in B$  with  $d_{g'_n(x)}(y, z) \geq 1/30$ . Let  $\bar{z} \in J$  be a point within  $\beta$  of  $z$ . Provided that  $\beta$  is sufficiently small there is a point  $\bar{w} \in J$  with  $d(\bar{y}, \bar{w}) \geq .031$ ,  $d(\bar{w}, \bar{z}) > 0.001$ , and  $\bar{w}$  separates  $\bar{y}$  and  $\bar{z}$  in  $J$ . Let  $w \in B$  be within  $\beta$  of  $\bar{w}$ . Then for any  $u \in U$ , the comparison angle  $\tilde{\angle} zwu$  is close to  $\pi$ , and the discrepancy  $d(\beta)$  from  $\pi$  goes to zero (uniformly for all  $u \in U$ ) with  $\beta$ . It follows that all geodesics from  $z$  to any point  $u \in U$  all have tangent vectors at  $u$  that make an angle at most  $d(\beta)$  with each other. Hence, there is a smooth vector field  $\chi$  on  $U$  such that for every  $u \in U$  the angle between  $\chi(u)$  and any geodesic from  $z$  to  $u$  is at least  $\pi - 2d(\beta)$ . This means that (again assuming that  $\beta$  is sufficiently small) that  $f$  is regular and hence the level sets of  $f$  are Lipschitz surfaces fibering  $U$ . Furthermore, the vector field  $\chi$  is transverse to these level sets in the sense that the level curves of  $\chi$  cross each level surface exactly once. Thus, these level curves can be used to define a product structure on  $U$  so that  $f$  is the projection onto the interval factor.

Now consider any geodesic  $\gamma$  of length at least  $1/4000$  ending at a point  $u \in U$ . By shortening  $\gamma$  if necessary we can suppose that the other endpoint  $w$  is contained in  $B(y, 1/25)$ . One of the comparison angles  $\tilde{\angle} zuw$  or  $\tilde{\angle} zwu$  is close to  $\pi$ . It then follows by monotonicity that the angle at  $u$  between  $\gamma$  and  $\chi$  is also close to either 0 or  $\pi$ .

Next, we argue that, provided that  $\beta > 0$  is sufficiently small, the fibers of  $p$  are either 2-spheres or 2-tori. If not we there is a sequence of  $x_k \in M_{n(k)}$ , constants  $\beta_k \rightarrow 0$  and examples  $f_k: U_k \rightarrow (-3/100, 3/100)$  with fibers  $L_k = p_k^{-1}(t_k)$  that are not 2-spheres or 2-tori. Fix points  $w_k \in L_k$ , let  $d_k$  be the diameter of  $L_k$  and rescale, forming  $\frac{1}{d_k}(U_k, w_k)$ , and, after passing to a subsequence take a limit. This limit is an Alexandrov space and since  $d_k \rightarrow 0$ , it is of dimension 2 or 3 and splits as a product  $\mathbb{R} \times Y$  where  $Y$  has diameter 1. If  $Y$  is 2-dimensional, then by Proposition 9.46 the convergence is smooth and  $Y$  is a surface of curvature  $\geq 0$ . Since  $Y$  is orientable, it follows in this case that  $Y$  and hence the fibers  $L_k$ , for all  $k$  sufficiently large, are homeomorphic to either 2-spheres or 2-tori, which is a contradiction.

Suppose that  $Y$  is 1-dimensional. Then it is either a closed interval or circle, and  $d_k^{-1}U_k$  converge to the product  $\mathbb{R} \times Y$ . If  $Y$  is a circle, we invoke Lemma 11.1 and Proposition 11.8 to see that for all  $k$  sufficiently large, any level set of  $f_k$  is contained in an open subset  $V_k \subset d_k^{-1}U_k$  that is the total space of a circle fibration. We can take a slightly smaller compact fibration  $W_k \subset V_k$  still containing the level set. The boundary components of  $W_k$  are tori and at least one of them separates the two ends of  $d_k^{-1}U_k$ . On the other hand, the level set  $L_k$  separates two of the boundary components of  $W_k$ . These two facts together imply that for all  $k$  sufficiently large,  $L_k$  is a 2-torus, in contradiction to our assumption.

Lastly, suppose that  $Y$  is a closed interval. Then invoking Lemma 11.1, Proposition 11.8 and Proposition 11.27 we see that for all  $k$  sufficiently large every level set of  $f_k$  is contained in the union of the total space of an  $S^1$ -fibration over an annulus and two disjoint, simply connected sets of the form  $\nu_\xi(\tilde{\gamma}_i)$  as in Proposition 11.27. Arguing as in the proof of Claim 11.38, we see that there is a map of the 2-sphere into  $U_k$  that separates its ends. It then follows that the fibers of  $f_k$  are 2-spheres.

The last item follows easily from the third item.  $\square$

**Definition 11.43.** A neighborhood  $U \subset B$ , and a projection mapping  $f: U \rightarrow J$  satisfying the 4 conditions in the conclusions of the previous lemma is called an *interval product structure  $\epsilon'$ -control* or an  *$\epsilon'$  interval product structure*. If  $y \in U$  and if the image of  $f$  is  $(-3/100, 3/100)$  with  $f(y) = 0$ , that the  $\epsilon$ -interval product structure is centered at  $y$ .

The content of the above lemma is that for  $\beta < \beta_1(\epsilon')$  if  $B_{g'_n(x)}(x, 1)$  is within  $\beta$  of a 1-dimensional Alexandrov ball  $J$  and if  $y \in B_{g'_n(x)}(x, 24/25)$  is within  $\beta$  of a point of  $J$  that has distance at least  $1/25$  from the endpoints (if any) of  $J$ , then there is an interval product structure with  $\epsilon'$ -control centered at  $y$ .

Now we need to understand what happens near the endpoints of the nearby interval.

**Proposition 11.44.** *There is  $a_1 > 0$  such that the following holds. Fix  $\epsilon' > 0$  and  $\zeta > 0$ . Then the following holds for all  $\beta$  less than a positive constant  $\beta_2(\epsilon', \zeta)$ . For some  $n$  suppose that the ball  $B_{g'_n(x)}(x, 1)$  is contained in the interior of  $M_n$  and this ball is within  $\beta$  in the Gromov-Hausdorff sense to an interval  $J$  and the  $x$  is within  $\beta$  of a point  $\bar{x}$  which is at distance at most  $1/25$  from the endpoint of  $J$ . Then for*

any point  $y \in B_{g'_n(x)}(x, 1)$  within  $\beta$  of the endpoint of  $J$ , setting  $f = d_{g'_n(x)}(y, \cdot)$ , the restriction of  $f$  to

$$B_{g'_n(x)}(y, 5/8) \setminus B_{g'_n(x)}(y, 1/1000)$$

is the projection mapping of an  $\epsilon'$  interval product structure, Furthermore, one of the following holds:

1. The closed ball  $\overline{B_{g'_n(x)}(y, 1/2)}$  is diffeomorphic to one of the following: a solid torus, to a twisted  $I$ -bundle over the Klein bottle, to a 3-ball, or to  $\mathbb{R}P^3 \setminus B^3$ .
2. There is a constant  $\lambda \gg 1$  such that the ball of radius 2 centered at  $y$  in  $\lambda B_{g'_n(x)}(x, 1)$  is within distance  $\zeta$  of the ball  $B(\bar{y}, 2)$  in a complete two-dimensional Alexandrov space  $(X, \bar{x})$  of curvature  $\geq 0$ . Furthermore, the function  $d_{g'_n(x)}(y, \cdot)$  is regular on  $B_{g'_n(x)}(y, 1/2) \setminus B_{g'_n(x)}(y, 1/3\lambda)$  and fibers this subset over  $[1/3\lambda, 1/2)$  with fibers either 2-spheres or 2-tori. Similarly,  $d(\bar{y}, \cdot)$  is regular on  $X \setminus B(\bar{y}, 1/3)$  and fibers this subset over  $[1/3, \infty)$  with fibers which are either topological intervals or simple closed curves. For any  $y \in \overline{B_{\lambda^2 g_n}(x, 1/3)}$  the ball  $B_{\lambda^2 g_n}(y, 1)$  is within  $4\zeta$  of the ball  $B(\bar{y}, 1) \subset B(\bar{x}, 2)$  for any  $\bar{y} \in B(\bar{x}, 1/2)$  within  $\zeta$  of  $y$ . Finally, for any  $\bar{y} \in B(\bar{x}, 1/2)$  the area of  $B(\bar{y}, 1)$  is at least  $a_1$ .

*Proof.* Fix  $\theta$  with  $\pi/2 < \theta < \pi$ , and let  $a_1 = a_1(\theta)$  as in Proposition 9.48. Fix  $\epsilon' > 0$ ,  $\zeta > 0$ , and suppose that there is no constant  $\beta_2(\epsilon', \zeta) > 0$  as required. Then we have a sequence  $\beta_k \rightarrow 0$  as  $k \rightarrow \infty$  and points  $x_{n_k} \in M_{n_k}$  such that  $B_{n_k} = B_{g'_{n_k}(x_{n_k})}(x_{n_k}, 1)$  is within  $\beta_k$  of a closed interval with  $x_{n_k}$  being within  $\beta_k$  of the endpoint of the interval but none of these examples satisfy the conclusion of the proposition. It follows that the  $B_{n_k}$  converge to an interval and the  $x_{n_k}$  converge to its endpoint. At many different steps in the proof we shall pass to a subsequence using the notation  $n_k$  for the subsequence. First notice that by Lemma 11.42 for all  $k$  sufficiently large there is an open subset  $U_{n_k}$  as required with an  $\epsilon'$ -product structure. Proposition 9.48 tells us that for all  $k$  sufficiently large there is another point  $\hat{x}_{n_k} \in M_{n_k}$ , such that the sequence  $\hat{x}_{n_k}$  also converges to the endpoint, such that one of two possibilities holds:

1. the distance function from  $\hat{x}_{n_k}$  has no points within distance  $1/2$  of  $\hat{x}_{n_k}$  (except of course  $\hat{x}_{n_k}$ ) at which the maximum value of the directional derivative of the distance function from  $\hat{x}_{n_k}$  is at most  $\theta$ , or
2. there is a sequence  $\zeta_{n_k} \rightarrow 0$  such that all points within distance  $1/2$  of  $\hat{x}_{n_k}$  where the maximum of the directions derivative of the distance function from  $\hat{x}_{n_k}$  is at most  $\theta$  are in fact within  $\zeta_{n_k}/3$  of  $\hat{x}_{n_k}$  and there is a such a point at distance  $\zeta_{n_k}/3$  from  $\hat{x}_{n_k}$ .

In the first case, the closed balls  $\overline{B_{g'_{n_k}(x_{n_k})}(\hat{x}_{n_k}, t)}$  are topological 3-balls for all  $0 < t < 1/2$ . Since for all  $k$  sufficiently large, the distance from  $x_{n_k}$  is regular and its level sets are close to the corresponding level sets of the distance function from  $\hat{x}_{n_k}$ , it follows that  $\overline{B_{g'_{n_k}(x_{n_k})}(x_{n_k}, 1/2)}$  is homeomorphic to a topological 3-ball. By



Lemma 11.42 its boundary separates the ends of  $U_{n_k}$ . This proves that the result (indeed as in 1.) holds for all  $n_k$  for  $k$  sufficiently large in this case, which is a contradiction.

In the second case we rescale by multiplying the metric by  $\zeta_{n_k}^{-2}$ , pass to a subsequence, and take a limit. The resulting limit is a complete Alexandrov space  $(X, \bar{x})$  of non-negative curvature and of dimension 2 or 3 with  $\bar{x}$  being the limit of the  $\hat{x}_{n_k}$ . We consider first the sub-case when the result is 3-dimensional. By Proposition 9.46 it is a complete 3-manifold of non-negative curvature, and as such it has a soul. If the soul is a point, then the limit is diffeomorphic to  $\mathbb{R}^3$  and level sets of the distance function from  $\bar{x}$  are 2-spheres. If the soul is a circle, then the limit is a solid torus and the level sets of the distance function from  $\bar{x}$  are 2-tori. If the soul is a Klein bottle, then the manifold is a twisted  $\mathbb{R}$ -bundle over the Klein bottle and the level sets of the distance function from  $\bar{x}$  are 2-tori. If the soul is  $\mathbb{R}P^2$ , then the limit is a punctured  $\mathbb{R}P^3$  and the level sets are 2-spheres. Thus, in these cases, for all  $k$  sufficiently large the original  $B(\hat{x}_{n_k}, 1/2)$  is diffeomorphic to the limiting complete manifold. All the distance function from  $\hat{x}_{n_k}$  is regular at distances between  $\zeta_{n_k}/3$  and  $1/2$  the level sets of the distance function from  $\hat{x}_{n_k}$  at distances between  $\zeta_{n_k}/3$  and  $1/2$  are parallel. Clearly, the level set  $d(\hat{x}_{n_k}, \cdot)^{-1}(1/2)$  is contained in  $U_{n_k}$  and separates the ends of  $U_{n_k}$ . For all  $k$  sufficiently large, the level set  $d(x_{n_k}, \cdot)^{-1}(1/2)$  is close to the level set  $d(\hat{x}_{n_k}, \cdot)^{-1}(1/2)$  and is parallel to it in  $U_{n_k}$ . Thus, all the above statements hold for the balls  $B(x_{n_k}, t)$  for all  $1/4 \leq t \leq 1/2$  for all  $k$  sufficiently large. This shows that in this sub-case the result holds (again as in 1.) for all  $k$  sufficiently large, which is a contradiction.

Suppose now that the limit of the rescalings is 2-dimensional  $(X, \bar{x})$ . Then by Proposition 9.48 for any  $\bar{y} \in B(\bar{x}, 1/2)$  the ball  $B(\bar{y}, 1)$  has area at least  $a_1$ . The fact that there are no critical points for the distance function from  $\hat{x}_{n_k}$  at distances greater than  $\zeta_{n_k}/3$ , and indeed no points in this range where the maximum directional derivative of the distance function is less than  $\theta$ , imply the statements about the fibration structure for both  $X \setminus \overline{B(\bar{x}, 1/3)}$  and for the

$$\overline{B_{g'_{n_k}(x_{n_k})}(\hat{x}_{n_k}, 1/2)} \setminus B_{g'_{n_k}(x_{n_k})}(\hat{x}_{n_k}, \zeta_{n_k}/3)$$

for all  $k$  sufficiently large. Since the boundary of  $\zeta_{n_k}^{-1} B_{g'(x_{n_k})}(x_{n_k}, \zeta_{n_k}/2)$  is parallel to the fibers of the  $\epsilon'$ -product structure on  $U_{n_k}$ , it is homeomorphic to either a 2-torus or a 2-sphere. Since these level sets converge in the Gromov-Hausdorff topology as  $k \rightarrow \infty$  to the level set of the distance function from  $\bar{x}$  at distance  $1/2$ , it follows that the latter level set is connected. Since it is a compact Lipschitz 1-manifold, it is either a simple closed curve or a closed interval. This then is true for all the level sets of the distance function from  $\bar{x}$  at distances greater than  $1/3$ . This shows that the result (as in 2.) holds, for all  $n_k$  for  $k$  sufficiently large. This is a contradiction.  $\square$

## 11.8 Determination of the Constants

We fix  $\epsilon' < 10^{-6}$  a universally small positive constant and let  $\epsilon > 0$  less than the minimum of the constants  $\epsilon_0(\epsilon')$  in Proposition 11.4,  $\epsilon_1(\epsilon')$  in Proposition 11.8,  $\epsilon_2(\epsilon')$  in Proposition 11.20, and sufficiently small so that Lemma 9.45 holds. Now we fix  $\xi$

with  $0 < \xi < \min(\xi_0, \xi_1(\epsilon), \xi_2)$  where  $\xi_0$  is defined near the end of Section 10,  $\xi_1(\epsilon)$  is given in Proposition 11.27 and  $\xi_2$  is given in Lemma 11.35. Having fixed  $\xi$  we also have  $\alpha = \alpha_0(\xi)$ .

Now we define a function  $\hat{\epsilon}(a)$  depending on a positive constant  $a$ . We do this as follows: Given  $a$  we choose  $\mu > 0$  sufficiently small so that Lemma 10.26 holds. We also take it to be less than the minimum of  $\delta(a'(a))/8$  where  $\delta$  is the constant from Lemma 10.7,  $(1/2)\mu_0''(10^{-6}, a'(a))$  from Proposition 10.18,  $\mu_1(a, \xi)$  from Theorem 10.30,  $\mu_2(\epsilon)$  from Lemma 11.1,  $\mu_3(\epsilon, a)$  from Proposition 11.20 as modified in Proposition 11.22,  $\bar{\mu}(\xi)$  from Proposition 11.27 and Addendum 11.29,  $\mu_4(\xi)$  from Lemma 11.31,  $\mu_5(\xi)$  from Lemma 11.34,  $\mu_6(\xi)$  from Lemma 11.35,  $\mu_7(\xi, a)$  from Proposition 11.37, and  $\mu_8(\epsilon, \xi, a)$  from Corollary 11.39. Now we choose  $\delta_0(a), r_0(a), r_1(a), r_2(a), s_0(a), s_1(a), s_2(a)$  positive functions of  $a$  as in Theorem 10.30 for the given values of  $\xi$  and  $\mu$ . With all of these determined, we are ready to define  $\hat{\epsilon}(a)$  for every  $a > 0$ . It is the minimum of:

$$r_2(a)\delta(a'(a))/20 \quad \text{where } \delta \text{ is the constant from Lemma 10.7,}$$

$$(r_1(a)/50)\mu_0''(10^{-6}, a'(a)) \quad \text{from Proposition 10.18,}$$

$$\hat{\epsilon}_0(\epsilon, \min(r_1 s_2, \xi^2 r_1 s_1/100, r_2 s_0)) \quad \text{from Lemma 11.1,}$$

$$\hat{\epsilon}'_0(\xi^2 r_1 s_1/100, a'(a)) \quad \text{from Lemma 11.26,}$$

$$\hat{\epsilon}_1(\epsilon, a, r_1, r_2) \quad \text{from Proposition 11.20 and Proposition 11.22,}$$

$$\hat{\epsilon}_2(\epsilon, \xi, r_1 s_1) \quad \text{from Proposition 11.27 and Addendum 11.29,}$$

$$\hat{\epsilon}_3(\epsilon, \xi, r_1 s_1) \quad \text{from Lemma 11.31,}$$

$$\hat{\epsilon}_4(\xi, \mu, r_1 s_1) \quad \text{from Lemma 11.34,}$$

$$\hat{\epsilon}_5(\xi, r_1 s_1) \quad \text{from Lemma 11.35,}$$

$$\hat{\epsilon}_6(\xi, a, \mu) \quad \text{from Proposition 11.37,}$$

$$\hat{\epsilon}_7(\epsilon, \xi, a, \mu) \quad \text{from Corollary 11.39,}$$

$$\hat{\epsilon}_8(\xi, a, \mu, r_0, r_1, s_1, s_2) \quad \text{from Lemma 11.41. and}$$

$$\xi^2 r_1 s_1/100C \quad \text{where } C \text{ is the constant in Corollary 11.7.}$$

Next we fix  $\zeta > 0$  to be less than  $\hat{\epsilon}(a_1)/3$  where  $a_1$  is the constant in Proposition 11.44. We then fix  $\beta > 0$  less than  $\min(\beta_0, \beta_1(\epsilon'), \beta_2(\epsilon', \zeta), 10^{-8})$  where  $\beta_0$  is the constant in Lemma 9.6,  $\beta_1(\epsilon')$  is as in Lemma 11.42, and  $\beta_2(\epsilon', \zeta)$  is as in Proposition 11.44. Now that we have fixed  $\beta$  we set  $a = \min(a_1, a_2(\beta/2))$ , the latter constant being as in Lemma 10.31. This fixes the constants  $\delta_0 = \delta_0(a), r_0 = r_0(a), r_1 = r_1(a), s_0 = s_0(a), s_1 = s_1(a), s_2 = s_2(a)$ .

We require  $0 < \hat{\epsilon} < \min(\hat{\epsilon}(a), \beta/2)$ . We fix  $\hat{\epsilon} > 0$  satisfying all these conditions. Now we pass to a subsequence of the  $M_n$  so that the constant  $\epsilon_n$  from Lemma 9.51 is  $\leq \hat{\epsilon}$  for all  $n$ , and also so that Proposition 11.19 holds for all  $n$  with  $\hat{\epsilon}$  taken equal to  $\beta$ .

### 11.8.1 Effect of these choices

By the definition of  $\epsilon_n$ , for every  $x \in M_n$  there is an Alexandrov ball  $B(\bar{x}, 1)$  of curvature  $\geq -1$  and of dimension either 1 or 2 such that  $B_{g'_n(x)}(x, 1)$  is within  $\epsilon_n$  in the Gromov-Hausdorff topology of  $B(\bar{x}, 1)$ . We divide into three cases:

1.  $B(\bar{x}, 1)$  is 2-dimensional and of area  $\geq a_2(\beta/2)$ .
2.  $B(\bar{x}, 1)$  is 2-dimensional and of area  $< a_2(\beta/2)$ .
3.  $B(\bar{x}, 1)$  is 1-dimensional.

In the second case by Lemma 10.31  $B(\bar{x}, 1)$  is within  $\beta/2$  of a 1-dimensional Alexandrov ball, and hence  $B_{g'_n(x)}(x, 1)$  is within  $\beta$  of a 1-dimensional Alexandrov ball. Thus, these three cases lead to the following two cases:

1.  $B_{g'_n(x)}(x, 1)$  is within  $\hat{\epsilon}$  of a 2-dimensional Alexandrov ball  $B(\bar{x}, 1)$  of curvature  $\geq -1$  and area  $\geq a$ , or
2.  $B_{g'_n(x)}(x, 1)$  is within  $\beta$  of an interval  $J$ .

As indicated in the definition below, having fixed the choices of all the constants from now on, we redefine the terms  $\epsilon$ -solid torus,  $\epsilon$ -solid cylinder, and 3-ball near a 2-dimensional corner so as to restrict to the cases of interest. First of all these notions will mean implicitly that they are with respect to all the constants that we just fixed. Also, in each case there will be one extra condition that was not originally required.

**Definition 11.45.** 1. An  $\epsilon$ -solid torus is a metric ball  $B_{g'_n(x)}(x, r/2) \subset M_n$  where  $B_{g'_n(x)}(x, 1)$  satisfies the conclusions of Proposition 11.20 with the given values of  $\epsilon', \epsilon, a, \mu, r_0, r_1, r_2, s_0$ , and  $\hat{\epsilon}$ , and with  $r_2 \leq r \leq r_1$ . The 2-dimensional Alexandrov ball  $B(\bar{x}, 1)$  as in that proposition, called the *associated* 2-dimensional Alexandrov ball, is  $\mu$ -good at  $\bar{x}$  on scale  $r$ . The extra condition in this case is that the cone angle of the close circular cone is required to be  $\leq 2\pi - \delta_0$ .

2. An  $\epsilon$ -solid cylinder is a subset of the form  $\nu_\xi(\tilde{\gamma}) \subset M_n$  satisfying Proposition 11.27. Thus, there are  $B = B_{g'_n(x)}(x, 1) \subset M_n$  containing  $\nu_\xi(\tilde{\gamma})$  and an *associated* 2-dimensional Alexandrov ball  $B(\bar{x}, 1)$  as in that proposition within  $\hat{\epsilon}$  of  $B$ , a point  $\bar{y} \in \partial B(\bar{x}, 1)$  with the property that  $B(\bar{x}, 1)$  is boundary  $\mu$ -flat near  $\bar{y}$  on all scales  $\leq r_1 s_1$  and there is a geodesic  $\bar{\gamma} \subset B(\bar{y}, r_1 s_1/3)$  of length  $r_1 s_1/4$  that is a  $\mu$ -approximation to the boundary with the endpoints of  $\tilde{\gamma}$  within  $\hat{\epsilon}$  of those of  $\bar{\gamma}$ . Lastly, the extra condition that we require in this case is  $d(\bar{x}, \bar{y}) < \xi^2 r_1 s_1/100$ .

3. A 3-ball near a 2-dimensional corner is a  $B_{g'_n(x)}(z, r/4) \subset B_{g'_n(x)}(x, 1)$  with  $r_1 \leq r \leq r_0$ , with an *associated* 2-dimensional Alexandrov ball  $B(\bar{x}, 1)$  and a point  $\bar{z} \in \partial B(\bar{x}, 1)$  as in Proposition 11.37. Thus,  $d(\bar{x}, \bar{z}) < \xi^2 r_1/100$  with  $B(\bar{x}, 1)$  being boundary  $\mu$ -good near  $\bar{z}$  on scale  $r$ . The extra conditions in this case is that we require  $B(\bar{x}, 1)$  to be boundary  $\mu$ -good near  $\bar{z}$  on scale  $r$  and angle  $\leq \pi - \delta_0$ .

Let us summarize what we have established so far.

**Theorem 11.46.** *The following hold for every  $n \geq 0$ :*

1. *Let  $Y_{1,n} \subset M_n$  be the open subset of all  $x \in M_n$  with the property that  $B_{g'_n(x)}(x, 1)$  is within  $\beta$  of an interval. Then for any  $x \in Y_{1,n}$  one of the following cases holds:*

- (a) *The ball  $B_{g'_n(x)}(x, 1)$  meets the boundary of  $M_n$  and there is an open subset  $V$  of  $M_n$  homeomorphic to  $T^2 \times [0, 1)$  that contains  $x$ . There is a neighborhood  $U$  of the end of  $V$  has an  $\epsilon'$  interval product structure with base being an interval of length  $3/50$  with fibers homeomorphic to 2-tori.*
- (b) *The ball  $B_{g'_n(x)}(x, 1)$  is disjoint from  $\partial M_n$ , and  $x$  is within  $\beta$  of a point of the interval whose distance to the endpoint of the interval is at least  $1/25$ . Then there is an open subset  $U \subset B_{g'_n(x)}(x, 1)$  that contains  $B_{g'_n(x)}(x, 1/50)$  with an  $\epsilon'$  interval product structure  $p: U \rightarrow (-3/100, 3/100)$  with fibers which are homeomorphic to either 2-tori or 2-spheres.*
- (c) *The ball  $B_{g'_n(x)}(x, 1)$  is disjoint from  $\partial M_n$  and  $x$  is within  $1/25$  of the endpoint. Then either:*
  - (i) *there is an open subset  $V \subset B_{g'_n(x)}(x, 1)$  containing  $B_{g'_n(x)}(x, 1/25)$  with a neighborhood  $U$  of the end of  $V$  that has an  $\epsilon'$  interval product structure as above, and  $V$  is homeomorphic to one of the following: a solid torus, a twisted  $I$ -bundle over the Klein bottle, a 3-ball, or the complement of a 3-ball in  $\mathbb{R}P^3$ , or*
  - (ii) *the Conclusion 2 of Proposition 11.44 holds for  $B_{g'_n(x)}(x, 1)$  with  $\zeta = \hat{\epsilon}(a_1)/3$ . In particular, there is a constant  $\lambda > \rho_n^{-1}(x)$  such that every point  $y$  in the closure of  $B_{\lambda^2 g_n}(x, 1/3)$  the ball  $B_{\lambda^2 g_n}(y, 1)$  is within  $\hat{\epsilon}(a_1)$  of a 2-dimensional Alexandrov ball of curvature  $\geq 0$  and area  $\geq a_1 \geq a$ .*

2. *Let  $Y_{2,n} \subset M_n$  be the open subset of all  $x \in M_n$  with the property that  $B_{g'_n(x)}(x, 1)$  is within  $\hat{\epsilon}$  of a 2-dimensional Alexandrov ball of curvature  $\geq -1$  and area  $\geq a$ . Then  $Y_{2,n}$  is covered by the union of the following sets: (i) the open subset  $U_{2,\text{gen}}$  of points that are centers of  $S^1$ -product neighborhoods with  $\epsilon$ -control, (ii) a collection of  $\epsilon$ -solid tori, (iii) a collection of 3-balls near 2-dimensional corners, and (iv) a collection of cores of  $\epsilon$ -solid cylinders.*

3.  $M_n = Y_{1,n} \cup Y_{n,2}$ .

*Proof.* Case 1, when  $B_{g'_n(x)}(x, 1)$  is within  $\beta$  of an interval, is immediate from Proposition 11.19, Lemma 11.42, and Proposition 11.44.

In the second case, if the 2-dimensional Alexandrov ball  $B(\bar{x}, 1)$  within  $\hat{\epsilon}$  of the ball  $B_{g'_n(x)}(x, 1)$  is interior  $\mu$ -flat at  $\bar{x}$  on scale  $r_2$ , then  $x$  is the center of an  $S^1$ -product structure with  $\epsilon$ -control. If this ball is interior  $\mu$ -good at  $\bar{x}$  on scale  $r$  with  $r_2 \leq r \leq r_1$  and angle  $\leq 2\pi - \delta_0$ , then  $B_{g'_n(x)}(x, r/2)$  is an  $\epsilon$ -solid torus. Otherwise, according to Theorem 10.30 there is a point  $\bar{y} \in \partial B(\bar{x}, 1)$  with  $d(\bar{x}, \bar{y}) < \xi^2 r_1 s_1 / 100$  and  $B(\bar{x}, 1)$  is either boundary  $\mu$ -flat near  $\bar{y}$  on all scales  $\leq r_1 s_1$  or it is boundary  $\mu$ -good near  $\bar{y}$  on some scale  $r$  with  $r_1 \leq r \leq r_0$  and angle  $\leq \pi - \delta_0$ . In the first case, there is an  $\epsilon$ -solid cylinder  $\nu_\xi(\tilde{\gamma})$  whose core contains  $x$  and indeed, given any  $-r_1 s_1 / 16 \leq c \leq r_1 s_1 / 16$  we can choose this  $\epsilon$ -solid cylinder so that  $f_{\tilde{\gamma}}(x) = c$ . In

the last case, for any  $y \in B_{g'_n(x)}(x, 1)$  within  $\hat{\epsilon}$  of  $\bar{y}$ , the ball  $B_{g'_n(x)}(y, r/4)$  is a 3-ball near a 2-dimensional corner containing  $x$ .  $\square$

Suppose that  $B = B_{g'(x)}(x, 1)$  is within  $\hat{\epsilon}$  of  $\bar{B} = B(\bar{x}, 1)$  and  $\bar{y} \in B(\bar{x}, 1/2)$ . Then:

1. If  $\bar{B}$  is interior  $\mu$ -good at  $\bar{y}$  on scale  $r$  with  $r_2 \leq r \leq r_1$ , then any point of  $B$  within  $\hat{\epsilon}$  of a point of  $B(\bar{y}, 7r/8) \setminus B(\bar{y}, r/8)$  is contained in  $U_{2,\text{gen}}$ .
2. If  $\bar{B}$  is boundary  $\mu$ -flat at  $\bar{y}$  on all scales  $\leq r_1 s_1$ , then any point of  $B$  that is within  $\hat{\epsilon}$  of a point of  $B(\bar{y}, r_1 s_1/2)$  at distance at least  $\xi^2 r_1 s_1/100$  from  $\partial B(\bar{y}, r_1 s_1)$  is contained in  $U_{2,\text{gen}}$ .
3. If  $\bar{B}$  is boundary  $\mu$ -good at  $\bar{y}$  on scale  $r$  with  $r_1 \leq r \leq r_0$  and angle  $\leq \pi - \delta_0$ , and if  $q \in B$  within  $\hat{\epsilon}$  of a point of  $\bar{q} \in (B(\bar{y}, 7r/8) \setminus B(\bar{y}, r/8))$  with the distance from  $\bar{q}$  to  $\partial B(\bar{y}, 1)$  being at least  $\xi^2 r_1 s_1/100$ , then  $q \in U_{2,\text{gen}}$ .

## 12 The global result

At this point we have fixed all the constants appearing in the last two sections in such a way that the conclusions of all the results from these two sections hold. As we have seen, this gives us complete control over the local nature of the  $(M_n, g_n)$  in the sense that we have complete control over a neighborhood of every  $x \in M_n$  whose size is determined by  $\rho_n(x)$ . The purpose of this section is to globalize these results establishing Theorem 6.2. Since we have arranged that  $\epsilon_n < \hat{\epsilon}$  for all  $n$ , the arguments of this section apply uniformly for all  $n$ . For this reason, for most of the rest of this section we drop  $n$  from the notation and denote by  $(M, g)$  one of the Riemannian manifolds  $(M_n, g_n)$ . We denote the function  $\rho_n: M_n \rightarrow \mathbb{R}$  by  $\rho$  and by  $g'(x)$  the Riemannian metric  $\rho^{-2}(x)g$ .

**Definition 12.1.** Given a ball  $B_{\lambda^2 g}(x, r)$  we say that  $r$  is its *rescaled radius* and  $r/\lambda$  is its *unrescaled radius*.

### 12.1 Regions of $M$ close to open intervals

We begin the globalization by studying the generic “1-dimensional” regions of  $M$ . We shall construct a compact submanifold with boundary  $W_1 \subset M$  which is a first approximation to the submanifold  $V_1 \subset M$  (dropping the subscript  $n$ ) referred to in Theorem 6.2. The manifold  $V_1$  will be obtained by deforming  $W_1$  by an isotopy supported near its boundary components.

**Definition 12.2.** We define  $X_1 \subset M$  to be the subset consisting of all points  $y \in M$  for which there is  $x \in M$  with  $d_{g'(x)}(x, y) < 1/10$  and with  $B_{g'(x)}(x, 1)$  being within  $\beta$  in the Gromov-Hausdorff distance of a 1-dimensional Alexandrov ball  $J$  in such a way that  $y$  is within  $\beta$  of a point  $\bar{y} \in J$  whose distance from any endpoint of  $J$  is greater than  $1/25$ . We say that the pair  $(x, y)$  satisfying these conditions is an  $X_1$  pair.

Notice that since  $B_{g'(x)}(x, 1)$  is non-compact, if the ball  $B_{g'(x)}(x, 1)$  is within  $\beta$  of an interval  $J$  and if  $d_{g'(x)}(x, y) < 1/10$ , then the distance from  $y$  any non-compact end of  $J$  is greater than  $9/10 - 2\beta$ .

**Definition 12.3.** We set  $\widehat{U} = \cup_{y \in X_1} B_{g'(y)}(y, 1/100)$ .

**Claim 12.4.** For any  $z \in \widehat{U}$  there is an  $X_1$ -pair  $(x, y)$  with  $d_{g'(x)}(y, z) < 0.011$  and hence  $z \in B_{g'(x)}(x, 0.111)$ .

*Proof.* Since  $y \in B_{g'(x)}(x, 1/10)$  it follows from Lemma 6.1 that  $\rho(y)/\rho(x) \leq 1.1$ . Since  $d_{g'(y)}(y, z) < 1/100$ , the claim is immediate.  $\square$

Thus,  $z$  is within  $\beta$  of a point  $\bar{z} \in J$  which is at least  $(1/25) - 0.011 - 2\beta$  from any endpoint of  $J$  and hence  $B_{g'(x)}(z, 1/50)$  is within  $4\beta$  of the sub interval  $B(\bar{z}, 1/50)$  of  $J$ . This is an open interval of length  $1/25$  centered at  $\bar{z}$ . Using the fact (from Lemma 6.1) that  $\rho(z)/\rho(x)$  is between 0.889 and 1.111 it follows that  $B_{g'(z)}(z, 1/100)$  is within  $5\beta$  of an open interval  $I(\bar{z})$  of length  $1/50$  centered at a point  $\bar{z}$  within  $2\beta$  of  $z$ . From this it follows that there is a smooth line field on  $\widehat{U}$  with the property that if  $\gamma$  is any geodesic ending at a point  $z \in \widehat{U}$  and if the length of  $\gamma$  in the metric  $g'(z)$  is at least  $10^{-3}$ , then the angle at  $y$  between  $\gamma$  and the line field is less than  $1/100$ .

Now we consider  $U^+ = \cup_{y \in X_1} (B_{g'(y)}(y, 1/400))$  and  $U^- = \cup_{y \in X_1} B_{g'(y)}(y, 1/500)$ . Clearly,  $U^- \subset U^+ \subset \widehat{U}$ . Suppose that  $z$  is in the frontier  $F$  of  $U^-$  in  $M$ . Then there is a sequence  $y_n \in X_1$  with the property that  $d_{g'(y_n)}(y_n, z) \rightarrow 1/500$  as  $n \rightarrow \infty$ . In particular, for all  $n$  sufficiently large we have  $y_n \in B_{g'(z)}(z, 1/100)$  and  $d_{g'(z)}(y_n, z) > 1/600$ . Thus,  $y$  is within  $5\beta$  of a point  $I(\bar{z})$  and all points of  $I(\bar{z})$  within  $5\beta$  of  $y_n$  lie in the same component of  $I(\bar{z}) \setminus \{\bar{z}\}$ .

**Definition 12.5.** We say that  $z$  is a *one-sided frontier point* of  $U^-$  if for every sequence  $y_n \in X_1$  with  $d_{g'(y_n)}(y_n, z) \rightarrow 1/500$  for all  $n$  sufficiently large the  $y_n$  are within  $5\beta$  of points  $\bar{y}_n$  of  $I(\bar{z})$  on the same side of  $\bar{z}$ . Otherwise we say that  $z$  is a *two-sided frontier point* of  $U^-$ .

Our goal is to expand  $U^-$  slightly until every point of its frontier is a one-sided point. It is easy to see that if  $z$  is a two-sided frontier point of  $U^-$  then  $B_{g'(z)}(z, 1/450)$  is contained in  $U^+$ . We form the union of  $U^-$  with the union of the  $B_{g'(z)}(z, 1/500)$  as  $z$  ranges over the two-sided frontier points of  $U^-$ . We call the result  $U_1$ .

**Claim 12.6.** The open subset  $U_1$  contains  $U^-$  and is contained in  $U^+$ . Any point of the frontier of  $U_1$  is a one-sided frontier point.

*Proof.* The first statement is clear. Let  $w$  be a point of the frontier of  $U_1$ , say  $w$  is the limit of  $y_n \in U_1$ . We claim that for all  $n$  sufficiently large  $y_n \in U^-$ . For, if  $z$  is a two-sided frontier point of  $U^-$  then  $B_{g'(z)}(z, 1/450) \subset B_{g'(z)}(z, 1/500) \cup U^-$ . Thus, the distance, measured in  $g'(z)$ , from  $y_n$  to the frontier of  $U_1$  is at least  $1/500 - 1/450$ , and hence the distance measured in  $g'(w)$  from  $w$  to  $y_n$  is bounded below by a positive constant independent of  $n$ . This is impossible. This shows that

the frontier of  $U_1$  is contained in the frontier of  $U^-$ . Clearly, no two-sided frontier point of  $U^-$  is contained in the frontier of  $U$ .  $\square$

Since  $U_1 \subset U^+ \subset \widehat{U}$ , there is a line field on  $U_1$  with the property that any geodesic ending at a point  $y \in U_1$  of length at least  $10^{-3}$  measured with respect to  $g'(y)$  makes angle at  $y$  less than  $1/100$  with the line field. In particular, for any  $x$  at distance at least  $10^{-3}$  from  $y$ , measured with respect to  $g'(y)$  the distance function  $d_{g'(x)}(x, \cdot)$  has directional derivative  $> .99$  in one of the two directions along the line field. In particular, the integral curves of the line field meet each component of  $U_1$  in a connected open set. Hence the components of  $U_1$  are diffeomorphic to  $T^2 \times (0, 1)$ ,  $S^2 \times (0, 1)$  or a bundle over the circle with fiber either  $T^2$  or  $S^2$ .

**Proposition 12.7.** *There is an open subset  $U'_1 \subset U_1 \subset M$  containing  $X_1$  with the following properties:*

1. *The closure  $\overline{U}'_1$  of  $U'_1$  is a compact submanifold with topologically locally flat boundary and  $U'_1$  is its interior.*
2. *The difference  $U_1 \setminus \overline{U}'_1$  is a disjoint union of connected neighborhoods of the ends of  $U_1$ . Each component of  $U_1 \setminus \overline{U}'_1$  has diameter less than  $100\beta$  and the integral curves of the line field on  $U_1$  foliate this difference by proper open intervals, so that  $U_1 \setminus \overline{U}'_1$  is diffeomorphic to a product of a surface with an open interval.*
3. *Each component of  $U'_1$  either is a 2-torus bundle over the circle, or is diffeomorphic to a product of either  $S^2$  or  $T^2$  with an interval.*
4. *For each end  $\mathcal{E}$  of  $U'_1$  there is an  $X_1$ -pair  $(x, x_{\mathcal{E}})$  and a neighborhood  $U(x_{\mathcal{E}}) \subset U'_1$  of  $\mathcal{E}$  for which there is an interval product structure  $p_{x_{\mathcal{E}}}: U(x_{\mathcal{E}}) \rightarrow J(x_{\mathcal{E}})$  with  $4\beta$ -control. Here  $U(x_{\mathcal{E}})$  is given the metric  $g'(x_{\mathcal{E}})$ , the length of the interval  $J(x_{\mathcal{E}})$  is  $1/250$ , and this interval is centered at  $\bar{x}_{\mathcal{E}} = p_{x_{\mathcal{E}}}$ .*
5.  *$U'_1$  is contained in the union of  $B_{g'(y)}(y, 1/400)$  for  $y \in X_1$ .*
6. *For each point  $y \in X_1$ , the ball  $B_{g'(y)}(y, 1/501)$  is contained in  $U'_1$ .*

*Proof.* For each end of  $U_1$  there are a point  $z$  in the corresponding component of the frontier of  $U_1$  and a sequence  $(x_n, y_n)$  of  $X_1$ -pairs converging to  $(x_{\infty}, y_{\infty})$  (which is not necessarily an  $(X_1$ -pair) such that  $d_{g'(y_{\infty})}(y_{\infty}, z) = 1/500$ . We take  $(x, x_{\mathcal{E}})$  to be  $(x_n, y_n)$  for some  $n$  sufficiently large that  $d_{g'(y_n)}(y_n, y_{\infty}) < \beta$ . Then we set  $U(x_{\mathcal{E}})$  to be  $B_{g'(x_{\mathcal{E}})}(x_{\mathcal{E}}, 1/500)$ . Of course,  $U(x_{\mathcal{E}}) \subset U^- \subset U_1$ . Then the distance function from  $x$  foliates the closure of  $U(x_{\mathcal{E}})$  by locally flat surfaces in  $M$  so that  $U(x_{\mathcal{E}})$  is the interior of a compact, codimension-0 submanifold with locally flat boundary. Exactly one of the boundary components of each  $U(x_{\mathcal{E}})$  is within  $10\beta$  of the frontier of  $U_1$ , when measured in the metric  $g'(x_{\mathcal{E}})$ . We call this the *exterior end* of  $U(x_{\mathcal{E}})$ . We define  $U'_1$  to be the open submanifold of  $U_1$  obtained by removing the region between the exterior ends of the  $U(x_{\mathcal{E}})$  and the corresponding frontier of  $U_1$ . The

complement  $U_1 \setminus \bar{U}'_1$  is diffeomorphic to the product with an interval – the product structure being given by the integral curves of the line field on  $U_1$ .

The third item is clear from the construction of  $U_1$  and  $U'_1$ . The  $4\beta$ -approximation is given by

$$U(x_{\mathcal{E}}) \xrightarrow{f} (-1/250, 1/250) \xrightarrow{\bar{f}} B(\bar{x}_{\mathcal{E}}, 1/250),$$

where  $\bar{x}_{\mathcal{E}}$  is any point of  $B(\bar{x}, 1)$  within  $\beta$  of  $x_{\mathcal{E}}$ , where  $f = .d(x, \cdot) - d(x, x_{\mathcal{E}}$  and  $\bar{f} = d(\bar{x}, \cdot) - d(\bar{x}, \bar{x}_{\mathcal{E}})$ .  $\square$

We fix  $U'_1 \subset M$  as in the above proposition. For each non-compact end  $\mathcal{E}$  of  $U'_1$  we fix an  $X_1$  pair  $(x, x_{\mathcal{E}})$  producing the neighborhood  $U(x_{\mathcal{E}})$  of the end together with a projection mapping  $p_{x_{\mathcal{E}}} : U(x_{\mathcal{E}}) \rightarrow J(x_{\mathcal{E}})$  as in Conclusion 3 of Proposition 12.7.

Now we begin the study of the complementary regions  $M \setminus U'_1$ .

**Definition 12.8.** Suppose that  $x \in M$  has the property that  $B_{g'(x)}(x, 1)$  is within  $\beta$  of an interval  $J$  and that  $x$  is within  $\beta$  of  $\bar{x} \in J$  with  $\bar{x}$  at distance at most  $1/25$  of an endpoint  $e$  of  $J$ . Then we say  $x$  is *close to a 1-dimensional endpoint*. In this case we define  $V(x)$  to be the open set of all points  $y \in B_{g'(x)}(x, 1)$  with the property  $y$  is within  $\beta$  of a point  $\bar{y} \in J$  within distance  $0.09$  of  $e$ .

**Claim 12.9.** Suppose that  $x \in M$  is close to a 1-dimensional endpoint and suppose that  $y \in B_{g'(x)}(x, 1)$  is within  $\beta$  of the endpoint of the corresponding interval. Then:

1.  $V(x)$  is an open subset of  $M$ .
2. The subset  $V(x) \cap d_{g'(x)}(y, \cdot)^{-1}(0.055, 0.099)$  contains the non-compact end of  $V(x)$  and is contained in  $U'_1$ .
3.  $d_{g'(x)}(y, \cdot)^{-1}(0.06, 0.08)$  is contained in  $V(x)$ .
4. The distance function  $d_{g'(x)}(y, \cdot)$  is regular on  $d_{g'(x)}(y, \cdot)^{-1}(0.06, 0.08)$  and each fiber of  $d_{g'(x)}(y, \cdot)$  in this open set is a surface isotopic in  $U'_1$  to the fiber of its fibration structure.
5. For every  $t \leq 0.1$  the fiber  $\Sigma_t = \{z \mid d_{g'(x)}(y, z) = t\}$  has diameter  $\leq 4\beta$  in the metric  $g'(x)$ .

*Proof.* The first item is clear. It is also clear that  $V(x) \cap d_{g'(x)}(y, \cdot)^{-1}(0.055, 0.099)$  contains the non-compact end of  $V(x)$ . By definition of  $X_1$ , this intersection is contained in  $X_1$  and hence it is contained in  $U'_1$ . Also, it is also clear that

$$Z(x) = d_{g'(x)}(y, \cdot)^{-1}(0.06, 0.08)$$

is contained in  $V(x)$ . Furthermore, clearly  $d_{g'(x)}(y, \cdot)$  is regular on  $Z(x)$  so that the level sets of this map are compact surfaces. The directional derivative of the distance function from  $y$  makes an angle close to either  $0$  or  $\pi$  with the vector field given in Lemma 11.42 and hence this vector field can be used to deform the level sets of  $d_{g'(x)}(y, \cdot)$  to a fiber of the fibration structure on  $U'_1$ . Lastly, for each  $t \leq 0.1$  the level set  $\Sigma_t$  is within  $2\beta$  of the point of the interval at distance  $t$  from the endpoint. It follows that  $\Sigma_t$  has diameter less than  $4\beta$ .  $\square$



**Claim 12.10.** *Suppose that  $x, x'$  are points in  $M$ , each close to a 1-dimensional endpoint. Then:*

1.  $V(x)$  is not contained in  $U'_1$ .
2. If  $V(x) \cap V(x') \neq \emptyset$ , then there is a connected component of  $M \setminus U'_1$  contained in  $V(x) \cap V(x')$ .

*Proof.* Suppose that  $B_{g'(x)}(x, 1)$  is within  $\beta$  of an interval  $J$  and that  $y \in V(x)$  is a point within  $\beta$  of the endpoint  $e$  of  $J$ . We shall prove the first statement by showing that  $y \notin U'_1$ . To do this we show that there is no point  $z \in X_1$  with  $d_{g'(z)}(z, y) < 1/400$  and invoke Condition 4 from Proposition 12.7. Suppose to the contrary that such a  $z$  exists. By the definition of  $X_1$  there is a point  $w$  with  $B_{g'(w)}(w, 1)$  within  $\beta$  of an interval  $J'$ , with  $d_{g'(w)}(w, z) < 1/10$  and with  $z$  within  $\beta$  of a point  $\bar{z} \in J'$  at distance greater than  $1/25$  from every endpoint of  $J$  (and also from every non-compact end of  $J'$ ). First, notice by Lemma 6.1 that  $g'(w)/g'(z)$  is equal to a constant between  $9/11$  and  $11/9$ , so that consequently  $d_{g'(w)}(y, z) < 1/200$ . Thus,  $y \in B_{g'(w)}(w, 1/9)$  and  $y$  is within  $\beta$  of a point  $\bar{y}' \in J'$  at distance at least  $1/30$  from the endpoints and non-compact ends of  $J'$ . Since  $B_{g'(w)}(w, 1/9) \cap B_{g'(x)}(x, 1/10) \neq \emptyset$ , it follows that for the constant  $R$  defined by  $R^2 = g'(w)/g'(x)$  we have  $(4/5) < R < (5/4)$ . Of course,  $R \cdot B_{g'(x)}(y, 1/2) = B_{g'(w)}(y, R/2)$ . Thus, the ball of radius  $R/2$  about  $e$  in  $R \cdot J$  and the ball of radius  $R/2$  about  $\bar{y}'$  in  $J'$  are within  $4(1+R)\beta$  of each other in the Gromov-Hausdorff topology. But this is absurd since  $e$  is an endpoint of  $J$  and  $\bar{y}'$  has distance at least  $1/30$  from the ends of  $J'$ . This contradiction shows that  $y \notin U'_1$ .

Suppose that  $x$  and  $x'$  are close to one-dimensional endpoints. Let  $y \in B_{g'(x)}(x, 1)$  and  $y' \in B_{g'(x')}(x', 1)$  be within  $\beta$  of the endpoints of the corresponding intervals. Also, suppose that  $V(x) \cap V(x') \neq \emptyset$ . This implies that  $g'(x)/g'(x')$  is a constant  $R^2$  with  $9/11 < R < 11/9$ . Let  $\Sigma$  be the level set  $d_{g'(x)}(y, \cdot)^{-1}(0.095)$  and let  $\Sigma'$  be a level set  $d_{g'(x')}(y', \cdot) = d$  for some  $0.095 < d < 0.098$ , chosen so that  $\Sigma \cap \Sigma' = \emptyset$ . Such a  $d$  exists since the diameter of  $\Sigma$ , respectively  $\Sigma'$ , is less than  $4\beta$  in the metric  $g'(x)$ , respectively  $g'(x')$  and  $g'(x)$ , and  $g'(x')$  differ by a multiplicative factor  $R^2$  with  $9/11 < R < 11/9$ . Both  $\Sigma$  and  $\Sigma'$  are contained in  $U'_1$ . Set  $V'(x) = \overline{B_{g'(x)}(y, 0.095)}$  and  $V'(x') = \overline{B_{g'(x')}(y', d)}$ . These are compact connected manifolds with connected boundary  $\Sigma$  and  $\Sigma'$ , respectively. Then  $V(x) \subset V'(x)$  and  $V(x') \subset V'(x')$  and the complements  $V'(x) \setminus V(x)$  and  $V'(x') \setminus V(x')$  are contained in  $X_1$  and hence are contained in  $U'_1$ .

Since  $V'(x)$  and  $V'(x')$  are compact, connected submanifolds with disjoint connected boundaries, either  $V'(x) \cap V'(x') = \emptyset$ ,  $V'(x) \subset V'(x')$ ,  $V'(x') \subset V'(x)$ , or  $V'(x) \cup V'(x')$  is a component of  $M$ . The last possibility cannot hold for it if did then the component would be the union of two open sets, each of diameter  $< 1/4$  with respect to  $g'(x)$  and this is impossible, since by our choice of  $\rho$ , the ball  $B_{g'(x)}(x, 1)$  is non-compact. According what was established in the first part of this proof,  $V(x)$  and  $V(x')$  each contain a connected component of  $M \setminus U'_1$ , and by the above, any such component is disjoint from  $(V'(x) \setminus V(x)) \cup (V'(x') \setminus V(x'))$ . Thus, the second

and third possibilities satisfy the conclusion of the claim. The first possibility is ruled out since it is contrary to the supposition that  $V(x) \cap V(x') \neq \emptyset$ .  $\square$

For each  $x$  close to a 1-dimensional endpoint,  $V(x)$  is the union of an open subset of  $U'_1$  and some finite, non-empty collection of complementary components. It follows from the previous claim that we can find a finite set of such points  $x_1, \dots, x_k$  close to 1-dimensional endpoints such that the  $V(x_i)$  are disjoint and every component of  $M \setminus U'_1$  that is contained in  $V(x)$  for any point  $x$  close to a 1-dimensional endpoint is contained in one of the  $V(x_i)$ . We fix these  $x_i$  and  $V(x_i)$ . For each  $i$  we fix a point  $y_i \in V(x_i)$  within  $\beta$  of the endpoint of the corresponding interval. We denote by  $V_0(x_i)$  the compact submanifold cut off by the surface  $\Sigma_i$  which is a level set of  $d_{g'(x_i)}(y_i, \cdot)$  at distance 0.07 from  $y_i$ .

The conclusions of Proposition 11.19 or Proposition 11.44 hold for  $V_0(x_i) \subset B_{g'(x_i)}(x_i, 1)$ :

**Corollary 12.11.** *One of the following hold:*

1. *Int  $V_0(x_i)$  is homeomorphic to (a) an open 3-ball, (b) the complement of a closed 3-ball in  $\mathbb{R}P^3$ , (c) an open solid torus, (d) an open twisted  $I$ -bundle over the Klein bottle, or (e)  $T^2 \times [0, 1)$  and its boundary is a boundary component of  $M$ .*
2. *Case 1 does not hold, and there is a constant  $\lambda_i \gg \rho^{-1}(x_i)$  and such that  $B = B_{\lambda_i^2 g}(y_i, 2)$  is within  $\hat{\epsilon}(a)$  of a ball of radius 2 in a complete 2-dimensional Alexandrov space of curvature  $\geq 0$  satisfying the Conclusion 2(b) of Lemma 11.44, and  $d(y_i, \cdot)$  has no critical points in  $V_0(x_i) \setminus \overline{B_{\lambda_i^2 g}(y_i, 1/3)}$ . (Recall that for every  $z \in B_{\lambda_i^2 g}(y_i, 1)$  the ball of radius 1 centered at  $z$  is within  $\hat{\epsilon}$  of an Alexandrov ball  $B(\bar{z}, 1)$  of curvature  $\geq 0$  and of area at least  $a_1$  and hence of area at least  $a$ .)*

Furthermore, for each  $i \leq k$  the surface  $\Sigma_i$  is contained in  $U'_1$  and  $\Sigma_i$  is isotopic in  $V(x_i) \cap U'_1$  to a fiber of the fibration structure of  $U'_1$  (of course  $\Sigma_i$  is either a 2-sphere or a 2-torus). Thus, in the first case the union of  $V_0(x_i)$  with the component of  $U'_1$  containing the boundary of  $V_0(x_i)$  is diffeomorphic to  $\text{Int } V_0(x_i)$ . In the second case, the region between  $\overline{B_{\lambda_i^2 g}(y_i, 1/3)}$  and  $\Sigma_i$  is a topological product.

### 12.1.1 An expansion of $U'_1$

After renumbering we can assume the subsets  $V_0(x_i)$ ,  $i = 1, \dots, \ell$ , satisfy the second conclusion in Corollary 12.11 and the subsets  $V_0(x_i)$ ,  $i = \ell + 1, \dots, k$ , satisfy the first. We define

$$U''_1 = U'_1 \cup \bigcup_{i=\ell+1}^k V_0(x_i).$$

Some of the components of  $U''_1$  are components of  $U'_1$  and some are strictly larger. Let us consider components of the latter type. Fix a component  $C''$  of  $U''_1$  that is not a component of  $U'_1$ . Then there is a  $V_0(x_i) \subset C''$ . If there is only one such  $V_0(x_i)$  contained in  $C''$ , then  $C''$  is the union of a component  $C'$  of  $U'_1$  and  $V_0(x_i)$ . Since

the boundary of  $V_0(x_i)$  is parallel in  $C'$  to the fiber of the fibration structure on  $C'$ , it follows that in this case  $C''$  is diffeomorphic to  $\text{Int } V_0(x_i)$ .

Suppose there are indices  $i \neq i'$  both greater than  $\ell$  such that  $V_0(x_i)$  and  $V_0(x_{i'})$  are both contained in  $C''$ . Since  $V_0(x_i) \cap V_0(x_{i'}) = \emptyset$  and since  $V_0(x_i)$  and  $V_0(x_{i'})$  each have only one non-compact end, it follows that  $C''$  is the union of  $V_0(x_i)$ ,  $V_0(x_{i'})$ , and a connected component  $C'$  of  $U'_1$ . Again using the fact that the boundaries  $V_0(x_i)$  and  $V_0(x_{i'})$  are parallel in  $C'$  to fibers of the fibration structure, we see that  $C''$  is a closed component of  $M$  and is diffeomorphic to the union of  $V_0(x_i)$  and  $V_0(x_{i'})$  along their boundary. Being the union of two manifolds each of which is homeomorphic to the closure of one of the five listed in Conclusion 1 of Corollary 12.11 glued together along their common boundary, every one of the prime factors of the closed manifold  $C''$  is geometric, or is diffeomorphic to  $T^2 \times I$ . (The manifold is prime unless it is  $S^3$  or  $\mathbb{R}P^3 \# \mathbb{R}P^3$ .)

From now on we work with  $U''_1$  and the  $V_0(x_i)$  satisfying the second conclusion of Corollary 12.11. Thus, when we refer to  $V_0(x_i)$  we implicitly are assuming that  $1 \leq i \leq \ell$ .

Invoking the hypothesis that the boundary of  $M$  consists of incompressible tori and that no closed component of  $M$  admits a Riemannian metric of non-negative sectional curvature, allows us to conclude the following:

**Proposition 12.12.** *The open subset  $U''_1 \subset M$  constructed in the previous paragraph satisfies the following:*

1. *Every component of  $U''_1$  is diffeomorphic to one of the following:*
  - (a) *a  $T^2$ -bundle or an  $S^2$ -bundle over either the circle or an interval with the fiber(s) over the endpoint(s) being boundary component(s) of  $M$ ,*
  - (b) *a twisted  $I$ -bundle over the Klein bottle whose boundary is a boundary component of  $M$ ,*
  - (c) *an open solid torus, an open twisted  $I$ -bundle over the Klein bottle, an open 3-ball, the complement of a closed 3-ball in  $\mathbb{R}P^3$ , or*
  - (d) *the union of two twisted  $I$ -bundles over the Klein bottle along their common boundary.*
2. *For each non-compact end of  $U''_1$  is also an end of  $U'_1$  and hence there is a neighborhood of each non-compact end of  $U''_1$  of the form  $U(x_\mathcal{E}) \subset U'_1$  as in Proposition 12.7.*

Now we turn to the complement of  $U''_1$ .

**Proposition 12.13.** *Let  $A$  be a connected component of  $M \setminus U''_1$ . Then one of the following two things holds.*

1. *For some  $1 \leq i \leq \ell$ , we have  $A \subset V_0(x_i)$ . Furthermore:*
  - (a)  *$A \subset B_{g'(x_i)}(y_i, 1/20)$ .*

- (b)  $V_0(x_i) \setminus \overline{B_{g'(x_i)}(y_i, 1/3\lambda_i)}$  is a topological product with an interval and the distance function from  $y_i$  is the projection mapping of this product structure.
  - (c)  $B_{\lambda_i^2 g}(y_i, 2)$  is within  $\hat{\epsilon}(a_1)$  of a 2-dimensional Alexandrov ball of  $B(\bar{y}_i, 2)$  curvature  $\geq -1$ . Every ball  $B(\bar{z}, 1)$  centered at a point of  $B(\bar{y}_i, 1)$  has area  $\geq a$ . Lastly, the distance from  $\bar{y}_i$  is the projection mapping of a product structure on  $B(\bar{y}_i, 2) \setminus \overline{B(\bar{y}_i, 1/3)}$ .
2.  $A$  is not contained in any  $V(x)$  for any point  $x$  near a 1-dimensional endpoint and for every point  $y \in A$  the ball  $B_{g'(y)}(y, 1)$  is within  $\hat{\epsilon}$  of a 2-dimensional ball  $B(\bar{y}, 1)$  of curvature  $\geq -1$  and area  $\geq a$ .

**Definition 12.14.** We call a component of  $M \setminus U_1'$  satisfying Conclusion 2 above a component close to a 2-dimensional Alexandrov space.

### 12.1.2 Compact submanifolds $W_1$ and $W_2$

Let  $A$  be a component of  $M \setminus U_1''$  that is close to a 2-dimensional Alexandrov space. We shall expand  $A$  to a slightly larger compact submanifold denoted  $\hat{A}$ . Let  $U(x_{\mathcal{E}})$  be a neighborhood of an end of  $U_1''$  whose closure meets  $A$ , and let  $p(x_{\mathcal{E}}): U(x_{\mathcal{E}}) \rightarrow J(x_{\mathcal{E}})$  be its  $4\beta$ -interval product structure. Recall that  $J(x_{\mathcal{E}}) = (-1/250, 1/250)$ , and suppose that  $A$  meets the closure of the negative end of this interval. We take any cross-section  $\Sigma(\mathcal{E}) = p_{x_{\mathcal{E}}}^{-1}(-1/300)$ . This surface is either a 2-sphere or a 2-torus and the region between it and the boundary component of  $A$  in the closure of  $U(x_{\mathcal{E}})$  is a product. We form  $\hat{A}$  by adding these product regions, one for each boundary component of  $A$ , to  $A$ . The region  $p_{x_{\mathcal{E}}}^{-1}(t - 10^{-4}, t + 10^{-4})$  is a collar neighborhood of  $\Sigma(\mathcal{E})$  in  $M$ . We set  $C(\hat{A})$  equal to the union of  $\hat{A}$  with the collar neighborhoods of each of its boundary components. Since the neighborhoods  $U(x_{\mathcal{E}})$  have width at least  $1/250$  in the metric used to define them, if  $A, A'$  are distinct connected components of  $M \setminus U_1''$  of the type under consideration here, then the closure  $\overline{C}(\hat{A})$  of  $C(\hat{A})$  and the closure  $\overline{C}(\hat{A}')$  of  $C(\hat{A}')$  are disjoint.

**Claim 12.15.** For no point  $y \in \overline{C}(\hat{A})$  is  $B_{g'(y)}(y, 1)$  within  $\beta$  of an interval. In particular, for every point  $y \in \overline{C}(\hat{A})$  the ball  $B_{g'(y)}(y, 1)$  is within  $\hat{\epsilon}$  of a 2-dimensional Alexandrov ball of curvature  $\geq -1$  and area  $\geq a$ .

*Proof.* Fix  $y \in \overline{C}(\hat{A})$ . Then  $y$  is within  $(1/1000) + (1/10,000)$  of  $A$  in the metric used to define  $U(x_{\mathcal{E}})$ , and hence, by Lemma 6.1, is within  $(1/900)$  of  $A$  in the metric  $g'(y)$ . Hence by Proposition 12.7  $y \notin X_1$ . This shows that the manifold  $\overline{C}(\hat{A})$  is disjoint from  $X_1$ . Thus, if  $B_{h(y)}(y, 1)$  is within  $\beta$  of an interval then  $y$  is within  $\beta$  of a point  $\bar{y}$  which is within  $1/25$  of its endpoint. But in this case the distance from  $y$  to the complement of  $V_0(y)$ , when measured using  $g'(y)$  is at least .01 and hence  $V_0(y)$  would meet  $A$ , which is a contradiction. This proves the first statement. The second follows immediately from this and the dichotomy set up in Section 11.8.1.  $\square$

In particular, the conclusion of Case 2 of Theorem 11.46 applies to  $\overline{C}(\hat{A})$  to give a covering of it by the four types of metric balls listed in that theorem.

**Claim 12.16.** *Any  $\epsilon$ -solid torus, any  $\epsilon$ -solid cylinder, and any 3-ball  $B$  near a 2-dimensional corner that has a point within  $2r_0$  of  $\hat{A}$  (with the distance measured by the metric used to define the element in question) is contained in  $C(\hat{A})$ .*

*Proof.* This is immediate from the fact that  $r_0$  is less than  $10^{-6}$  and Lemma 6.1.  $\square$

Now consider one of the  $V_0(x_i)$ ,  $1 \leq i \leq \ell$ , containing complementary components of  $U_1''$ . In this case we set  $\hat{A}_i = \overline{B_{\lambda_i^2 g}(y_i, 1/3)}$ . We say that  $\hat{A}_i$  is a *component near a 1-dimensional endpoint*. The open set  $U_1''$  contains  $\partial V_0(x_i)$  and this boundary is parallel to the fibers of  $U_1''$ . Since  $d(y_i, \cdot)$  is regular on  $V_0(x_i) \setminus \hat{A}_i$ , this region is a product region. Also,  $\partial V_0(x_i) \subset U_1''$  and is isotopic in  $U_1''$  to a fiber of its fibration structure. Thus, it follows that every component of  $\cup_{i=1}^{\ell} (V_0(x_i) \setminus \hat{A}_i) \cup U_1''$  satisfies Condition 1 in Proposition 12.12.

In this case we set  $C(\hat{A}_i)$  equal to  $B_{\lambda_i^2 g}(y_i, 1)$ . In this case for any point  $y \in C(\hat{A}_i)$  we see that  $B_{\lambda_i^2 g}(y, 1)$  is within  $\hat{\epsilon}(a_1)$  of 2-dimensional Alexandrov ball of curvature  $\geq -1$  and area  $\geq a_1$ . In particular, the conclusion of Case 2 of Theorem 11.46 applies to  $\overline{C}(\hat{A}_i)$  to give a covering by the four types of metric balls listed in that theorem, when we use the metric  $\lambda_i^2 g$  at each point of  $C(\hat{A}_i)$ . As before, any  $\epsilon$ -solid torus, any  $\epsilon$ -solid cylinder, and any 3-ball near a 2-dimensional corner (each of these defined using the metric  $\lambda_i^2 g$ ) that has a point within  $r_0$  of  $\hat{A}_i$  is contained in  $C(\hat{A}_i)$ . The  $C(\hat{A}_i)$  are pairwise disjoint and are also disjoint from the  $\hat{A}$  associated to components of  $M \setminus U_1''$  near 2-dimensional Alexandrov spaces.

**Definition 12.17.** We define  $W_2$  to be the (disjoint) union of the  $\hat{A}$ , one for each complementary component  $A$  for  $M \setminus U_1''$  that is near a 2-dimensional Alexandrov space and the  $\hat{A}_i$ ,  $1 \leq i \leq \ell$ . **At this point we shift notation and use the symbol  $\hat{A}$  to refer to any component of  $W_2$ .** We set  $C(W_2)$  equal to the union of the  $C(\hat{A})$  as  $A$  ranges over the connected components of  $W_2$ , and  $\overline{C}(W_2)$  denotes the closure of  $C(W_2)$ . We define  $W_1$  to be the complement of the relative interior of  $W_2$  in  $M$ . Then  $W_1$  and  $W_2$  are compact manifolds with  $W_1 \cap W_2 = \partial W_2$  which in turn is the union of those components of  $\partial W_1$  that are not boundary components of  $M$ .

We define a metric  $h(x)$  on  $\overline{C}(W_2)$  as follows. For  $x \in \overline{C}(\hat{A})$  with  $A$  being a component close to a 2-dimensional we set  $h(x) = g'(x)$ . For  $x \in \overline{C}(\hat{A}_i)$  with  $A \subset V_0(x_i)$ ,  $1 \leq i \leq \ell$ , we set  $h(x) = \lambda_i^2 g$  where  $\lambda_i$  is the constant associated to this component by Proposition 12.13. Since  $\hat{\epsilon} < \hat{\epsilon}(a)$ , we see that every point  $y \in C(W_2)$  has the property that  $B_{h(y)}(y, 1)$  is within  $\hat{\epsilon}(a)$  of a 2-dimensional ball  $B(\bar{y}, 1)$  of curvature  $\geq -1$  and area  $\geq a$ .

**Proposition 12.18.** *Every component of  $W_1$  is one of the following:*

1. a  $T^2$ -bundle or an  $S^2$ -bundle over either the circle or a closed interval,
2. a twisted  $I$ -bundle over the Klein bottle,
3. a compact solid torus, a compact 3-ball, or the complement of an open 3-ball in  $\mathbb{R}P^3$ , or

4. the union of two twisted  $I$ -bundles over the Klein bottle along their common boundary.

*Proof.* This follows from Proposition 12.12, and the fact that the differences between  $A$  and  $\hat{A}$  are collar neighborhoods of the boundary.  $\square$

The following lemma gives the structure of  $W_2$  near each of its boundary components.

**Lemma 12.19.** *Let  $\Sigma$  be a boundary component of a connected component  $\hat{A}$  of  $W_2$ . Then there is a point  $x \in W_2$  such that the following hold:*

1.  $B_{h(x)}(x, 1)$  is within  $\hat{\epsilon} < \hat{\epsilon}(a)$  of a 2-dimensional Alexandrov ball  $B(\bar{x}, 1)$  of curvature  $\geq -1$  and area  $\geq a$ .
2.  $B_{h(x)}(x, 1/2)$  contains all points  $y \in \hat{A}$  within distance  $10^{-5}$  of  $\Sigma$  in the metric  $h(x)$ .
3. If  $\hat{A}$  is close to a 2-dimensional Alexandrov space then  $d_{h(x)}(x, \Sigma) < 2 \times 10^{-4}$ .
4. There are  $0 < a < b < 1$  with  $b - a = 1/8000$  such that, setting  $N(\Sigma)$  equal to the connected component of  $d_{h(x)}(x, \cdot)^{-1}(a, b)$  that contains  $\Sigma$ , the function  $d_{h(x)}(x, \cdot)$  is regular on  $N(\Sigma)$  and defines the projection map of a topological product structure  $N(\Sigma) \rightarrow (a, b)$ .
5. The boundary component  $\Sigma$  is isotopic in  $N(\Sigma)$  to the fiber  $d_{h(x)}(x, \cdot)^{-1}(t)$ , for every  $t \in (a, b)$ .
6. There is a connected component  $\bar{N}(\Sigma)$  of  $d(\bar{x}, \cdot)^{-1}(a, b) \subset B(\bar{x}, 1)$  that is within  $4\hat{\epsilon}$  of  $N(\Sigma)$  and on which  $d(\bar{x}, \cdot)$  is regular and defines the projection mapping of a topological product structure  $\bar{N}(\Sigma) \rightarrow (a, b)$ .

*Proof.* We denote by  $\hat{A}$  the component that has  $\Sigma$  as a boundary component. First suppose that the corresponding component  $A$  of  $M \setminus U_1''$  is contained in one of the  $V_0(x_i)$ ,  $1 \leq i \leq k$ . Then the result is immediate from Proposition 12.13 using the point  $y_i$ .

Now suppose that  $\hat{A}$  corresponds to a component  $A$  of  $M \setminus U_1''$  that is close to a 2-dimensional Alexandrov space. Let  $U(x_{\mathcal{E}})$  be the neighborhood of an end of  $U_1''$  that contains  $\Sigma$ . In this case we choose a point  $x$  in the component of the frontier of  $U(x_{\mathcal{E}})$  that is not contained in  $U_1''$ . Then the distance from  $x$  to  $\Sigma$  is within  $2\epsilon'$  of  $1/8000$  when measured using  $d_{g'(x_{\mathcal{E}})}$ , and  $\Sigma$  has diameter at most  $\epsilon'$  in this metric. Since the distance  $d_{g'(x_{\mathcal{E}})}(x_{\mathcal{E}}, x)$  is within  $2\hat{\epsilon}$  of  $1/8000$ , which is between  $(1.1)/8000$  and  $(0.9)/8000$  by Lemma 6.1, the ratio of  $d_{h(x)} = d_{g'(x)}$  and  $d_{g'(x_{\mathcal{E}})}$  is between  $26/25$  and  $25/26$ . Since  $\epsilon' < 10^{-6}$ , it follows that the distance between any point of  $\Sigma$  and  $x$ , measured using  $h(x) = g'(x)$ , is between  $1/7000$  and  $1/9000$ . We denote the distance from  $\Sigma$  to  $x$  by  $d$ , set  $a = d - (1/16,000)$ , and set  $b = d + (1/16,000)$ . We set  $N(\Sigma)$  equal to the connected component of  $d_{h(x)}(x, \cdot)^{-1}(a, b)$  containing  $\Sigma$ . Then  $N(\Sigma)$  contains the neighborhood of size  $10^{-5}$  (when measured in  $h(x)$ ) about  $\Sigma$ . Notice

that  $N(\Sigma)$  is contained in  $U(x_{\mathcal{E}})$  and the distance, measured using  $g'(x_{\mathcal{E}})$ , from  $x$  to any point of  $N(\Sigma)$  is greater than  $2 \cdot 10^{-5}$ . It then follows from Lemma 11.42 that  $d_{h(x)}(x, \cdot)$  is regular on  $N(\Sigma)$  and that the fibers  $d_{h(x)}(x, \cdot)^{-1}(t)$  are isotopic in  $N(\Sigma)$  to  $\Sigma$  for all  $t \in (d - (1/16, 000), d + (1/16, 000))$ .

We know that  $B_{h(x)}(x, 1)$  is within  $\hat{\epsilon}$  of a 2-dimensional Alexandrov ball  $B(\bar{x}, 1)$  of curvature  $\geq -1$  and area  $\geq a$ . Since  $B_{g'(x_{\mathcal{E}})}(x_{\mathcal{E}}, 1)$  is within  $\beta$  of an interval, it follows that the ball  $B(\bar{x}, 1/2)$  is within  $5(\hat{\epsilon} + \beta)$  of an interval. Thus,  $d(\bar{x}, \cdot)$  is regular on  $B(\bar{x}, 1/2) \setminus B(\bar{x}, 10^{-4})$ . Hence, this function is regular on  $\bar{N}(\Sigma)$  which is defined to be the connected component of  $d(\bar{x}, \cdot)^{-1}(a, b)$  within  $4\hat{\epsilon}$  of  $N(\Sigma)$ . This completes the proof of the lemma in this case.  $\square$

## 12.2 A covering of $\bar{C}(W_2)$

According to Theorem 11.46 and the remark before Definition 12.17,  $\bar{C}(W_2)$  has an opening covering consisting of  $U_{2,\text{gen}}$ ,  $\epsilon$ -solid tori, cores of  $\epsilon$ -solid cylinders, and 3-balls near 2-dimensional corners. Furthermore, since  $r_0 \leq 10^{-6}$  any  $\epsilon$ -solid torus,  $\epsilon$ -solid cylinder or 3-ball near a 2-dimensional corner that meets the  $r_0$ -neighborhood of  $W_2$  (measured in the metric used to define the element) is contained in  $C(W_2)$ . Of course, by compactness we need only finitely many such open sets to cover  $\bar{C}(W_2)$ .

### 12.2.1 Seifert fibrations

It will be important in the following to know that any compact subset contained in the union of  $U_{2,\text{gen}}$  and  $\epsilon$ -solid tori is in fact contained in the total space of a Seifert fibration.

**Proposition 12.20.** *Suppose that  $X$  is any compact subset of the union of  $U_{2,\text{gen}}$  and a collection of  $\epsilon$ -solid tori. Then there is an open subset  $Z$  containing  $X$  that is the total space of a Seifert fibration. There is a disjoint union of solid tori in  $X$ , each of the solid tori is saturated under the Seifert fibration and is an unknotted solid torus in an  $\epsilon$ -solid torus neighborhood,  $B_{h(z_i)}(z_i, r_i/4)$ . The complement of these solid tori in  $Z$  is saturated under the Seifert fibration and is contained in  $U_{2,\text{gen}}$ , and the restriction of the Seifert fibration to the complement is an  $S^1$ -fibration with fibers within  $\epsilon'$  of vertical with respect to  $S^1$ -product structures with  $\epsilon$ -control.*

*Proof.*  $X$  is contained in the union of  $U_{2,\text{gen}}$  and a finite number of  $\epsilon$ -solid tori neighborhoods  $B_{h(z_i)}(z_i, r_i/4)$ . Suppose two of these solid tori  $B_1$  and  $B_2$  meet. We number things so that the unrescaled radius of  $B_1$  is great than or equal to that of  $B_2$ . Then  $B_2$  is contained in the metric ball with center  $x_1$  and radius  $3r_1/4$  (measured in the metric  $h(x_1)$ ). Of course  $U_{2,\text{gen}}$  contains the union of the  $B_{h(z_i)}(z_i, 7r_i/8) \setminus B_{h(z_i)}(z_i, r_i/8)$ . Hence, the expense of expanding these metric balls to have radius  $3r_i/4$  we can assume that the  $\epsilon$ -solid tori  $B_{h(z_i)}(z_i, r_i/4)$  are disjoint. Let  $X'$  be the complement in  $X$  of the  $B_{h(z_i)}(z_i, r_i/4)$  and let  $X_1$  be the union of  $X'$  with the  $\overline{B_{h(z_i)}(z_i, 3r_i/4)} \setminus B_{h(z_i)}(z_i, r_i/4)$ . This is a compact subset of  $U_{2,\text{gen}}$ , and hence by Proposition 11.8 it is contained in an open subset  $U_0 \subset U_{2,\text{gen}}$  that is fibered by circles that are within  $\epsilon'$  of vertical in the  $S^1$ -product structures.

By Corollary 11.23 this fibration extends to a Seifert fibration over the union of  $U_0$  with the  $B_{h(z_i)}(z_i, r_i/4)$  with at most one exceptional fiber in each of these balls. This is the required Seifert fibration.  $\square$

**Corollary 12.21.** *Let  $\nu_\xi(\tilde{\gamma})$  be an  $\epsilon$ -solid cylinder, and let  $D_0$  be a spanning disk for its core  $\nu_{\xi^2}(\tilde{\gamma})$ . Then  $D_0$  is not contained in the union of  $U_{2,\text{gen}}$  and  $\epsilon$ -solid tori.*

*Proof.* First, let us suppose that the disk  $D_0$  is contained in  $U_{2,\text{gen}}$ . Then  $U_{2,\text{gen}}$  contains the closure of  $X = D_0 \cup \nu_\xi(\tilde{\gamma}) \setminus \nu_{\xi^2/2}(\tilde{\gamma})$ . There is an  $S^1$ -fibration structure on an open subset containing  $X$  with the property that each fiber is within  $\epsilon'$  of any  $S^1$ -product structure centered at any point of  $X$ . This implies that the boundary of  $D_0$  is isotopic to a fiber of this  $S^1$ -fibration. But the only  $S^1$ -fibrations whose generic fibers are homotopically trivial have total space  $S^3$ . But this is ruled out since no component of  $M$  is homeomorphic to  $S^3$ . This proves that  $D_0$  is not contained in  $U_{2,\text{gen}}$ .

Now suppose that  $D_0$  is contained in the union of  $U_{2,\text{gen}}$  and a collection of  $\epsilon$ -solid tori. By the above,  $D_0$  meets an  $\epsilon$ -solid torus  $T = B_{h(z)}(z, r(z)/4)$ .

**Claim 12.22.**  $r(z) \leq 50\xi^2 r_1 s_1$ .

Given this claim, it follows that any  $\epsilon$ -solid torus  $T$  that meets  $D_0$  is disjoint from  $A = \nu_\xi(\tilde{\gamma}) \setminus \nu_{51\xi^2}(\tilde{\gamma})$ . Thus, we can cover  $D_0 \cup (\nu_\xi(\tilde{\gamma}) \setminus \nu_{\xi^2}(\tilde{\gamma}))$  by  $\epsilon$ -solid tori and  $U_{2,\text{gen}}$  in such a way that  $A$  is disjoint from all the  $\epsilon$ -solid tori in the covering. Then there is a Seifert fibration structure on an open set containing this union, and the level circles of  $\nu_\xi(\tilde{\gamma}) \setminus \nu_{21\xi^2}(\tilde{\gamma})$  are homotopic to a generic fiber of this Seifert fibration. As before, this is only possible if the component of the total space of the Seifert fibration is a closed 3-manifold is a 3-dimensional spherical space form.

It remains to prove the claim, which follows immediately from the next claim.

**Claim 12.23.** *Suppose that  $T = B_{h(z)}(z, r(z)/4)$  contains a point of  $\nu_\xi(\tilde{\gamma})$  at distance  $d$  (in the metric used to define  $\nu_\xi(\tilde{\gamma})$ ) from  $\tilde{\gamma}$ . Then  $r(z) < 20d + \xi^2 r_1 s_1$ .*

*Proof.* Suppose that  $r(z) \geq 20d + \xi^2 r_1 s_1$ . Let  $h(x)$  be the metric used to define  $\nu_\xi(\tilde{\gamma})$ . First notice that there is a constant  $R$  such that  $h(z) = R^2 h(x)$  and by Lemma 6.1 we have  $(1.1)^{-1} < R < (1.1)$ . Thus, if  $T$  meets  $\nu = \nu_\xi(\tilde{\gamma})$  and contains a point at distance  $d$  (measured in the metric used to define  $\nu$ ) from  $\tilde{\gamma}$ , then  $B_{h(z)}(z, r(z)/4 + (1.1)d)$  contains a point  $q \in \tilde{\gamma}$ . According to Part 5 of Lemma 11.31, there is a point  $\bar{q} \in \partial B(\bar{x}, 1)$  within  $\xi^2 r_1 s_1 / 100$  of  $q$  and there is a point  $q' \in B_{h(z)}(z, r(z)/4 + (1.1)d + 3\xi^2 r_1 s_1 / 100)$  within  $\hat{\epsilon}$  of  $\bar{q}$ . Under our hypothesis  $r(z)/4 + (1.1)d + 3\xi^2 r_1 s_1 / 100 < r(z)/3$ , so that  $q' \in B_{h(z)}(z, r(z)/3)$ . Now  $(1/r(z))B_{h(z)}(z, r(z))$  is within  $\hat{\epsilon}/r(z)$  of  $(1/r(z))B(\bar{z}, r(z))$  which is within  $\mu$  of a circular cone  $C$  with cone point  $\bar{z}$ . The point  $q'$  is within  $(\hat{\epsilon}/r(z)) + \mu$  of a point  $\bar{q}' \in C$  with  $d(\bar{z}, \bar{q}') < (0.34)$ . Hence,  $(1/r(z))B_{h(z)}(q', r(z)/2)$  is within  $4[(\hat{\epsilon}/r(z)) + \mu]$  of  $B(\bar{q}', 1/2) \subset C$ . On the other hand,  $(1/r(z))B_{h(z)}(q', r(z)/2) = (1/r(z))B_{R^2 h(x)}(q', r(z)/2) = (R/r(z))B_{h(x)}(q', r(z)/2R)$ , and this ball is within  $4R\hat{\epsilon}/r(z)$  of  $(R/r(z))B(\bar{q}, r(z)/2R)$ . It follows that  $(1/r(z))B(\bar{q}', r(z)/2)$  and the ball  $(R/r(z))B(\bar{q}, r(z)/2R)$  are within  $4((R+1)\hat{\epsilon}/r(z) + \mu)$  of each other. But we have



$\mu < \delta(a'(a))/8$ ,  $\hat{\epsilon}/r(z) < (r_2/20)\delta(a'(a))/r(z)$ ,  $r(z) \geq r_2$ , and  $R \leq 1.1$ . This implies that these two balls are within  $\delta(a'(a))$  of each other in the Gromov-Hausdorff distance. By construction  $\bar{q} \in \partial B(\bar{q}, 1/2)$ , and  $C$ , being a circular cone, has no boundary. This contradicts Lemma 10.7.  $\square$

This completes the proof of the corollary.  $\square$

### 12.3 Fixing the 3-balls

**Lemma 12.24.** *There is a finite set of balls near 2-dimensional corners,*

$$B_1 = B_{h(x_1)}(w_1, r(w_1)/8), \dots, B_N = B_{h(x_N)}(w_N, r(w_N)/8),$$

*each meeting  $\overline{C}(W_2)$ , such that the following hold:*

1. *The closures of the  $B_{h(x_i)}(w_i, 3r(w_i)/16)$  are disjoint.*
2. *Every ball near a 2-dimensional corner,  $B_{h(x)}(w, r(w)/8)$  that meets  $\overline{C}(W_2)$ , is contained in one of the  $B_{h(x_i)}(w_i, 7r(w_i)/8)$ ,  $i = 1, \dots, N$ .*
3. *If  $B = B_{h(x)}(w, r(w)/8)$  is a 3-ball near a 2-dimensional corner, then  $B$  is contained in the union of the  $B_i$ ,  $U_{2,\text{gen}}$ , and the union of the cores of  $\epsilon$ -solid cylinders.*

*Proof.* Among all balls  $B_{h(x_i)}(w_i, r(w_i)/8)$  in  $M$  near 2-dimensional corners that meet  $\overline{C}(W_2)$ , choose one whose rescaled radius is at least  $(0.9)$  times the supremum of the rescaled radii of all such balls. Call this  $B_1 = B_{h(x_1)}(w_1, r(w_1)/8)$ . Now among all balls  $B_{h(x)}(w, r(w)/8)$  near 2-dimensional corners meeting  $\overline{C}(W_2)$  with the property that the closure of  $B_{h(x)}(w, 3r(w)/16)$  is disjoint from the closure of  $B_{h(x_1)}(w_1, 3r(w_1)/16)$  choose one whose rescaled radius is at least  $(0.9)$  times the supremum of the rescaled radii of all such balls. Call this  $B_2$ . Continue in this fashion constructing  $B_1, B_2, \dots$ . First notice that since the rescaled radii of all balls under consideration are at least  $r_1$  and since  $p$  is bounded on the compact manifold  $M$ , this process must terminate after a finite number of steps, say after  $B_N$ . Now suppose that  $B = B_{h(x)}(w, r(w)/8)$  is a ball near a 2-dimensional corner that meets  $\overline{C}(W_2)$ . Then the closure of the ball with the same center and with radius  $3r(w)/16$  must meet at least one of the closures of the  $B_{h(x_j)}(w_j, 3r(w_j)/16)$ . Take the smallest index  $j$  for which this is true. Then by the inductive construction of  $B_i$ , we have  $(0.9)r(w) \leq r(w_j)$ . On the other hand, since the balls have closures that meet, since  $r_1, s_1, \xi, \hat{\epsilon} \leq 10^{-6}$ , and since  $d_{h(x)}(x, y) < 2\hat{\epsilon} + \xi^2 r_1 s_1 / 100$ , it follows from Lemma 6.1 that  $p^{-1}(x)/p^{-1}(x_j) \leq 1.01$ . It then follows that  $B_{h(x)}(w, 3r(w)/16) \subset B_{h(x_j)}(w_j, 7r(w_j)/8)$  and hence by Corollary 11.39 that  $B$  is contained in  $B_j$  and the union of  $U_{2,\text{gen}}$  and the union of cores of  $\epsilon$ -solid cylinders. This shows that the collection  $\{B_1, \dots, B_N\}$  satisfies the conclusion of the lemma.  $\square$

For the rest of this section we fix a set of 3-balls  $B_i = B_{h(x_i)}(w_i, r(w_i)/8)$ ,  $1 \leq i \leq N$ , near 2-dimensional corner points satisfying the conclusion of the previous lemma.

### 12.3.1 Attaching $\epsilon$ -solid cylinders to each 3-ball

According to Corollary 11.39 for each ball  $B_i$  we can choose two disjoint  $\epsilon$ -solid cylinders  $\nu(i)^\pm$  with width factor  $(0.9)\xi$  such that the boundary sphere of  $B_i$  passes through the central point of each of the defining geodesics of the  $\nu(i)^\pm$ , and such that every point of the boundary sphere not contained in the cores of these two  $\epsilon$ -solid cylinders is contained in  $U_{2,\text{gen}}$ . Since the closures of the balls with the same centers and radii  $3r(w_i)/16$  are disjoint, making these choices results in a pairwise disjoint collection of  $\epsilon$ -solid cylinders. The  $\nu(i)^\pm$  are called *the  $\epsilon$ -solid cylinders bisected by  $S_i$* , the metric sphere bounding  $B_i$ , see FIG. 9.

**Definition 12.25.** For  $1 \leq i \leq N$  we define

$$\tilde{B}_i = B_i \cup \nu(i)^+ \cup \nu(i)^-.$$

The  $\tilde{B}_i$  are fixed for the rest of the argument.

Since the  $B_{h(x_i)}(w_i, 3r(w_i)/16)$  are disjoint, the following is clear from Lemma 6.1.

**Claim 12.26.** *There is no  $\epsilon$ -solid cylinder that meets two of the  $\tilde{B}_i$ . Furthermore, each  $\tilde{B}_i$  is contained in the  $r_0$ -neighborhood of  $W_2$  using the metric  $h(x_i)$ .*

**Lemma 12.27.** *Suppose that an  $\epsilon$ -solid cylinder,  $\nu$ , meets  $\tilde{B}_i$  for some  $i$ . Then the intersection of the defining geodesic  $\tilde{\gamma}$  for  $\nu$  with  $\tilde{B}_i$  is contained in the union of the cores of  $\nu(i)^\pm$  and the ball  $B_{h(x_i)}(w_i, (r(w_i)/8) - (r_1 s_1/18))$ . Also, the intersection of the core of  $\nu$  with  $\tilde{B}_i$  is contained in  $\nu(i)^+ \cup B_{h(x_i)}(w_i, t(w_i)) \cup \nu(i)^-$ , where  $t(w_i) = (r(w_i)/8) - (r_1 s_1/20)$ .*

*Proof.* Clearly, the second statement follows from the first. We establish the first. By Lemma 11.34 the intersection of  $\tilde{\gamma}$  with  $\nu(i)^\pm$  is an interval with each endpoint either being an endpoint of  $\tilde{\gamma}$  or an intersection of  $\tilde{\gamma}$  with an end of  $\nu(i)^\pm$ . Also, this intersection is contained in the core of  $\nu(i)^\pm$ . The result will follow once we show that the intersection of  $\tilde{\gamma}$  with the annular region  $B_i \setminus B_{h(x_i)}(w_i, (r(w_i)/8) - (r_1 s_1/18))$  is contained in the union of the cores of  $\nu(i)^\pm$ . If  $\tilde{\gamma}$  meets this annular region, then according to Lemma 11.41 it is within  $\xi^2 r_1 s_1/50$  of the boundary of the associated 2-dimensional Alexandrov space  $B(\bar{x}, 1)$ . On the other hand, since the defining geodesics for  $\nu(i)^+$  and  $\nu(i)^-$  are within  $\hat{\epsilon}$  of  $\mu$ -approximations to  $\partial B(\bar{x}, 1)$  of length  $r_1 s_1/4$  and midpoint at distance  $r(w_i)/8$  from  $\bar{x}$ , it follows that the union of the cores of  $\nu(i)^+$  and  $\nu(i)^-$  contains the middle sub-geodesic of  $\tilde{\gamma}$  of length  $r_1 s_1/8$  and hence contains all points of the annular region within  $\xi^2 r_1 s_1/9$  of  $\partial B(\bar{x}, 1)$ , and hence contains the intersection of  $\tilde{\gamma}$  with this annular region.  $\square$

## 12.4 $\epsilon$ -Chains

At this point we must introduce the notion of chains of  $\epsilon$ -solid cylinders and 3-balls near 2-dimensional corners.

### 12.4.1 Good intersections of $\epsilon$ -solid cylinders

**Definition 12.28.** Suppose that for  $i = 1, 2$  we have  $\epsilon$ -solid cylinders  $\nu(i) = \nu_{c_i \xi, [a_i, b_i]}(\tilde{\gamma}_i) \subset B_{h(x_i)}(y_i, 1)$ . (Recall that implicitly  $c_i \in [1/10, 1]$  and  $b_i - a_i \geq \ell_i/5$ ,  $y_i$  is the control point for  $\nu(i)$ , and  $\ell_i$  is the length of  $\tilde{\gamma}_i$  with respect to the metric  $h(x_i)$ .) We say that the  $\nu(2)$  has *good intersection with  $\nu(1)$*  if the following hold with appropriate orientations of the  $\tilde{\gamma}_i$ :

1. There is a point in the negative end of  $\nu(2)$  that is contained in

$$f_{\tilde{\gamma}_1}^{-1}(b_1 - (0.009)\ell_1, b_1 - (0.006)\ell_1)$$

in  $\nu(1)$ , and the positive end of  $\nu(2)$  is at distance at least  $(0.1)\ell_2$  from  $\nu(1)$  when measured in the metric  $h(y_2)$ .

2.  $c_1\ell_1 p(x_1)$  is either at least  $(1.1)c_2\ell_2 p(x_2)$  or is at most  $(1.1)^{-1}c_2\ell_2 p(x_2)$ .

**Lemma 12.29.** *With the notation above, suppose that for  $i = 1, 2$  the  $\epsilon$ -solid cylinders  $\nu(i) = \nu_{c_i \xi, [a_i, b_i]}(\tilde{\gamma}_i)$  have the property that  $\nu(2)$  has good intersection with  $\nu(1)$ . Then the closure of that intersection is homeomorphic to a closed 3-ball. If*

$$c_1\ell_1 p(x_1) < c_2\ell_2 p(x_2), \tag{12.1}$$

*then that 3-ball meets the boundary of the closure  $\bar{\nu}(2)$  of  $\nu(2)$  in a 2-disk contained in the negative end of  $\bar{\nu}(2)$  and the rest of the boundary consists of an annulus in the side of  $\bar{\nu}(1)$  together with the positive end of  $\bar{\nu}(1)$ . If the reverse inequality holds in 12.1, the similar statements hold with the roles of  $\bar{\nu}(1)$  and  $\bar{\nu}(2)$  and ‘positive’ and ‘negative’ reversed. See FIG. 10.*

*Proof.* We suppose that Inequality 12.1 holds. It follows from Lemma 10.27 that the sides of  $\bar{\nu}(1)$  and of  $\bar{\nu}(2)$  do not intersect and in fact the side of  $\bar{\nu}(2)$  is disjoint from  $\bar{\nu}(1)$ . Thus, the intersection of  $\bar{\nu}(1)$  and  $\partial\bar{\nu}(2)$  is contained in the negative end of  $\bar{\nu}(2)$ . By Part 3 of Lemma 11.34, this intersection is a 2-disk. Hence, it cuts off a 3-ball in  $\bar{\nu}(1)$ .

The other case is analogous. □

**Corollary 12.30.** *With notation and assumptions above, suppose that Inequality 12.1 holds. Then the boundary of  $\bar{\nu}(1) \cup \bar{\nu}(2)$  consists of the union of two subsets: (i) the disjoint union of two 2-disks: the negative end of  $\bar{\nu}(1)$  and the positive end of  $\bar{\nu}(2)$ , and (ii) an annulus  $E$ . These two subsets are glued together along their boundaries. The annulus  $E$  consists of the union of three annuli glued together along their boundaries. The first is the intersection of the side of  $\bar{\nu}(1)$  with the complement of the interior of  $\bar{\nu}(2)$ . The second is the negative end of  $\bar{\nu}(2)$  minus its intersection with the interior of  $\bar{\nu}(1)$  and the third is the side of  $\bar{\nu}(2)$ . If the opposite inequality to Inequality (12.1) holds, then there are similar statements with the roles of  $\bar{\nu}(1)$  and  $\bar{\nu}(2)$  and ‘positive’ and ‘negative’ reversed.*

### 12.4.2 Chains of $\epsilon$ -solid cylinders

Now suppose that we have a sequence of  $\epsilon$ -solid cylinders  $\{\nu(1), \dots, \nu(k)\}$ , with  $\nu(i) = \nu_{c_i \xi, [a_i, b_i]}(\tilde{\gamma}_i)$  with the geodesics  $\tilde{\gamma}_i$  oriented. We say that these form a *linear chain of  $\epsilon$ -solid cylinders* if:

1. For each  $1 \leq i < k$  the  $\epsilon$ -solid cylinder  $\nu(i+1)$  has good intersection with  $\nu(i)$ .
2. If  $\nu(i) \cap \nu(j) \neq \emptyset$  for some  $i \neq j$ , then  $|i - j| = 1$ .

In addition to linear chains there are circular chains.

**Definition 12.31.** A *circular chain of  $\epsilon$ -solid cylinders* is a sequence  $\{\nu(1), \dots, \nu(k)\}$  of  $\epsilon$ -solid cylinders, indexed by the integers modulo  $k$ , such that for each  $i$ ,  $1 \leq i \leq k$ , the  $\epsilon$ -solid cylinder  $\nu(i+1)$  has good intersection with  $\nu(i)$ , and for each  $i, j$  if  $\nu(i) \cap \nu(j) \neq \emptyset$  then  $j \equiv i-1, i$  or  $i+1 \pmod{k}$ .

**Lemma 12.32.** Suppose that  $\{\nu(1), \dots, \nu(k)\}$  is a linear chain of  $\epsilon$ -solid cylinders. Then  $\bar{\nu}(1) \cup \dots \cup \bar{\nu}(k)$  is homeomorphic to a 3-ball and its boundary is the union of the negative end of  $\bar{\nu}(1)$ , the positive end of  $\bar{\nu}(k)$  and an annulus  $E$ .

*Proof.* This is proved easily by induction. □

The same arguments establish the analogue for circular chains.

**Lemma 12.33.** Let  $\{\nu(1), \dots, \nu(k)\}$  be a circular chain of  $\epsilon$ -solid cylinders contained in  $M$ . Then  $\cup_i \bar{\nu}(i)$  is homeomorphic to a solid torus.

**Definition 12.34.** Suppose that  $\nu(1), \dots, \nu(k)$  is a linear chain of  $\epsilon$ -solid cylinders. The  $\nu(i)$  are the *elements* of the chain. The *extremal elements* are  $\nu(1)$  and  $\nu(k)$  and its *free ends* are the end of  $\nu(1)$  disjoint from  $\nu(2)$  and the end of  $\nu(k)$  disjoint from  $\nu(k-1)$ . For a chain  $C$  of  $\epsilon$ -solid cylinders, we denote by  $U(C)$  the union of the  $\epsilon$ -solid cylinders in  $C$ . The subset  $U(C)$  is also called the *total space of the chain*.

### 12.4.3 Definition of $\epsilon$ -chains and their topology

Now we are ready to construct chains made up of  $\epsilon$ -solid cylinders and the  $\tilde{B}_i$  which have been fixed earlier in the discuss (with good intersections).

**Definition 12.35.** A *linear  $\epsilon$ -chain* consists of an ordered set

$$\{C_1, \tilde{B}_{i_1}, C_2, \tilde{B}_{i_2}, \dots, \tilde{B}_{i_{k-1}}, C_k\}, \quad k \geq 1,$$

where:

1. Each  $C_j$ ,  $1 \leq j \leq k$ , is a linear chain of  $\epsilon$ -solid cylinders with good intersection.
2. For  $j \neq j'$  we have  $U(C_j) \cap U(C_{j'}) = \emptyset$ .
3. For each  $j$ ,  $1 \leq j \leq k-1$ , the ordered collection of  $\epsilon$ -solid cylinders

$$\{\nu(i_{j-1})^+, C_j, \nu(i_{j+1})^-\}$$

is a linear chain of  $\epsilon$ -solid cylinders with good intersection.

4. The  $\tilde{B}_{i_j}$  are distinct balls chosen from the  $\tilde{B}_1 \dots, \tilde{B}_N$ .
5. For every  $j < k$  the intersection of  $B_{i_j}$  with  $\cup_m U(C_m)$  is equal to the intersection of  $\nu(i_j)^+ \cup \nu(i_j)^-$  with  $B_{i_j}$ .

The *elements* of the linear  $\epsilon$ -chain are the  $\nu(i_j)^\pm, B_{i_j}$  and the elements of the  $C_j$ . The *free ends* of a linear  $\epsilon$ -chain  $\{C_1, \tilde{B}_{i_1}, \dots, \tilde{B}_{i_{k-1}}, C_k\}$  is the end of  $C_1$  disjoint from  $\tilde{B}_{i_1}$  and the end of  $C_k$  disjoint from  $\tilde{B}_{i_{k-1}}$ , and the extremal elements are the two  $\epsilon$ -solid cylinders containing the free ends.

An *circular  $\epsilon$ -chain* consists either (a) of an ordered set (up to cyclic permutation shifting by an even number of terms)  $\{C_1, \tilde{B}_{i_1}, \dots, C_k, \tilde{B}_{i_k}\}$  satisfying the conditions above except that in the third item the indices are taken modulo  $k$ , so that the end of  $C_1$  disjoint from  $\tilde{B}_{i_1}$  is  $\nu(i_k)^+$  or (b) of a circular chain of  $\epsilon$ -solid cylinders up to cyclic permutation. The *elements* of the circular  $\epsilon$ -chain are the  $\nu(i_j)^\pm, B_{i_j}$  and the elements of the  $C_j$ .

An  $\epsilon$ -chain is either a linear  $\epsilon$ -chain or a circular  $\epsilon$ -chain.

Given an  $\epsilon$ -chain  $\mathcal{C}$  we define the *total space*,  $U(\mathcal{C})$ , of the chain to be the union of the  $U(C_i)$  as  $C_i$  ranges over the chains of  $\epsilon$ -solid cylinders that are elements of  $\mathcal{C}$ , and the balls  $\tilde{B}_{i_j}$  that are elements of  $\mathcal{C}$ . See FIG. 11.

The next two lemmas describe the topology of  $\epsilon$ -chains.

**Lemma 12.36.** *Let  $\mathcal{C}$  be a linear  $\epsilon$ -chain. Then  $U(\mathcal{C})$  is homeomorphic to a 3-ball.*

*Proof.* Since each  $U(C_j)$  is homeomorphic to a 3-ball and the intersection of  $U(C_j)$  with the boundary of each of  $\tilde{B}_{i_{j-1}}$  and  $\tilde{B}_{i_j}$  is a 2-disk, the first statement is easily proved by induction.  $\square$

The same argument shows the following:

**Lemma 12.37.** *Let  $\mathcal{C}$  be a circular  $\epsilon$ -chain. Then  $U(\mathcal{C})$  is homeomorphic to a solid torus.*

## 12.5 Existence theorem for a complete set of $\epsilon$ -chains

Now we shall show that we can cover all of  $W_2$  by  $U_{2,\text{gen}}$ , a finite set of  $\epsilon$ -solid tori, and a finite disjoint collection of  $\epsilon$ -chains.

**Theorem 12.38.** *There are a finite number of  $\epsilon$ -chains  $\mathcal{C}_1, \dots, \mathcal{C}_K$  satisfying the following conditions:*

1.  $W_2$  is contained in the union of  $\cup_{i=1}^K U(\mathcal{C}_i)$ ,  $U_{2,\text{gen}}$ , and a finite collection of  $\epsilon$ -solid tori.
2. For  $i = 1, \dots, K$ , the  $\mathcal{C}_i$  are contained in  $C(W_2)$ .
3.  $U(\mathcal{C}_i) \cap U(\mathcal{C}_j) = \emptyset$  for all  $i \neq j$ .
4. The width factor in each  $\epsilon$ -solid cylinder element of each  $\mathcal{C}_i$  is between  $(0.7)\xi$  and  $(0.9)\xi$ .

5. The free ends of the  $\mathcal{C}_i$  are at distance greater than  $r_0$  from  $W_2$ .

*Proof.* The proof of this theorem takes up the entire subsection. Let us begin with some basic definitions in this context.

**Definition 12.39.** Let  $\mathcal{C}$  be an  $\epsilon$ -chain. We say that  $\mathcal{C}$  is *calibrated* if:

- (a)  $U(\mathcal{C}) \subset C(W_2)$ .
- (b) The width of every  $\epsilon$ -solid cylinder element of  $\mathcal{C}$  is between  $(0.7)\xi$  and  $(0.9)\xi$  and any extremal element of  $\mathcal{C}$  has width either  $(0.7)\xi$  or  $(0.9)\xi$ .
- (c) If  $\nu$  is an extremal element of  $\mathcal{C}$ , then either  $\nu$  is one of the  $\epsilon$ -solid cylinders  $\nu(j)^\pm$  bisected by one of the  $B_j$  or  $\nu = \nu_{c\xi, [-r_1 s_1/16, 0]}(\tilde{\gamma})$  and with the corresponding free end of  $\mathcal{C}$  contained in  $f_{\tilde{\gamma}}^{-1}(0)$ .

We say that a collection of  $\epsilon$ -chains is a *calibrated collection* if each individual  $\epsilon$ -chain in the collection is calibrated, if the total spaces of the  $\epsilon$ -chains in the collection are pairwise disjoint, and if for every  $i$  the three elements making up  $\tilde{B}_i$  are all elements of one of the  $\epsilon$ -chains.

**Claim 12.40.** *Let  $\nu$  be an  $\epsilon$ -solid cylinder with generating geodesic  $\tilde{\gamma}$ , and suppose that the core of  $\nu$  meets the total space of a calibrated  $\epsilon$ -chain  $\mathcal{C}$  and also meets its complement. Then the intersection of  $\tilde{\gamma}$  with  $\mathcal{C}$  consists of either one or two intervals and each endpoint of each interval is either contained in a free end of  $\mathcal{C}$  or is an endpoint of  $\tilde{\gamma}$ . Furthermore, the intersection of  $\tilde{\gamma}$  with  $\mathcal{C}$  is contained in the union of the cores of the  $\epsilon$ -solid cylinders in  $\mathcal{C}$  and the balls  $B_{h(x_i)}(w_i, (r(w_i)/8) - (r_1 s_1/18))$ . Lastly, if  $\tilde{\gamma}$  meets one of the  $\tilde{B}_i$  then its intersection with  $\tilde{B}_i$  is an interval with one endpoint in the free end of  $\nu^\pm(i)$  and the other an endpoint of  $\tilde{\gamma}$ .*

*Proof.* If  $\nu$  meets an  $\epsilon$ -solid torus  $\nu'$  and also meets its complement, then it follows from Lemma 11.34 that  $\tilde{\gamma} \cap \nu'$  is contained in the core of  $\nu'$  and one endpoint of intersection of  $\tilde{\gamma}$  with  $\nu'$  is a point in the core of an end of  $\nu'$ . Also, it follows from Lemma 11.41 that if  $\nu$  meets one of the  $B_j$ , then since it meets both  $B_j$  and its complement, the geodesic  $\tilde{\gamma}$  is within  $\xi^2 r_1 s_1/50$  of an arc on  $\partial B(\bar{x}_i, 1)$ , an arc that contains a point at distance  $r_j/8$  from  $\bar{x}_i$ . Then, by Lemma 11.39,  $\nu$  meets one of  $\nu(j)^\pm$ , for definiteness let us say  $\nu(j)^+$  and its intersection with  $B_j$  is contained in  $\nu(j)^+ \cup B_{h(x_j)}(w_j, r(w_j)/8 - r_1 s_1/18)$ . Again since  $\nu$  meets the complement of  $\tilde{B}_j$ , its defining geodesic must meet the end of  $\nu(j)^+$  disjoint from  $B_j$ . This shows that, since  $\nu$  meets both  $U(\mathcal{C})$  and its complement, the geodesic  $\tilde{\gamma}$  meets  $U(\mathcal{C})$  and that this intersection is as claimed in the last statement of the claim. If  $\tilde{\gamma}$  is completely contained in  $U(\mathcal{C})$ , then it follows easily that the core of  $\nu$  is contained in  $U(\mathcal{C})$  which contradicts our hypothesis. Hence,  $\tilde{\gamma}$  must also have a point  $p$  not contained in  $U(\mathcal{C})$ . Fix an orientation for  $\tilde{\gamma}$ . Consider the sub-geodesic of  $\tilde{\gamma}$  on the positive side of  $p$ . It may be disjoint from  $U(\mathcal{C})$ . Otherwise, beginning at  $p \in \tilde{\gamma}$  and moving in the positive direction, the first point  $q$  of  $U(\mathcal{C})$  that  $\tilde{\gamma}$  meets is contained in a free end of  $\mathcal{C}$ . If the sub-geodesic on the positive side of  $q$  meets one of the  $B_j$  then its positive endpoint is contained in  $B_j$  and the entire sub-geodesic on the positive

side of  $q$  is contained in  $U(\mathcal{C})$ . Otherwise, it follows from Lemma 11.34 that the intersection of  $\tilde{\gamma}$  with  $U(\mathcal{C})$  is contained in sub-chain of  $\epsilon$ -solid cylinders and is an interval whose other endpoint either is contained in a free end of  $\mathcal{C}$  or is an endpoint of  $\tilde{\gamma}$ . Exactly the same analysis applies to the sub-geodesic of  $\tilde{\gamma}$  on the negative side of  $p$ . Of course, if both sides of  $p$  intersect  $U(\mathcal{C})$ , then the endpoints of  $\tilde{\gamma}$  are contained in  $U(\mathcal{C})$ . This proves the first statement in the claim.

According to Lemma 11.34 the intersection of  $\tilde{\gamma}$  with any  $\epsilon$ -solid cylinder in  $\mathcal{C}$  is contained in the core of that  $\epsilon$ -solid cylinder and as we saw above, by Lemma 11.41 and Corollary 11.39 the intersection of  $\tilde{\gamma}$  with any  $B_i$  is contained in the union of  $B_{h(x_i)}(w_i, (r(w_i)/8) - (r_1 s_1/18))$  and the cores of  $\nu(i)^\pm$ . From all of this, the last statement in the claim follows easily.  $\square$

**Definition 12.41.** When we say that a free end of an  $\epsilon$ -chain is within  $r$  of  $W_2$  implicitly we are measuring distances with the metric used to define the extremal  $\epsilon$ -solid cylinder in the chain having the free end as one of its ends.

**Claim 12.42.** *Suppose that we have a calibrated collection of  $\epsilon$ -chains. Suppose that one of the free ends,  $D^+$ , of one of the chains  $\mathcal{C}$  in the calibrated collection has a point at distance  $\leq r_0$  from  $W_2$ . Let  $\nu$  be the  $\epsilon$ -solid cylinder in  $\mathcal{C}$  that has  $D^+$  as a free end. Then there is an  $\epsilon$ -solid cylinder contained in  $C(W_2)$  that has good intersection with  $\nu$ .*

*Proof.* According to Corollary 12.21 there is a point  $x \in D^+$  that is not contained in  $U_{2,\text{gen}}$  and not contained in any  $\epsilon$ -solid torus. This means that  $x$  is either contained in the core of an  $\epsilon$ -solid cylinder or in a 3-ball near a 2-dimensional corner. Since  $x \in C(W_2)$ , we know that  $B_{h(x)}(x, 1)$  is within  $\hat{\epsilon}$  of a 2-dimensional Alexandrov ball  $B(\bar{x}, 1)$ . Since  $x$  is not contained in  $U_{2,\text{gen}}$  nor in an  $\epsilon$ -solid torus, it follows from Theorem 10.30, Lemma 11.1, and Proposition 11.20 and that  $x$  is within  $\xi^2 r_1 s_1/50$  of a point  $y \in \partial B(\bar{x}, 1)$ . If  $B(\bar{x}, 1)$  is boundary  $\mu$ -flat at  $y$  on scale  $r_1 s_1$  then by Proposition 11.27 there is an  $\epsilon$ -solid cylinder with generating geodesic  $\tilde{\gamma}'$  with  $y$  in the core of  $\nu_\xi(\tilde{\gamma}')$  and with  $f_{\tilde{\gamma}}(y) = -r_1 s_1/16 + (0.0075)r_1 s_1$  when  $\tilde{\gamma}'$  is oriented so that its positive direction exists from  $\nu$  through  $D^+$ . We set  $\nu' = \nu_{c\xi, [-r_1 s_1/16, 0]}$ , where  $c \in [(0.7), (0.9)]$  is chosen so that Condition 2 in Definition 12.28 holds for  $\nu$  and  $\nu'$ . Then  $\nu$  and  $\nu'$  have good intersection and  $\nu' \subset C(W_2)$ .

Lastly, consider the case when  $B(\bar{x}, 1)$  is not  $\mu$ -flat at  $y$  on scale  $r_1 s_1$ . Then by Proposition 10.18  $x$  is contained in a 3-ball  $B_{g'(x')}(w, r(w)/8)$  near a 2-dimensional corner and hence by Lemma 12.24  $x$  is contained in  $B_{h(x_i)}(w_i, 7r(w_i)/8)$ , for some  $i, 1 \leq i \leq N$ . Since we are supposing that  $B(\bar{x}, 1)$  is not boundary  $\mu$ -flat  $y$  on scale  $r_1 s_1$  and that  $x$  is not contained in  $U_{2,\text{gen}}$ , according to Proposition 10.18 this means that  $x$  is contained in  $B_i$ . But this is impossible since  $x$  is contained in a free end of the  $\epsilon$ -chain and since the  $\epsilon$ -chains are calibrated, together they contain all the  $\tilde{B}_i$ .  $\square$

Now suppose that we have a calibrated collection  $\epsilon$ -chains  $\mathcal{C}_1, \dots, \mathcal{C}_k$  with the property that at least one free end of one of these chains, say  $D^+$ , has a point within  $r_0$  of  $W_2$ . Let  $\nu$  be the  $\epsilon$ -solid cylinder that contains  $D^+$ , and let  $\mathcal{C}_i$  be the

$\epsilon$ -chain that  $\nu$  belongs to. Then by the above claim there is an  $\epsilon$ -solid cylinder  $\nu'$  with good intersection with  $\nu$ . If  $\nu'$  meets  $\cup_j U(\mathcal{C}_j)$  only in  $\nu$ , then we extend  $\mathcal{C}_i$  by adding  $\nu'_{c\xi, [-r_1 s_1/16, 0]}$ , (where  $c$  is either (0.7) or (0.9) chosen so that Condition 2 of Definition 12.28 holds) to the end of this calibrated  $\epsilon$ -chain, creating a new calibrated collection of  $\epsilon$ -chains.

Let us suppose now that  $\nu'$  meets  $\cup_j U(\mathcal{C}_j)$  in some point not contained in  $\nu$ . Then by Claim 12.40 we see that, orienting the generating geodesic  $\tilde{\gamma}'$  for  $\nu'$  so that at  $z = \tilde{\gamma}' \cap D^+$  the positive orientation points out of  $\nu$ , and setting  $\alpha$  equal to the open interval in  $\tilde{\gamma}'$  whose closure has endpoints  $z$  and the positive endpoint of  $\tilde{\gamma}'$ , the following hold:

1.  $\alpha$  meets  $U = \cup_j U(\mathcal{C}_j)$ .
2. Let  $p \in \alpha$  be the first point (as we move in the positive direction) meeting the closure of  $U$ . Then  $p$  is contained in the core of a free end,  $D''$ , of one of the  $\mathcal{C}_j$  and  $\nu'$  meets the extremal  $\epsilon$ -solid cylinder, denoted  $\nu''$  and contained in the  $\epsilon$ -chain  $\mathcal{C}_j$ , having  $D''$  as an end.

Denote the generating geodesic of  $\nu''$  by  $\tilde{\gamma}''$ . Let  $D''_1 \subset \nu''$  be the level set of  $f_{\tilde{\gamma}''}$  with the property that the distance from  $D'' \cap \tilde{\gamma}''$  to  $D''_1 \cap \tilde{\gamma}''$  is  $(0.0075)(r_1 s_1/4)$  (in the defining metric for  $\nu''$ ). (Recall that the length of  $\tilde{\gamma}''$  is  $r_1 s_1/4$ .) We divide into two cases.

**Case 1:  $\tilde{\gamma}' \cap D''_1$  is not contained in  $\nu'$ .** In this case we can extend  $\nu'$  so that its positive end contains  $D''_1 \cap \tilde{\gamma}'$ . Since  $\nu'$  meets  $\nu''$  by Lemma 6.1 the metrics defining  $\nu'$  and  $\nu''$  differ by a multiplicative factor between  $(1 + 3(r_1 s_1)^{-1})^2$  and  $(1 - 3r_1 s_1)^{-1}$ . Thus, since the positive end of  $\nu'$  was contained in the level set  $f_{\tilde{\gamma}'}^{-1}(0)$ , after this extension the positive end of  $\nu'$  lies in the level set  $f_{\tilde{\gamma}'}^{-1}(b)$  for some  $0 < b < r_1 s_1/16$ . Thus, the extension produces an allowable  $\epsilon$ -solid cylinder. By construction and by Lemma 6.1 the first condition in Definition 12.28 holds for  $\nu'$  and  $\nu''$ . Since both  $\nu$  and  $\nu''$  are extremal  $\epsilon$ -solid cylinders in the calibrated  $\epsilon$ -chains to which they belong to, each of their width factors is either  $(0.7)\xi$  or  $(0.9)\xi$ . Thus, taking the width factor of the extended version of  $\nu'$  to be  $(0.8)\xi$ , and using Lemma 6.1 we see that  $\nu'_{(0.8)\xi}$  has good intersection with  $\nu''$  and  $\nu'$ . Clearly,  $\nu'$  meets only  $\nu$  and  $\nu''$  and in this case  $\nu'$  has spanned between two calibrated  $\epsilon$ -chains and, with them, forms a single  $\epsilon$ -chain or possibly  $\nu'$  has joined an calibrated  $\epsilon$ -chain to itself creating a circular  $\epsilon$ -chain out of a linear one. Notice that the free ends of the newly formed  $\epsilon$ -chain are also free ends of the original set of  $\epsilon$ -chains. It then follows that the new collection of  $\epsilon$ -chains is calibrated.

**Case 2:  $\tilde{\gamma}' \cap D''_1$  is contained in  $\nu'$ .** In this case arguing as above we can extend  $\nu$  to  $\nu_{c\xi, [-r_1 s_1/16, b]}$  with  $0 < b < r_1 s_1/16$  in such a way that  $\tilde{\gamma}' \cap D''_1$  is contained in its positive end. The extended version of  $\nu_{(0.8)\xi, [-r_1 s_1/16, b]}$  has good intersection with  $\nu''$ . In this fashion, by extending  $\nu$  we have joined two of the calibrated  $\epsilon$ -chains together into one, or possibly we have joined a calibrated  $\epsilon$ -chain to itself to form a circular calibrated  $\epsilon$ -chain out of a linear one.

Thus, in either case, given a calibrated collection of  $\epsilon$ -chains with at least one free end that has a point within distance  $r_0$  of  $W_2$ , we are able to create a new



calibrated collection such that the total space of core is strictly larger. Beginning with  $\coprod_i \tilde{B}_i$  we continue this inductive process until, by compactness of  $W_2$ , we have a calibrated collection of  $\epsilon$ -chains,  $\mathcal{C}_1, \dots, \mathcal{C}_{K_0}$ , with the property that both the free ends of every linear  $\epsilon$ -chain  $\mathcal{C}_i$  have no points within distance  $r_0$  of  $W_2$ .

We set  $U_0 = \cup_{i=1}^{K_0} U(\mathcal{C}_i)$ . There may still be points of  $W_2$  that are not contained in the union of  $U_{2,\text{gen}}$ ,  $\epsilon$ -solid tori, and  $U_0$ . Suppose that  $x \in W_2$  is such a point. Then the ball  $B_{h(x)}(x, 1)$  is within  $\hat{\epsilon}$  of a 2-dimensional Alexandrov ball  $B(\bar{x}, 1)$  of curvature  $\geq -1$  and area  $\geq a$ , and as we have argued before,  $x$  is within  $\xi^2 r_1 s_1 / 50$  of a point  $\bar{y} \in \partial B(\bar{x}, 1)$ . If  $B(\bar{x}, 1)$  is boundary  $\mu$ -flat near  $\bar{y}$  then there is an  $\epsilon$ -solid cylinder  $\nu = \nu_{(0.9)\xi, [-r_1 s_1 / 16, r_1 s_1 / 16]}$  whose core contains  $x$ , and in fact the level set  $f_{\tilde{\gamma}}^{-1}(0)$  contains  $x$ . Since the generating geodesic  $\tilde{\gamma}$  for  $\nu$  is contained in the  $r_0/2$ -neighborhood of  $W_2$ , it does not meet any of the free ends of the  $\mathcal{C}_i$ , and hence by Claim 12.40  $\nu$  is disjoint from  $V_0$ .

Now suppose that  $B(\bar{x}, 1)$  is not boundary  $\mu$ -flat near  $\bar{y}$ . Then  $y \in W_2$  is contained in a ball  $B_{h(w)}(w, r(w)/8)$  near a 2-dimensional corner. As we have seen, this implies that  $y$  contained is one of the  $B_{h(x_i)}(w_i, 7r(w_i)/8)$ . Let  $B(\bar{x}_i, 1)$  be the 2-dimensional of area  $\geq a$  and curvature  $\geq -1$ . Since  $y$  is not contained in  $U_{2,\text{gen}}$ , it follows that  $y$  is close to a point  $\bar{y}' \in \partial B(\bar{x}, 1)$ , and hence either  $B(\bar{x}, 1)$  is boundary  $\mu$ -flat near  $\bar{y}'$  or  $y \in B_i$ . If the first possibility holds then the above shows that  $y$  is contained in the core of an  $\epsilon$ -solid cylinder. The second possibility contradicts the fact that  $y \notin \cup_j U(\mathcal{C}_j)$ . This proves that any point  $y \in W_2$  not contained in  $U_{2,\text{gen}}$ , an  $\epsilon$ -solid torus, or  $U_0$  is in the central disk of the core of an  $\epsilon$ -solid cylinder. Suppose that there is such a point and let  $\nu$  be an  $\epsilon$ -solid cylinder containing the point in the center 2-disk of its core.

**Claim 12.43.** *Let  $D$  be an end of  $\nu$ . Then there is an  $\epsilon$ -solid cylinder  $\nu'_{(0.7)\xi}$  which has good intersection with  $\nu_{(0.9)\xi}$ , which contains the core of  $D$ , and which is also disjoint from  $V_0$ .*

*Proof.* By Claim 12.42 there is an  $\epsilon$ -solid cylinder  $\nu'_{(0.7)\xi}$  with good intersection with  $\nu$  containing the core of  $D$ . Let  $\tilde{\gamma}'$  be the generating geodesic for  $\nu'$ . Then  $\tilde{\gamma}'$  passes within  $2r_1 s_1$  of  $x$  and hence is contained in the  $r_0/2$  neighborhood of  $W_2$  (all distances measured in the defining metric for  $\nu'$ ). As a result  $\tilde{\gamma}'$  does not meet any free end of any of the  $\mathcal{C}_i$ . It follows from Claim 12.40 that  $\nu'$  is disjoint from  $V_0$ .  $\square$

We replace  $\nu'$  by  $\nu'_{(0.7)\xi, [-r_1 s_1 / 16, 0]}$ . Performing the analogous construction for the other end  $D''$  of  $\nu$  produces a calibrated  $\epsilon$ -chain  $\mathcal{C}'$  consisting of three  $\epsilon$ -solid cylinders, with the property that  $\{\mathcal{C}', \mathcal{C}_1, \dots, \mathcal{C}_{K_0}\}$  forms a calibrated collection of  $\epsilon$ -chains. We then repeat the construction above to expand  $\mathcal{C}'$  by adding  $\epsilon$ -solid cylinders to form a calibrated collection of  $\epsilon$ -chains whose free ends are at distance  $\geq r_0$  from  $W_2$ . (Notice that it is possible in the process that we join the  $\epsilon$ -chain  $\mathcal{C}'$  to one of more of the existing calibrated  $\epsilon$ -chains.) By the compactness of  $W_2$ , after a finite number of repetitions of this construction we arrive at a situation where we have a finite collection of  $\epsilon$ -chains  $\mathcal{C}_1, \dots, \mathcal{C}_K$ , which in addition to satisfying Conditions (a), (b), and (c) in Definition 12.39 also satisfy:

- (d) For  $i = 1, \dots, K$ , the free ends of the  $\mathcal{C}_i$  have no points within distance  $r_0$  of  $W_2$ .
- (e)  $W_2$  is contained in the union of  $U_{2,\text{gen}}$ , a finite set of  $\epsilon$ -solid tori, and  $\cup_{i=1}^K U(\mathcal{C}_i)$ .

We say that such a collection is a *complete calibrated* collection. Clearly, a complete calibrated collection of  $\epsilon$ -chains satisfies the conclusion of Theorem 12.38. This completes the proof of the theorem.  $\square$

We now fix the complete calibrated collection  $\{\mathcal{C}_i\}_{i=1}^K$

### 12.5.1 Smaller versions of $\epsilon$ -chains

The next step is to construct smaller versions of the  $\epsilon$ -chains that lie between the  $\epsilon$ -chains and their cores, see FIG. 12.

**Definition 12.44.** Let  $\nu(i) = \nu_{c_i \xi, [a_i, b_i]}(\tilde{\gamma}_i)$ , for  $i = 1, \dots, k$  be a chain of  $\epsilon$ -solid cylinders. Consider a consecutive pair  $\nu(i), \nu(i+1)$  with  $y_i$  being the  $\epsilon$ -control point for  $\nu(i)$  and  $\ell_i$  being the length of  $\tilde{\gamma}_i$ . If Inequality 12.1 holds, i.e., if  $c_i \ell_i p(y_i) < c_{i+1} \ell_{i+1} p(y_{i+1})$ , then we set

$$\nu'(i) = \nu_{(c_i \xi/2), [a_i, b_i]}(\tilde{\gamma}_i)$$

and

$$\nu'(i+1) = \nu_{(c_{i+1} \xi/2), [a_{i+1} + (0.001)\ell_{i+1}, b_{i+1}]}(\tilde{\gamma}_{i+1}).$$

If the opposite inequality holds then we set

$$\nu'(i) = \nu_{(c_i \xi/2), [a_i, b_i - (0.001)\ell_i]}(\tilde{\gamma}_i)$$

and

$$\nu'(i+1) = \nu_{(c_{i+1} \xi/2), [a_{i+1}, b_{i+1}]}(\tilde{\gamma}_{i+1}).$$

Thus, we halve the width of both the  $\epsilon$ -solid cylinders and the truncate the end of the larger one by  $10^{-3}$  times the length of its defining geodesic. We perform an analogous operation for each pair of successive  $\xi$ -boxes, so that it is possible that both ends of  $\nu(i)$  are truncated, only one end is truncated, or neither end is truncated. In all cases the width factor of  $\nu(i)$  is halved so as to become  $c_i \xi/2$ . Notice that in this process we do not truncate any extremal end of the chain.

The result is denoted  $\{\nu'(1), \dots, \nu'(k)\}$ . It is easy to see that the smaller version of a chain of  $\epsilon$ -solid cylinders is also a chain of  $\epsilon$ -solid cylinders. The boundary of  $\bar{\nu}'(1) \cup \dots \cup \bar{\nu}'(k)$  consists of the negative end of  $\bar{\nu}'(1)$  union the positive end of  $\bar{\nu}'(k)$  union an annulus  $E'$  (analogous to the annulus  $E$  from Lemma 12.32), an annulus which is properly embedded in  $\nu(1) \cup \dots \cup \nu(k)$ .

Now let us consider an  $\epsilon$ -chain. It contains a finite number of disjoint chains of  $\epsilon$ -solid cylinders,  $C_1, \dots, C_k$ . We have constructed a smaller version  $C'_i$  of each of the  $C_i$ . Now for each ball  $\tilde{B}_i = \nu(i)^- \cup B_{h(x_i)}(w_i, r(w_i)/8) \cup \nu(i)^+$  we perform the construction analogous to the one above on the  $\nu(i)^\pm$ , possibly shifting the

end not contained in  $B_i$  and cutting its width in half. Also we replace  $B_i$  with  $B'_i = B_{h(x_i)}(w_i, r'(w_i)/8)$  where  $r'(w_i) = r(w_i) - 0.001r_1s_1$ . We set  $\tilde{B}'_i$  equal to the union of  $B'_i$  and the modified versions of the  $\nu^\pm(i)$ . We define the *smaller version* of  $\mathcal{C}$ , denoted  $\mathcal{C}'$ , by taking the union of the  $\mathcal{C}'_i$  and the  $\tilde{B}'_i$ .

**Claim 12.45.** *Let  $\mathcal{C}$  be one of the  $\epsilon$ -chains in the complete calibrated collection.*

1. *The smaller version  $\mathcal{C}'$  of  $\mathcal{C}$  has the property that  $U(\mathcal{C}) \setminus U(\mathcal{C}') \subset U_{2,\text{gen}}$ .*
2. *If  $\mathcal{C}$  is a linear chain then  $\overline{U(\mathcal{C})} \setminus U(\mathcal{C}')$  is homeomorphic to  $S^1 \times I \times I$ , and it meets the union of the two free ends in  $S^1 \times I \times \partial I$ .*
3. *If  $\mathcal{C}$  is a circular chain, the  $\overline{U(\mathcal{C})} \setminus U(\mathcal{C}')$  is homeomorphic to  $T^2 \times I$ .*
4. *Suppose that  $y$  is a point contained in an element of  $\mathcal{C}'$  which is defined using the metric  $h(x)$ . Then  $B_{h(x)}(y, \xi r_1 s_1 / 20) \subset U(\mathcal{C})$ .*

*Proof.* All these results, except the last, are easily established by induction given that the smaller version of a chain of  $\epsilon$ -solid cylinders is itself a chain of  $\epsilon$ -solid cylinders. The last is immediate from the construction and Lemma 6.1 which implies that neighboring elements of  $\mathcal{C}$  are defined using metrics that differ from each other by a multiplicative factor  $R^2$  for some  $(1.1)^{-1} < R < (1.1)$ .  $\square$

Since any point of  $U(\mathcal{C}) \setminus U(\mathcal{C}')$  is contained in  $U_{2,\text{gen}}$ , it follows that we have a finite number of  $\epsilon$ -chains  $\mathcal{C}_i$  with smaller versions  $\mathcal{C}'_i$  of  $\mathcal{C}_i$  with the property that (i) the  $U(\mathcal{C}_i)$  are pairwise disjoint and (ii)  $W_2$  is contained in the union of the  $U(\mathcal{C}'_i)$ ,  $U_{2,\text{gen}}$ , and a finite number of  $\epsilon$ -solid tori.

## 12.6 The Seifert fibration containing $W_2 \setminus \cup_i U(\mathcal{C}'_i)$

We have constructed  $\epsilon$ -chains  $\mathcal{C}_1, \dots, \mathcal{C}_k$  and smaller versions  $\mathcal{C}'_i \subset \mathcal{C}_i$  such that  $W_2$  is contained in the union of  $\cup_i U(\mathcal{C}'_i)$ ,  $U_{2,\text{gen}}$  and a finite number of  $\epsilon$ -solid tori,  $\{B_{h(z_j)}(z_j, r(z_j)/4)\}_{j=1}^N$ , each of which meets  $W_2$ .

**Lemma 12.46.** *We can choose the covering referred to above so that  $\epsilon$ -solid tori are pairwise disjoint and are disjoint from  $\cup_i U(\mathcal{C}_i)$ .*

*Proof.* Suppose two  $\epsilon$ -solid tori  $T_1 = B_{h(z_1)}(z_1, r(z_1)/4)$  and  $T_2 = B_{h(z_2)}(z_2, r(z_2)/4)$  meet. By symmetry we can suppose that the unrescaled radius of  $T_1$  is at least as large as that of  $T_2$ . Then  $T_2$  is contained in  $B_{h(z_1)}(z_1, 3r(z_1)/4)$ , and hence  $T_2$  is contained in the union of  $T_1$  and  $U_{2,\text{gen}}$ . Consequently, we can remove  $T_2$  from the collection keeping it a covering. This allows us to assume that the  $T_i$  are disjoint.

Suppose that  $T_1$  meets an  $\epsilon$ -solid cylinder  $\nu \subset B_{h(x)}(x, 1)$  with generating geodesic  $\tilde{\gamma}$ , with  $\nu$  being one of the elements of one of the  $\epsilon$ -chains  $\mathcal{C}_i$ . Let  $\gamma \subset B(\bar{y}, r_1 s_1 / 3) \subset B(\bar{x}, 1)$  where  $\gamma$  is a  $\mu$  approximation to  $\partial B(\bar{y}, r_1 s_1)$  of length  $r_1 s_1 / 4$  be the generating geodesic for the associated 2-dimensional  $\xi$ -box. Then it follows from Claim 12.23 that  $r(z_1) < 21\xi r_1 s_1$ . Suppose  $p \in T_1 \cap \nu$ . Consider the difference of the  $f_{\tilde{\gamma}}(p)$  and  $f_{\gamma}$  on the end of  $\nu$  closest to  $p$ . If this difference is at least  $50\xi r_1 s_1$ , then the value

of  $f_{\tilde{\gamma}}$  at any point of  $T_1$  is strictly between the values of  $f_{\tilde{\gamma}}$  on the two ends of  $\nu$  and differs by at least  $20\xi r_1 s_1$  from these two values. It then follows that every point  $q$  of  $T_1$  is within  $\hat{\epsilon}$  of a point  $\bar{q}$  of  $B(\bar{y}, r_1 s_1/3)$  that is either contained in  $\nu(\gamma)$  or is distance more than  $\xi r_1 s_1/20$  from  $\partial B(\bar{y}, r_1 s_1)$ . Thus, in this case  $T_1$  is contained in the union of  $\nu$  and  $U_{2,\text{gen}}$ , and hence can be removed from the collection without destroying the covering property.

Suppose the value of  $f_{\tilde{\gamma}}(p)$  is within  $(0.001)r_1 s_1$  of the value of  $f_{\tilde{\gamma}}$  on one of the ends of  $\nu$ . This end is either contained in a neighboring  $\epsilon$ -solid cylinder  $\nu'$  in the  $\epsilon$ -chain, or  $\nu$  is one of the  $\nu(i)^\pm$  and the end in question is contained in  $B_i$ . In the first case let the generating geodesic for  $\nu'$  be denoted by  $\tilde{\gamma}'$ . Then because of the amount of overlap of  $\nu$  and  $\nu'$  and Lemma 6.1,  $f_{\tilde{\gamma}'}(p)$  is strictly between the value of  $f_{\tilde{\gamma}'}$  on the ends of  $\nu'$  and this value differs by at least  $(0.001)r_1 s_1$  from the value of  $f_{\tilde{\gamma}'}$  on either end of  $\nu'$ . Thus, the above argument applies to show that  $T_1 \subset \nu' \cup U_{2,\text{gen}}$ . If  $\nu = \nu(i)^\pm$  and the end in question is contained in  $B_i$ , then because  $r(z_i) \leq 21\xi r_1 s_1$  it follows from Lemma 6.1 that  $T_1$  is contained in  $B_i$ . This proves that if  $T_1$  meets one of the  $\epsilon$ -solid cylinders in  $U(\mathcal{C}_i)$ , then  $T_1$  is contained in the union of  $U(\mathcal{C}_i)$  and  $U_{2,\text{gen}}$  and hence can be removed from the collection.

Now suppose that  $T_1$  meets one of the  $B_i = B_{h(x_i)}(w_i, r(w_i)/8)$ . Since  $r(z_1) < r_1 < r(w_i)$ , it follows that  $T_1$  is contained in  $B_{h(x_i)}(w_i, 7r(w_i)/8)$ . If the intersection of  $T_1$  with  $A = B_{h(x_i)}(w_i, 7r(w_i)/8) \setminus B_i$  contains a point  $q$  within  $\hat{\epsilon}$  of a point  $\bar{q} \in B(\bar{x}, 1)$  which itself is within  $\xi^2 r_1 s_1/100$  of a point  $\bar{q}' \in \partial B(\bar{x}_i, 1)$ , then  $B(\bar{x}_i, 1)$  is boundary  $\mu$ -flat at  $\bar{q}'$  on scale  $r_1 s_1$  and the above argument shows that  $r(z_1) \leq 21\xi r_1 s_1$ . Since  $T_1$  meets  $B_i$ , this implies that  $d_{h(x_1)}(p, w_i) < r(w_i)/8 + (0.001)r_1 s_1$  and hence  $q$  is contained in  $\nu(i)^\pm$ . Any point of  $T_1 \cap A$  that is not within  $\hat{\epsilon}$  of a point in the  $\xi^2 r_1 s_1/100$ -neighborhood  $\partial B(\bar{x}_i, 1)$  belongs to  $U_{2,\text{gen}}$ . This proves that  $T_1$  is contained in  $\tilde{B}_i \cup U_{2,\text{gen}}$  and hence can be removed from the collection.  $\square$

We set

$$W' = (W_2 \cup_i \overline{U(\mathcal{C}_i)} \cup_j \overline{B_{h(z_j)}(z_j, r(z_j)/4)}) \setminus (\cup_i U(\mathcal{C}'_i) \coprod \cup_j B_{h(z_j)}(z_j, r(z_j)/8)).$$

This is a compact set contained in  $U_{2,\text{gen}}$ . Thus, by Proposition 11.8 there is an  $S^1$ -fibration  $V' \rightarrow F'$  whose total space,  $V'$ , contains  $W'$ . The fibers of this fibration are within  $\epsilon'$  of the fibers of any  $\epsilon$  local  $S^1$ -product structure centered at any point of  $W'$ . By Corollary 11.22 there is a Seifert fibration  $V \rightarrow F$ , where

$$V = V' \cup_{i=1}^N B_{h(z_i)}(z_i, r(z_i)/8),$$

which agrees with the restriction of the  $S^1$ -fibration on  $V'$  to a saturated open subset  $V^0 \subset V'$  that contains  $V' \setminus \cup_j B_{h(z_j)}(z_j, r(z_j)/4)$ . Clearly  $W_2 \subset \cup_i U(\mathcal{C}'_i) \cup V$ . The total space  $V^0$  is called the *regular part* of  $V$ . For the rest of the argument, in addition to fixing the complete calibrated chains  $\mathcal{C}_i$  and the  $\epsilon$ -solid tori,  $B_{h(z_j)}(z_j, r(z_j)/4)$ , we fix this Seifert fibration  $V \rightarrow F$ .

## 12.7 Deforming the boundary of $W_2$

Our next step is to deform  $W_2$  by a small isotopy until its boundary is in good position with respect to the Seifert fibration  $V \rightarrow F$  introduced in the previous

subsection. In the case of a torus boundary component, this means deforming that boundary component by an isotopy until it is contained in the regular part,  $V^0$ , of the Seifert fibration and is invariant under the  $S^1$ -fibration structure on  $V^0$ . In the case of a 2-sphere boundary component, this means that deforming the  $S^2$ -sphere by an isotopy until it is the (overlapping) union of spanning disks for two  $\epsilon$ -solid cylinder elements of the  $\{\mathcal{C}\}_i$  and an annulus  $E$  in  $V^0$  that is invariant under the  $S^1$ -fibration. To produce these isotopies, we find appropriate surfaces near to and isotopic to each boundary component  $\Sigma$  of  $W_2$ .

Let  $\Sigma$  be a boundary component of  $W_2$ . Then according to Lemma 12.19 there is a point  $x \in W_2$  and a ball  $B_{h(x)}(x, 1)$  and a 2-dimensional Alexandrov ball  $B(\bar{x}, 1)$  within  $\hat{\epsilon}$  of  $B_{h(x)}(x, 1)$  satisfying the conclusions of that lemma for  $\Sigma$ . In particular, there is in an interval  $(a, b)$  of length  $\geq 1/8000$  and a connected component of  $d_{h(x)}(x, \cdot)^{-1}(a, b)$  that contains all points within  $10^{-5}$  of  $\Sigma$  in the metric  $h(x)$  and on which  $d_{h(x)}(x, \cdot)$  is the projection mapping of a topological product structure. Since  $\Sigma$  is the fiber of an  $\epsilon'$  projection mapping to an interval, it follows that the diameter of  $\Sigma$ , measured in  $g'(x_{\mathcal{E}})$  is at most  $\epsilon' < 10^{-8}$ . Also, the metric  $h(x)$  is either much larger than  $g'(x_{\mathcal{E}})$  (when  $\hat{A}$  is a component close to an interval but which expands to be close to a 2-dimensional ball) or, by Lemma 6.1, is greater than  $(0.9)$  times  $g'(x_{\mathcal{E}})$ . It follows that there is an interval  $(a', b') \subset (a, b)$  with  $b' - a' > 10^{-5}$  such that the pre-image  $Z = d_{h(x)}(x, \cdot)^{-1}(a', b')$  is contained in  $W_2$  and each fiber of the restriction of the distance function  $d_{h(x)}(x, \cdot)$  to  $Z$  is a surface parallel to  $\Sigma$ . We let  $Z' \subset Z$  be the pre-image of an interval  $I$  of length  $(b' - a')/2$  centered in  $(a', b')$ . Similarly, we have  $\bar{Z} \subset d(\bar{x}, \cdot)^{-1}(a', b')$ , and  $d(\bar{x}, \cdot): \bar{Z} \rightarrow (a', b')$  is a topological fibration. We shall see the boundary component  $\Sigma$  of  $W_2$  is either a 2-torus or a 2-sphere depending on whether the level sets of  $d(\bar{x}, \cdot)|_{\bar{Z}}$  are circles or intervals. The easier case, which we deal with first, is when the level sets are circles.

### 12.7.1 Case 1: The fibers of $d(\bar{x}, \cdot)|_{\bar{Z}}$ are circles.

**Proposition 12.47.** *In this case  $\Sigma$  is homeomorphic to a 2-torus and there is a 2-torus  $\Sigma' \subset Z \cap V^0$  that is saturated under the  $S^1$ -fibration structure on  $V^0$  and is parallel in  $Z$  to  $\Sigma$ , see FIG. 13.*

*Proof.* The first thing to see in this case is that every  $\epsilon$ -chain  $\mathcal{C}_i$  is disjoint from  $Z'$ . For suppose that one of the  $\epsilon$ -solid cylinders or one of the  $\tilde{B}_i$  meets  $Z'$ . Then, measured with respect to the metric used to define it, this element has diameter less than  $r_0 \leq 10^{-6}$ . Hence, its diameter with respect to the metric used to define  $Z$  is less than  $(1.2)r_0$ . Thus, this element is contained in  $Z$ . On the other hand, since  $\bar{Z}$  has no boundary,  $Z$  is contained in the union of  $U_{2, \text{gen}}$  and a finite number of  $\epsilon$ -solid tori. Hence, the closure of the  $2r_0$ -neighborhood of  $Z'$  (measured in the metric defining  $Z$ ) is contained in the total space of a Seifert fibration (possibly a different Seifert fibration from the given Seifert fibration on  $V$ ). At the same time, this closure contains either an  $\epsilon$ -solid cylinder or a ball near a 2-dimensional corner. This contradicts Corollary 12.21.

This means that  $Z' \subset V$ . Since  $r(z_i) \leq 10^{-6}$ , any  $\epsilon$ -solid torus  $B_{h(z_i)}(z_i, r(z_i)/4)$  that meets  $Z'$  has closure contained in  $Z$ . Since the diameters of the  $S^1$ -fibers are

at most  $2C\hat{\epsilon} \leq 2 \cdot 10^{-6}$ , it also follows that any  $S^1$ -fiber through a point of  $Z'$  is contained in  $Z$ . Thus, there is a compact sub-Seifert fibration of the given Seifert fibration structure on  $V$  with total space  $X \subset Z$ , whose interior contains  $Z'$  and contains every  $B_{h(z_i)}(z_i, r(z_i)/4)$  that meets  $Z'$  and whose boundary is contained in  $V^0$ . Since each boundary component of the closure of  $Z'$  separates the ends of  $Z$ , there are two boundary components of  $X$ ,  $\partial_{\pm}X$ , on opposite sides of  $Z'$ , each of which separates  $Z'$  from an end of  $Z$ . Each of these boundary components is a 2-torus contained in  $Z \cap V^0$  and is saturated under the  $S^1$ -fibration structure on  $V^0$ . Since a 2-sphere in a Seifert fibration cannot separate two of its boundary components, it follows that the fibers of the restriction of  $d_{h(x)}(x, \cdot)$  to  $Z'$  are not 2-spheres; so neither is  $\Sigma$ . Consequently,  $\Sigma$  is homeomorphic to a 2-torus, and  $Z$  and  $Z'$  are each homeomorphic to the product of  $T^2$  with an interval. Since each of  $\partial_{\pm}X$  separates the ends of  $Z$  and is also homeomorphic to a 2-torus, each is parallel to  $\Sigma$ . We choose  $\Sigma'$  to be  $\partial_+X$ . The surface  $\Sigma'$  is a 2-torus is parallel to  $\Sigma$  in  $Z$ , contained in  $Z \cap V^0$ , and invariant under the  $S^1$ -fibration structure on  $V^0$ . This completes the proof of the proposition and completes the analysis of Case 1.  $\square$

### 12.7.2 Case 2: The fibers of $d(\bar{x}, \cdot)|_{\bar{Z}}$ are intervals.

**Proposition 12.48.** *In this case  $\Sigma$  is homeomorphic to a 2-sphere and is isotopic in  $Z$  to a 2-sphere  $\Sigma'$  that is the overlapping union of spanning disks  $\Delta^{\pm}$  for two  $\epsilon$ -solid cylinder elements of the  $\mathcal{C}_j$  and an annulus  $E$  contained in  $V^0 \cap Z$  and saturated under the  $S^1$ -fibration structure of  $V^0$ .*

The proof of this proposition takes the rest of this subsection.

In this case  $\partial\bar{Z}$  consists of 2 (topological) intervals; say,  $\partial_{\pm}\bar{Z}$ , and consequently  $\partial\bar{Z}'$  is the disjoint union of two intervals  $\partial_+\bar{Z}' \amalg \partial_-\bar{Z}'$  where  $\partial_{\pm}\bar{Z}' = \bar{Z}' \cap \partial_{\pm}\bar{Z}$ . Also, any point of  $\partial B(\bar{x}, 1)$  within  $2r_0$  of  $\partial_{\pm}\bar{Z}'$  is contained in  $\partial_{\pm}\bar{Z}$ .

**Claim 12.49.** *Every point in  $\partial\bar{Z}'$  is within  $\hat{\epsilon}$  of a point of  $\cup_{i=1}^N U(\mathcal{C}'_i)$*

*Proof.* (of the claim) Fix  $\bar{y} \in \partial\bar{Z}'$  and let  $y \in B_{h(x)}(x, 1)$  be a point within distance  $\hat{\epsilon}$  of  $\bar{y}$ . First, let us suppose that  $B(\bar{x}, 1)$  is boundary  $\mu$ -flat at  $\bar{y}$  on scale  $r_1 s_1$ . Then there is an  $\epsilon$ -solid cylinder  $\nu = \nu_{\xi}(\tilde{\gamma})$  with the property that  $y$  is contained in  $D_0$ , the central disk of the core of  $\nu$ . (Notice that we do not claim that  $\nu_{\xi}(\tilde{\gamma})$  is an element of one of the  $\mathcal{C}_i$ .) According to Corollary 12.21,  $D_0$  is not contained in the union of  $U_{2,\text{gen}}$  and  $\epsilon$ -solid tori, and hence there is either an  $\epsilon$ -solid cylinder element  $C$  in one of the chains  $\mathcal{C}_i$  whose core meets  $D_0$  or one of the balls  $B_i = B_{h(x_i)}(w_i, r(w_i))$  has the property that the sub-ball  $B_{h(x_i)}(w_i, r(w_i))/8 - (0.001)r_1 s_1$  meets  $D_0$ . Let  $h(z)$  be the metric used to define either  $C$  or  $B_i$ . If  $\hat{A}$  is a component close to an interval but which expands to be close to a 2-dimensional Alexandrov ball, then  $h(x) = h(z)$ . In this case we set  $R = 1$ . If  $\hat{A}$  is a component close to a 2-dimensional Alexandrov space, then  $y \in B_{h(x)}(x, 2 \times 10^{-5})$ , hence by Lemma 6.1  $h(z) = R^2 h(x)$  for some  $0.99 \leq R \leq 1.01$ . This implies that the diameter of  $D_0$  in the metric used to define  $C$  or  $B_i$  is at most  $(1.01)\xi^2 r_1 s_1$ . It follows that  $D_0 \subset \cup_i U(\mathcal{C}'_i)$ , and hence  $y \in \cup_i U(\mathcal{C}'_i)$ .

Now suppose that  $B(\bar{x}, 1)$  is boundary  $\mu$ -good  $\bar{y}$  on some scale  $r$  with  $r_2 \leq r \leq r_1$  and angle  $\leq \pi - \delta_0$ . Then the ball  $B = B_{h(x)}(y, r(y)/8)$  is a 3-ball near a 2-dimensional corner with  $\bar{y} \in \partial B(\bar{x}, 1)$ . According to Lemma 12.24, one of the 3-balls near 2-dimensional corners that are elements of the  $\mathcal{C}_j$ , say  $B_{h(x_i)}(w_i, r(w_i)/8)$ , has the property that  $B$  is contained in  $B_{h(x_i)}(w_i, 7r(w_i)/8)$ . In particular,  $y \in B_{h(x_i)}(w_i, 7r(w_i)/8)$ . Let  $B(\bar{x}_i, 1)$  be the 2-dimensional Alexandrov ball associated to  $B_{h(x_i)}(x_i, 1)$ . Apply Lemma 11.26 to  $B_{h(x)}(y, r(y))$  and  $B_{h(x_i)}(x_i, 1)$ . Since  $y$  within  $\hat{\epsilon}$  of a point  $\bar{y} \in \partial B(\bar{x}, 1/3)$  and  $\hat{\epsilon} \leq \hat{\epsilon}'_0(\xi^2 r_1 s_1 / 100, a)$ , there is a point  $z$  with  $d_{h(x_i)}(y, z) < \xi^2 r_1 s_1 / 100$  with the property that  $z$  is within  $\hat{\epsilon}$  of a point  $\bar{z} \in \partial B(\bar{x}_i, 7r(w_i)/8)$ . If  $y \in B_{h(x_i)}(x_i, r(x_i)/8 - r_1 s_1)$ , then  $y \in \cup_i U(\mathcal{C}'_j)$ . Otherwise,  $\bar{z} \in (B(\bar{x}_i, 15r(x_i)/16) \setminus B(\bar{x}_i, r(x_i)/16))$  and consequently, by Proposition 10.18  $B(\bar{x}_i, 1)$  is  $\mu$ -flat at  $\bar{z}$  on scale  $r_1 s_1$ . In this case,  $y$  is contained in the intersection,  $D_0$ , of the central disk and core of an  $\epsilon$ -solid cylinder. The argument in the previous paragraph shows that  $D_0 \subset \cup_i U(\mathcal{C}'_i)$ , and hence  $y \in \cup_i U(\mathcal{C}'_i)$ .  $\square$

**Claim 12.50.** *Set  $\bar{Z}'' = d(\bar{x}, \cdot)^{-1}(a' + 3r_0, b' - 3r_0)$  and let  $\mathcal{D}''_{\pm}$  be all the elements of the complete, calibrated  $\epsilon$ -chains  $\{\mathcal{C}_i\}_{i=1}^K$  containing points within  $\hat{\epsilon}$  of  $\partial_{\pm} \bar{Z}''$ . Then  $\mathcal{D}''_+$  and  $\mathcal{D}''_-$  have no elements in common.*

*Proof.* Since the ratio of the metrics used in defining the elements of  $\mathcal{D}''_{\pm}$  are at least 0.99 times  $h(x)$ , it follows that the  $h(x)$ -diameter of any element of  $\mathcal{D}''_{\pm}$  is less than  $(1.02)r_0$ . Thus, every element of  $\mathcal{D}''_{\pm}$  contains no points of  $\partial B(\bar{x}, 1)$  outside  $\bar{Z}$ . Let us show that there is no  $\epsilon$ -solid cylinder common to  $\mathcal{D}''_+$  and  $\mathcal{D}''_-$ . For if there were there would be points of opposite components of  $\partial \bar{Z}''$  within  $3r_1 s_1 / 4$  of each other. But since  $B(\bar{x}, 1)$  has curvature  $\geq -1$  and area  $\geq a$ , it follows from Proposition 10.22 that every point of  $\partial \bar{Z}''$  has a neighborhood of size at least  $r_1 s_1$  and at most  $r_0$  that meets  $\partial \bar{Z}''$  only points in the same component of  $\partial \bar{Z}$ . This contradiction shows that there is no  $\epsilon$ -solid cylinder in common to  $\mathcal{D}''_+$  and  $\mathcal{D}''_-$ . This means the only elements that can be common to  $\mathcal{D}''_+$  and  $\mathcal{D}''_-$  are 3-balls near 2-dimensional corners.

Suppose that there is a  $B_i = B_{h(x_i)}(w_i, r(w_i)/8)$  common to  $\mathcal{D}''_+$  and  $\mathcal{D}''_-$ . Let  $B(\bar{x}_i, 1)$  be the associated 2-dimensional Alexandrov ball, so that  $\bar{w}_i \in \partial B(\bar{x}_i, 1)$  is within  $\hat{\epsilon}$  of  $w_i$  and  $B(\bar{x}_i, 1)$  is boundary  $\mu$ -good at  $\bar{w}_i$  on scale  $r(w_i)$  with  $r_1 \leq r(w_i) \leq r_0$ . Let  $\bar{w}'_i \in \bar{Z}$  be a point within  $\hat{\epsilon}$  of  $w_i \in Z \subset B_{h(x)}(x, 1)$ . Then arguing as above, we see that there is a constant  $R$  with  $(0.99) \leq R_i \leq (1.01)$  such that  $h(x_i) = R_i^2 h(x)$ . Hence,  $R \cdot B(\bar{w}'_i, r(w_i)/8R)$  is within  $4(R+1)\hat{\epsilon}$  of  $B(\bar{w}_i, r(w_i)/8)$ . Hence  $(R/r(w_i))B(\bar{w}'_i, r(w_i)/8R)$  is within  $4r^{-1}(w_i)((R+1)\hat{\epsilon} + \mu)$  of a disk centered at the cone point in a flat cone in  $\mathbb{R}^2$ . Since  $\mu < (1/2)\mu''_0(10^{-6}, a'(a))$ ,  $\hat{\epsilon} \leq (r_1/50)\mu''_0(10^{-6}, a'(a))$ , and  $r_1 \leq r(w_i)$ , according to Proposition 10.18 this implies that for every  $r(w_i)/8R \leq r \leq 7r(w_i)/8R$  the metric ball  $B(\bar{w}'_i, r)$  is a disk and its boundary metric sphere is an arc. In particular, this ball meets only one of  $\partial_{\pm} \bar{Z}''$ . Since  $B_i \subset B_{R^2 h(x)}(w'_i, 7r(w_i)/8\max(R, 1))$ , the ball  $B_i$  contains points within  $\hat{\epsilon}$  of only one of  $\partial_{\pm} \bar{Z}''$ . This is a contradiction and completes the proof.  $\square$

**Claim 12.51.** *There are sub-chains  $\mathcal{C}_+$  and  $\mathcal{C}_-$  of the  $\{\mathcal{C}_i\}_{i=1}^K$ , with no elements in common such that every point of  $\partial_{\pm} \bar{Z}''$  is within  $\hat{\epsilon}$  of  $U(\mathcal{C}_{\pm})$ .*

*Proof.* Let  $\mathcal{C}_\pm$  be all the elements of the  $\epsilon$ -chains  $\{\mathcal{C}_j\}$  that have points within  $\hat{\epsilon}$  of  $\partial_\pm \overline{Z}''$ , respectively. We have seen that these are disjoint collections. It remains to show that each is a chain. A point of  $\partial_\pm \overline{Z}''$  is within  $\hat{\epsilon}$  of points of at most two elements of the  $\{\mathcal{C}_j\}$  and if it is within  $\hat{\epsilon}$  of points in two distinct elements then these elements are neighboring (i.e., intersecting) elements in one of the original chains. Let  $C_1, \dots, C_T$  be the elements of  $\mathcal{C}_+$ . For  $\bar{y} \in \partial_+ \overline{Z}'$ , for any  $j$ , and any  $y \in U(C_j)$  with  $d(\bar{y}, y) < \hat{\epsilon}$  then  $y$  is contained in the smaller version of  $C'_j$  of  $C_j$  so that  $B_{h(x)}(y, \xi r_1 s_1 / 20)$  is either contained in  $C_j$  or is contained in the union of  $C_j$  with one of the neighboring elements  $C_{j'}$  in the chain containing  $C_j$ . Since every point of  $\partial_+ \overline{Z}''$  is within  $\hat{\epsilon}$  of some point in one of the  $\{\mathcal{C}_j\}$ , it follows immediately that the  $C_1, \dots, C_T$  form a sub-chain of one of the  $\mathcal{C}_j$ . By symmetry the argument is the same for  $\partial_- \overline{Z}'$ .  $\square$

Now we are ready to construct the surface in  $Z$  separating its ends and isotopic to  $\Sigma$ . It will be a 2-sphere that is the union of an annulus in  $Z \cap V^0$ , an annulus invariant under the circle fibration, and two spanning disks for  $\epsilon$ -solid cylinders. First, we need a lemma about spanning disks for  $\epsilon$ -solid cylinders, see FIG. 14.

**Lemma 12.52.** *Let  $\nu = \nu_{c\xi, [a, b]}(\tilde{\gamma})$  be an  $\epsilon$ -solid cylinder which is an element of one of the chains  $\mathcal{C}_i$ . Let  $h(y)$  be the metric used to define  $\nu$ . Then there is a spanning disk  $D$  for  $\nu$  that is contained in the interior of a larger disk  $\Delta$  with the following properties:*

1.  $\Delta$  is the (overlapping) union of the spanning disk  $D_0$  for  $\nu_{c\xi, [a, b]}(\tilde{\gamma})$  and an annulus  $E$  that is contained in  $V^0$  and is saturated under the circle fibration structure on  $V^0$ .
2.  $D_0 \cap E$  is an annulus and is a regular neighborhood in  $D_0$  of  $\partial D_0$  and is a regular neighborhood in  $E$  of one of its boundary components. The intersection of  $E$  with the level sets  $h_{\tilde{\gamma}}^{-1}(e\xi)$  are circles for every  $e \in [\xi/10, c\xi]$ .
3. The diameter of  $\Delta$  in the metric  $h(y)$  is less than  $r_1 s_1$ .

*Proof.* Take a point  $w$  on the level set  $f_{\tilde{\gamma}}^{-1}((a+b)/2)$  at distance  $(1.1)c\xi\ell(\tilde{\gamma})$  from  $\tilde{\gamma}$  and take a minimal length curve  $\alpha$  from  $\tilde{\gamma}$  to  $w$ . Set  $\alpha_0$  be the intersection of  $\alpha$  with  $h_{\tilde{\gamma}}^{-1}([\xi\ell(\tilde{\gamma})/20, (1.1)c\xi\ell(\tilde{\gamma})]$ . Then  $\alpha_0 \subset V^0$  and we set  $E'$  equal to the saturation of  $\alpha_0$  under the  $S^1$ -fibration structure on  $V^0$ . It is a smoothly embedded annulus in  $M$  of diameter less than  $2\xi r_1 s_1$ .

**Claim 12.53.** *For every  $e \in [1/10, c]$  the intersection of  $E'$  with the level surface  $h_{\tilde{\gamma}}^{-1}(e\xi\ell(\tilde{\gamma}))$  is a circle separating the boundary components of  $E'$ . Also, every flow line of the vector field  $\chi$  meets  $E'$  in at most one point.*

*Proof.* According to Addendum 11.29 we can choose the coordinates on the local  $S^1$ -product structure centered at any point of  $\alpha_0$ , Euclidean coordinates  $(x, y)$  for the ball in  $\mathbb{R}^2$  and a coordinate  $\theta$  for the circle, such that the intersection of  $\alpha_0$  with this neighborhood is within  $\epsilon'$  in the  $C^1$ -topology of a horizontal line parallel



to the  $y$ -axis oriented so that as we move along  $\alpha_0$  away from  $\tilde{\gamma}$  we are increasing  $y$ . Then using the fact that the  $S^1$ -fibration on the saturation of a neighborhood of  $\alpha_0$  is oriented and that the fibers have an induced metric, there is a natural  $S^1$ -action inducing the fibration. The  $S^1$ -fibration structure in this region is  $C^\infty$  close to the product structure. Thus, we have an  $S^1$ -equivariant map  $\alpha_0 \times S^1 \rightarrow E'$  and for every  $\theta \in S^1$  the arcs  $\alpha_0 \times \{\theta\}$  are also within  $\epsilon'$  in the  $C^1$ -topology of a horizontal line parallel to the  $y$ -axis. Since the directional derivative of  $h_{\tilde{\gamma}}$  is close to 1 on tangent vectors pointing in the positive  $y$ -direction and close to 0 on the  $x$ - and  $\theta$ -axes of this local product structure, it follows that  $h_{\tilde{\gamma}}$  is strictly increasing on  $\alpha_0 \times \{\theta\}$  for every  $\theta \in S^1$  and hence that the intersection of the level sets of  $h_{\tilde{\gamma}}$  with  $E'$  are circles separating the boundary components of  $E'$ .

It follows from this argument that the tangent space to  $E'$  at every point is close to the  $y$ - $\theta$  plane in the local  $S^1$ -product structures. It follows from Addendum 11.29 that in the local  $S^1$ -product structures the vector field  $\chi$  is close to the positive  $x$ -direction. Thus,  $\chi$  is transverse to  $E'$  and any integral curve for this vector field crosses  $E'$  at most once.  $\square$

Now we return to the proof of Lemma 12.52. Let  $D'$  be the level set  $f_{\tilde{\gamma}}^{-1}((a + 3b)/4)$ . It is disjoint from  $E'$ . Since each integral curve of the vector field  $\chi$  meets  $E'$  in at most one point, flowing along the flow lines of this vector field defines a deformation  $H: E'' \times I \hookrightarrow \nu$ , where  $E'' = E' \cap h_{\tilde{\gamma}}^{-1}[5\xi\ell(\tilde{\gamma})/80, 3\xi\ell(\tilde{\gamma})/40]$  such that  $H|_{(E'' \times \{0\})}$  is the inclusion of  $E'' \subset E'$  and  $H(E'' \times \{1\})$  is an annulus  $A' \subset D'$ . Let  $u$  be a weakly monotone  $C^\infty$  function on  $[5\xi\ell(\tilde{\gamma})/80, 3\xi\ell(\tilde{\gamma})/40]$  is identically 1 near  $5\xi\ell(\tilde{\gamma})/80$  and identically 0 near  $3\xi\ell(\tilde{\gamma})/40$ . Then we have the embedding  $H(e', u(h_{\tilde{\gamma}}(e')))$  which is an annulus  $E'''$  connecting the circle  $E' \cap h_{\tilde{\gamma}}^{-1}(3\xi\ell(\tilde{\gamma})/40)$  to the simple closed curve  $C' = H((E' \cap h_{\tilde{\gamma}}^{-1}(5\xi\ell(\tilde{\gamma})/80), 1))$  in  $D'$ . Let  $D'_0 \subset D'$  be the sub-disk bounded by  $C'$ . By Proposition 11.32 the annulus  $E'''$  is contained in  $h_{\tilde{\gamma}}^{-1}([0, \xi\ell(\tilde{\gamma})/10])$ . We define

$$\Delta = E' \cap h_{\tilde{\gamma}}^{-1}([3\xi\ell(\tilde{\gamma})/40, (1.1)c\xi]) \cup E''' \cup \Delta_0.$$

Then  $\Delta$  is an embedded disk. It is the union of the saturation of  $\alpha_0 \cap h_{\tilde{\gamma}}^{-1}(3\xi\ell(\tilde{\gamma})/40, (1.1)c\xi\ell(\tilde{\gamma}))$  under the  $S^1$ -fibration and the spanning disk for  $\nu_{c\xi}(\tilde{\gamma})$ . Clearly, the diameter of  $\Delta$  measured using  $h(y)$  is bounded by  $r_1 s_1$ , and  $\Delta \cap \nu_{c'\xi, [a, b]}(\tilde{\gamma})$  is a spanning disk for all  $c' \in [1/10, c]$ .  $\square$

We fix an  $\epsilon$ -solid cylinder elements  $\nu^\pm = \nu_{c_\pm \xi}(\tilde{\gamma}^\pm)$  of the chains  $\mathcal{C}'_\pm$ ,  $\epsilon$ -solid cylinders that have points within  $\hat{\epsilon}$  of  $\partial_\pm \bar{Z}$ . For each, invoking Lemma 12.52 we construct a disk  $\Delta^\pm$  which is the union of a spanning disk of  $\nu_{c_\pm \xi/2}(\tilde{\gamma}^\pm)$  and an annulus  $E^\pm$  saturated under the  $S^1$ -fibration structure on  $V^0$ . Since  $\Delta^\pm$  contains points that are close to points of  $\partial_\pm \bar{Z}$  and has diameter less than  $r_1 s_1$  it follows that  $\Delta^\pm \subset Z''$ . Consider the open subset  $U^0$  of  $V^0$  consisting of the union of all fibers of the  $S^1$ -fibration structure that meet  $Z'' \cap (V^0 \setminus (U(\mathcal{C}'_+) \cup U(\mathcal{C}'_-)))$ . Let  $B$  denote the base of this fibration and let  $\alpha^\pm \subset B$  be image under the quotient mapping of  $E^\pm \cap U^0$ . Each  $\alpha^\pm$  is a union of arcs. Let  $a^\pm$  be the endpoint of  $\alpha^\pm$  which is the image of the boundary circle of  $\Delta^\pm$ .

**Claim 12.54.** *The points  $a^\pm$  are contained in the same connected component of  $B$ .*

*Proof.* Since  $\overline{Z}''$  is connected, there is an arc in  $Z''$  connecting  $\partial\Delta^+$  and  $\partial\Delta^-$ . We deform this arc slightly, relative to its endpoints, so that it is disjoint from  $\cup_j U(\mathcal{C}'_j)$  and all the  $\epsilon$ -solid tori  $B_{h(z_i)}(z_i, r(z_i)/8)$  in the covering we have fixed. (This is possible since the inclusion of the frontier of each of these elements into the element induces a bijection on connected components.) This deformation can be chosen so that it does not move the arc more than distance  $r_0$  in the metrics defining the elements of the  $\mathcal{C}_j$  and the  $\epsilon$ -solid tori, and hence does not move the arc outside  $Z''$ . The result is an arc in  $U^0$ . Projecting this arc to  $B$  establishes the claim.  $\square$

The image of each of  $\Delta^\pm \cap U^0$  in  $B$  is a disjoint union of embedded arcs, denoted  $\delta^\pm$ , and  $\delta^+ \cap \delta^- = \emptyset$ . Since the points  $a^\pm$  (which are endpoints of one of the components of  $\delta^\pm$ ) are in the same component of  $Z$ , it follows that there is an embedded arc  $\alpha$  in  $B$  from  $a^+$  to  $a^-$  that is otherwise disjoint from  $\delta^\pm$ . We can choose  $\alpha$  so that  $\delta^+ \cup \alpha \cup \delta^-$  is a disjoint union of smoothly embedded arcs in  $B$ . The pre-image of  $\alpha$  is an annulus  $E_1$  with the property that  $\Delta^+ \cup E_1 \cup \Delta^-$  is a smoothly embedded 2-sphere  $\Sigma' \subset Z''$ . Since  $\Sigma'$  meets both  $\mathcal{C}'_\pm$  in a spanning disk, it follows that  $\Sigma'$  separates the ends of  $Z''$ . Since  $Z''$  is fibered over an interval, it then follows that  $Z''$  is homeomorphic to  $S^2 \times (a', b')$ , that  $\Sigma$  is a 2-sphere, and that the region in  $Z''$  between  $\Sigma'$  and  $\Sigma$  is a product region. By construction  $\Sigma'$  is the (overlapping) union of two spanning disks  $\Delta^\pm \cap \nu^\pm$  for  $\epsilon$ -solid cylinder elements and an annulus  $E = E^- \cup E_1 \cup E^+$  contained in  $V^0$  and saturated under the  $S^1$ -fibration. For each  $\Delta^\pm$  the intersection  $E \cap \Delta^\pm$  is a regular neighborhood in  $\Delta^\pm$  of  $\partial\Delta^\pm$  and is a regular neighborhood in  $E$  of the corresponding boundary component of  $E$ , see FIG. 15.

This completes the construction for each component  $\Sigma$  of  $W_1 \cap W_2$  of a surface  $\Sigma' \subset W_2$  isotopic to  $\Sigma$  in  $W_2$  and satisfying the conditions stated in Proposition 12.48.

### 12.7.3 Modification of $W_1$ and $W_2$ by isotopy

We modify the decomposition  $M = W_1 \cap W_2$  by isotopy supported near  $W_1 \cap W_2 = \partial W_2$ , an isotopy that moves each boundary component  $\Sigma$  of  $W_2$  onto the corresponding  $\Sigma'$  just constructed. Of course, this deformation does not change the topological type of any component of  $W_1$  or  $W_2$ .

### 12.8 Removing the solid tori and solid cylinders from $W_2$

The last step in the argument is to remove a disjoint union of solid tori and solid cylinders from  $W_2$  with the following three properties: (i) the union of these solid tori and solid cylinders contains the intersection  $W_2 \cap (\cup_i U(\mathcal{C}'_i))$ ; (ii) the boundaries of the solid tori and the sides of the solid cylinders are contained in  $V^0$  and are saturated under the  $S^1$ -fibration structure; and (iii) the ends of the solid cylinders are contained in the 2-sphere boundary components of  $W_2$  and the complement of the ends of these solid cylinders in each 2-sphere is contained in  $V^0$  and is fibered under the  $S^1$ -fibration structure on  $V^0$ .

### 12.8.1 The $S^1$ -invariant torus boundary components associated with circular $\epsilon$ -chains

Suppose that  $\mathcal{C}_i \subset W_2$  is a circular  $\epsilon$ -chain. Then according to Claim 12.45 the submanifold  $G = \overline{U(\mathcal{C}_i)} \setminus U(\mathcal{C}'_i)$  is homeomorphic to  $T^2 \times I$ . We have a Seifert fibration  $V \rightarrow F$  with  $G$  contained in the regular part  $V^0$  of this Seifert fibration. Furthermore, since the fibers of the  $S^1$ -fibration on  $V^0$  are within  $\epsilon'$  of vertical in the local  $S^1$ -product structures with  $\epsilon$ -control center at each point of  $G$ , and since these latter fibers have length at most  $C\hat{\epsilon}$ , it follows that no fiber of the  $S^1$ -fibration structure on  $V^0$  meets both boundary components of  $G$ . Consequently, we have an open saturated subset of  $V$  that contains the complement in  $W_2$  of  $\cup_j U(\mathcal{C}_j)$  and which is disjoint from  $\cup_j U(\mathcal{C}'_j)$ . Hence, there is a compact sub-Seifert fibration  $X \rightarrow \bar{F}'$  of  $V \rightarrow F$  that contains  $W_2 \setminus \cup_j U(\mathcal{C}_j)$  and is disjoint from  $\cup_j U(\mathcal{C}'_j)$ . One of the boundary components,  $T_i$ , of  $X$  separates the boundary of  $U(\mathcal{C}'_i)$  from the complement of  $U(\mathcal{C}_i)$ . Since it is fibered by circles and since it is a boundary component of an orientable 3-manifold,  $T_i$  is homeomorphic to a 2-torus. Since it separates the boundary components of  $G$  and since  $G$  is diffeomorphic to  $T^2 \times I$ ,  $T_i$  is parallel in  $G$  to  $\partial U(\mathcal{C}'_i)$ . As such  $T_i$  bounds a solid torus  $\tau_i$  in  $U(\mathcal{C}_i)$  containing  $U(\mathcal{C}'_i)$ .

We perform a similar construction for each circular  $\epsilon$ -chain  $\mathcal{C}_i$  producing an  $S^1$ -invariant torus  $T_i$  in  $U(\mathcal{C}_i)$ , a torus that bounds a solid torus  $\tau_i$  in the circular  $\epsilon$ -chain  $U(\mathcal{C}_i)$ . Then we truncate the Seifert fibration along these tori. This produces a partially compactified sub-Seifert fibration whose boundary components are tori, the  $T_i$ , one for each circular  $\epsilon$ -chain. Each torus boundary component  $T_i$  of the sub-Seifert fibration bounds a complementary component  $\tau_i$  of  $W_2$  that is a solid torus in  $U(\mathcal{C}_i)$ .

### 12.8.2 Construction for linear $\epsilon$ -chains

Now let  $\mathcal{C}_i$  be a linear  $\epsilon$ -chain. Then we have the smaller version  $\mathcal{C}'_i$  and the difference  $\overline{U(\mathcal{C}_i)} \setminus U(\mathcal{C}'_i)$  is homeomorphic to  $S^1 \times I \times I$ . Furthermore, since the 2-sphere boundary components of  $W_2$  meet both the  $\epsilon$ -chains and their smaller versions in spanning disks, the intersection of this difference with  $W_2$  is also homeomorphic to  $S^1 \times I \times I$  and is contained in  $V^0$ . We consider the open subset of  $V^0$  consisting of all  $S^1$ -orbits that are closer to the complement of  $U(\mathcal{C}_i)$  than they are to  $U(\mathcal{C}'_i)$ . This open  $S^1$ -saturated subset of  $V^0$  contains the side of  $U(\mathcal{C}_i)$  and is disjoint from  $U(\mathcal{C}'_i)$ . Hence, there is a compact  $S^1$ -saturated subset  $X$  of  $V^0$  that contains the side of  $U(\mathcal{C}_i)$  and is disjoint from  $U(\mathcal{C}'_i)$ . One of the boundary components  $\partial_i X$  of  $X$  separates the side of  $U(\mathcal{C}_i)$  from  $U(\mathcal{C}'_i)$ . As such,  $\partial_i X$  contains an annulus  $E_i$  contained in  $\overline{U(\mathcal{C}_i)}$  with one boundary circle in each end of  $U(\mathcal{C}_i)$ , an annulus that is saturated under the  $S^1$ -fibration. The annulus  $\hat{E}_i$  is the frontier in  $\overline{U(\mathcal{C}_i)}$  of a solid cylinder in  $\overline{U(\mathcal{C}_i)}$ . The intersection of  $E_i = \hat{E}_i \cap W_2$  is also a saturated annulus and bounds a solid cylinder  $K_i$  in  $W_2$  containing  $U(\mathcal{C}'_i) \cap W_2$  with each end of  $K_i$  being disk in one of the 2-sphere components  $\Sigma'$  of  $\partial W_2$ . This disk contains a component of the intersection of  $U(\mathcal{C}_i)$  with  $\Sigma'$  and is contained in a component of

the intersection of  $U(\mathcal{C}_i)$  with  $\Sigma'$ .

## 12.9 Completion of the Proof

We set  $Y \subset W_2$  equal to the union of the solid tori  $\tau_i$ , one for each circular  $\epsilon$ -chain  $\mathcal{C}_i$ , and the solid cylinders  $K_j$ , one for each linear  $\epsilon$ -chain  $\mathcal{C}_j$ . Recall that  $T_i = \partial\tau_i$  and the sides  $E_j$  of the  $K_j$  are contained in  $V^0$  and are saturated under the  $S^1$ -fibration, and ends of the  $K_j$  are 2-disks in 2-sphere boundary components of  $W_2$ . We define  $V_1 = W_1$  and we define  $V_2$  to be the complement in  $W_2$  of the relative interior of  $Y$ .

Now let us revert to the original notation of the Riemannian manifolds  $(M_n, g_n)$  and recap our progress to date. We change notation so that  $W_1, V_1, W_2$  and  $V_2$  become  $W_{n,1}, V_{n,1}, W_{n,2}$  and  $V_{n,2}$ , respectively. By Proposition 12.18 we see that the topological type of each component of  $V_{n,1}$  is one of the types list in Part 1 of Theorem 6.2. Also, by construction  $V_{n,2}$  is the total space of a compact Seifert fibration and  $V_{n,2} \cap V_{n,1}$  consists of an  $S^1$ -saturated family of torus boundary components and  $S^1$ -saturated annuli contained in 2-sphere boundary components of  $V_{n,1}$ . Furthermore, each  $S^2$ -boundary component of  $V_{n,1}$  meets  $V_{n,2}$  in exactly one annulus, and each  $T^2$ -boundary component of  $V_{n,1}$  is a boundary component of  $V_{n,2}$  if and only if it is not a boundary component of  $M_n$ . Lastly,  $M_n \setminus (\text{int}(V_{n,1} \cup V_{n,2}))$  is a disjoint union of solid tori and solid cylinders. The boundary components of the solid tori are boundary components of  $V_{n,2}$  and the solid cylinders meet the boundary of  $V_{n,2}$  in annuli saturated under the  $S^1$ -fibration structure and meet the boundary of  $V_{n,1}$  in their end disks. This completes the proof that  $V_{n,1}$  and  $V_{n,2}$  satisfy of all the conditions required by Theorem 6.2.

This completes the proof of Theorem 6.2 and hence of the Geometrization Conjecture (Corollary 5.6).

Notice that we have one extra condition not stated (nor required) in Theorem 6.2. Namely, for each solid torus component  $\tau_i$  of  $M_n \setminus (V_{n,1} \cup V_{n,2})$ , the fibers of the  $S^1$ -fibration on its boundary,  $T_i$ , bound disks in  $\tau_i$ . This means that the Seifert fibration structure on  $V_{n,2}$  does not extend over any of the  $\tau_i$ .

## 13 The Equivariant Case

In this last section we establish that the decomposition and the locally homogeneous metrics in the Geometrization Conjecture can be taken to be equivariant with respect to any compact group action. Arguments here are very similar to ones in [6] where the case of manifolds with either hyperbolic or elliptic geometry are considered and rely on those in [17] where the case of actions on locally homogeneous manifolds modelled Solv, Nil, Flat,  $\mathbb{H}^2 \times \mathbb{R}$  or  $\widetilde{PSL}_2(\mathbb{R})$  was considered. Also, the result is closely related to those of [16] where similar, but slightly, stronger results are established in the context of 3-dimensional orbifolds. Let us begin with the notion of a linear action on a family of balls and the definition of equivariant connected sum.

**Definition 13.1.** A linear action of a compact group  $K$  on a compact  $n$ -ball  $B$  is

the action induced from the defining action on a closed ball  $B$  in  $\mathbb{R}^n$  centered at the origin by an embedding  $K \subset O(n)$ . The induced action of  $K$  on the boundary of  $B$  is said to be a *linear action of  $K$  on a sphere of dimension  $(n - 1)$* . A *linear action of a compact group  $H$  on a family of compact  $n$ -balls* is an action for which there is a subgroup  $K \subset H$  of finite index such that the action is the natural left action of  $H$  on  $H \times_K B^n$  where the action of  $K$  on  $B$  is linear. Notice that  $H \times_K B$  is diffeomorphic to a disjoint union of  $n$ -balls, the number of balls being the cardinality of  $H/K$ . We also say that the restriction of this action to  $H \times_K \partial B$  is a *linear action of  $H$  on a family of  $(n - 1)$ -spheres*.

A *linear action on  $T^2$*  is an action preserving a flat metric.

First, we have an elementary result from dimension 2, see [30]:

**Proposition 13.2.** *Any compact group action on a closed surface  $\Sigma$  leaves invariant a metric of constant curvature on  $\Sigma$ . In particular, any compact group action on  $S^2$  or  $T^2$  is equivariantly diffeomorphic to a linear action.*

*Proof.* The case of finite groups follows from the classification of 2-dimensional orbifolds. The case of actions of positive dimensional groups is elementary.  $\square$

**Definition 13.3.** Suppose that  $M$  is an oriented manifold  $M$  with a smooth action  $H \times M \rightarrow M$  of a compact group  $H$ . Suppose that there is a homomorphism from  $\mathcal{O}: H \rightarrow \mathbb{Z}/2\mathbb{Z}$  which is the orientation character of this action in the sense that for any  $h \in H$  and any  $x \in M$  the map  $dh_x: T_x M \rightarrow T_{hx} M$  is orientation-preserving if and only if  $\mathcal{O}(h) = +1$ . Suppose that there is a disjoint family of compact smooth balls  $\mathcal{B} \subset M$  invariant under  $H$  and an  $H$ -equivariant diffeomorphism from  $\mathcal{B}$  to a linear action of  $H$  on a disjoint union of balls. Lastly, suppose that there is an orientation-reversing  $H$ -equivariant involution  $\psi: \mathcal{B} \rightarrow \mathcal{B}$  that induces a free involution on the components of  $\mathcal{B}$ . Given  $M, \mathcal{B}$  and  $\psi$ , we form the  $H$ -connected sum as follows. Let  $c: \partial \mathcal{B} \times [0, 1] \rightarrow \mathcal{B}$  with  $c|_{\partial \mathcal{B}} = \text{Id}_{\partial \mathcal{B}}$ , be an  $H$ -equivariant collar that is identified by the isomorphism to the linear model with the radial collars. Any two such neighborhoods (coming from different isomorphisms to the linear action) are equivariantly isotopic in  $\mathcal{B}$  by an isotopy that is the identity on the boundary. Given an  $H$ -equivariant collar, set  $C = c(\partial \mathcal{B} \times (0, 1))$ . Let  $\mathcal{B}' \subset \mathcal{B}$  be the complement of  $c(\partial \mathcal{B} \times [0, 1])$ . Then the connected sum is defined as the quotient of  $M \setminus \mathcal{B}'$  by the involution on  $C$  defined by  $(x, t) \mapsto (\psi|_{\partial \mathcal{B}}(x), 1 - t)$ . The result is an oriented manifold, with the orientation compatible with that on the  $M \setminus \mathcal{B}'$ , that carries a natural  $H$ -action compatible with the action on the  $M \setminus \mathcal{B}'$ . This action has the same orientation character as the original actions. Since the collars are unique up to equivariant isotopy the resulting connected sum is unique up to equivariant diffeomorphism. It is the  *$H$ -connected sum defined by  $\mathcal{B}$  and  $\psi$* .

Here is the main equivariant result, which we establish in this section.

**Theorem 13.4.** *Suppose that  $M$  is a compact, orientable smooth 3-manifold and that  $H \times M \rightarrow M$  is a smooth action of a compact group  $H$  on  $M$  with an orientation character  $\mathcal{O}_M$ . Then there is a disjoint union of oriented, prime 3-manifolds*

$P = P_1 \amalg \cdots \amalg P_k$ , an  $H$ -action  $H \times P \rightarrow P$  with an orientation character  $\mathcal{O}_P$ , an  $H$ -invariant family of balls  $\mathcal{B}$  on which the  $H$ -action is equivariantly diffeomorphic to a linear action, and an orientation-reversing involution  $\psi: \mathcal{B} \rightarrow \mathcal{B}$  acting freely on the connected components of  $\mathcal{B}$  such that  $H \times M \rightarrow M$  is  $H$ -equivariantly diffeomorphic to the  $H$ -connected sum of  $P$  determined by  $\mathcal{B}$  and  $\psi$ , and with  $\mathcal{O}_M$  induced by  $\mathcal{O}_P$ . Furthermore, in  $P$  there is an  $H$ -invariant disjoint union of incompressible tori and Klein bottles,  $\widehat{T}(P) \subset P$ , such that  $P \setminus \widehat{T}(P)$  admits an  $H$ -invariant Riemannian metric with the property that the restriction of this metric to each connected component is a complete, locally homogeneous metric of finite volume.

### 13.1 Preliminary results on compact group actions in dimension 3.

We begin with sub-actions of standard actions.

**Proposition 13.5.** *Suppose that  $H \times (S^2 \times I) \rightarrow S^2 \times I$  is the product of the linear action of a compact group  $H$  on  $S^2$  with the trivial action on  $I$ . Suppose that  $\Sigma \subset S^2 \times \text{int } I$  is an  $H$ -invariant smoothly embedded 2-sphere separating the ends of  $S^2 \times I$ . Let  $X$  be the compact, connected submanifold of  $S^2 \times I$  with boundary  $(S^2 \times \{0\}) \amalg \Sigma$ . Then the identity map from  $S^2 \times \{0\}$  to itself extends to an  $H$ -equivariant diffeomorphism from  $S^2 \times I$  to  $X$ .*

*Proof.* Let  $Z$  be a non-trivial cyclic subgroup (finite or infinite) of  $H$  acting in an orientation-preserving fashion on  $S^2 \times I$ . Then the fixed set of  $Z$  is the product of 2 points in  $S^2$  with the interval. Since  $\Sigma$  separates the ends of  $S^2 \times I$ , each of these intervals meets  $\Sigma$ . Also, the action of  $Z$  on  $\Sigma$  is orientation-preserving and hence has exactly two fixed points. Thus,  $\Sigma$  meets each arc of fixed points for  $Z$  in a single point. Its tangent plane is  $Z$ -invariant and thus transverse to the arc of fixed points. Similarly, if  $Z$  is generated by reflection in a codimension-1 subspace of  $S^2$ , then the fixed set  $F_Z$  of  $Z$  is the product of a circle in  $S^2$  with  $I$  and hence meets  $\Sigma$ . The action of the reflection on  $\Sigma$  is orientation-reversing and hence is reflection in a circle in  $\Sigma$ . Once again, an examination of the tangent planes shows that  $\Sigma$  is transverse to  $F_Z$ . The circle  $\Sigma \cap F_Z$  separates the ends of the annulus  $F_Z$ .

Now let us turn to the proof of the result.

**Case when  $H$  has dimension 3.** In this case the group contains  $SO(3)$  and every orbit is a two-sphere of the form  $S^2 \times \{t\}$  for some  $t \in I$ . Being an  $H$ -invariant,  $\Sigma$  is one of these 2-spheres and the restriction of the product structure is  $H$ -invariant.

If  $H$  is not of dimension 3, then it is of dimension  $\leq 1$ .

**Sub-case when  $H$  has dimension 1.** Then  $H$  contains a circle subgroup and the quotient of  $H$  by this normal subgroup has order 1, 2, or 4. The circle subgroup of  $H$  acts with two fixed points on  $S^2$ , and hence the fixed points of the circle action on  $S^2 \times I$  are two vertical arcs. The 2-sphere  $\Sigma$  is transverse to each arc and meets each in a single point. We can deform  $\Sigma$  by an equivariant isotopy until  $\Sigma$  meets a regular neighborhood of each arc in a disk at level  $1/2$ . The complement of these regular neighborhoods  $V$  is an annulus times  $I$  and  $\Sigma$  meets it in an annulus separating the ends. The quotient  $V/S^1$  is a rectangle and the image of  $\Sigma \cap V$  is an interval connecting the sides of the rectangle. The induced action of  $H/S^1$  on

this rectangle is either trivial or is a reflection about the mid-line. Of course  $\Sigma/S^1$  is invariant under this action. Hence, there is an isotopy of  $\Sigma/S^1$ , equivariant with respect to the induced action of  $H$  on the rectangle and relative to a neighborhood of its endpoints to the interval at height  $1/2$ . This isotopy lifts to an  $S^1$ -equivariant isotopy of  $\Sigma$ , relative to a neighborhood of its intersection with the fixed point arcs to the 2-sphere at level  $t$ . This completes the proof when  $H$  is 1-dimensional.

**Sub-case when  $H$  has dimension 0.** In this case  $H$  is a finite group and there are finitely many components of fixed points of non-trivial elements of  $H$ . The local models are: (i) arcs of fixed points of cyclic group actions: (ii) annular regions of reflection fixed points, and (iii) ‘pin wheels’ of  $n$  annular regions fixed under reflections meeting along an arc with the centralizer of the central arc being a dihedral group.  $\Sigma$  crosses each of these components transversely, either in a single point if the component is an arc, in a single circle separating the ends if the component is an annulus, and in a pin wheel of circles in the pin wheel of annuli in the dihedral case. There is an equivariant tubular neighborhood  $\nu$  of the union of these fixed point components given by a product of a tubular neighborhood of the fixed set in  $S^2 \times \{0\}$  with the interval. Choosing the neighborhood small enough, we can assume that the intersection  $\Sigma \cap \nu$  divides  $\nu$  into two components each of which is diffeomorphic to a product of  $\Sigma \cap \nu$  with an interval. Deforming by an  $H$ -equivariant isotopy, we can make this intersection have a constant  $I$ -coordinate  $1/2$ . Now we consider the complement of an open tubular neighborhood. The 3-manifold in question is a product of a compact subsurface  $Y \subset S^2 \times \{0\}$  with  $I$ , and  $(Y \times I) \cap \Sigma$  is a surface diffeomorphic to  $Y$  embedded into  $Y \times I$  in such a way that its boundary is  $\partial Y \times \{1/2\}$ .

Let  $Y_0$  be a connected component of  $Y$ . The intersection of  $\Sigma \cap (Y_0 \times I)$  is diffeomorphic to  $Y_0$  and its boundary is embedded at a constant level  $1/2$ . We claim that the inclusion  $\Sigma \cap (Y_0 \times I) \rightarrow Y_0$  induces an isomorphism on fundamental groups. First, let us show that it is injective. If not, then by Dehn’s lemma and the loop theorem there is a disk embedded in  $Y_0 \times I$  whose boundary is a non-trivial embedded loop in  $\Sigma$ . Since  $\Sigma \cap (Y_0 \times I)$  is a planar surface, every non-trivial embedded loop separates the boundary components into two non-empty sets. But then no such loop can bound in  $Y_0 \times I$ . Once we know that the fundamental group of  $\Sigma \cap (Y_0 \times I)$  injects into  $\pi_1(Y_0 \times I)$  we have the Seifert-Van Kampen theorem giving the fundamental group of  $Y_0 \times I$  as a free product with amalgamation over in the maps on fundamental groups induced by the inclusions of  $\Sigma \cap (Y_0 \times I)$  into the two sides. If one of these inclusions does not induce a surjective homomorphism then the fundamental group of the other components does not surject onto  $\pi_1(Y_0 \times I)$ , but this is absurd since each component contains a copy of  $Y_0$ .

Once we know that  $\pi_1(\Sigma \cap (Y_0 \times I))$  maps isomorphically onto  $\pi_1(Y_0 \times I)$ , the same is true of the quotients by the stabilizer of  $Y_0$  in  $H$ . To complete the proof in this case we invoke an elementary consequence of Dehn’s lemma and the loop theorem.

**Lemma 13.6.** *Suppose that  $Y$  is a compact, connected smooth 3-manifold with corners. We suppose that  $Y$  is irreducible in the sense every embedded 2-sphere in  $Y$*

bounds a 3-ball in  $Y$ . We suppose that  $\partial Y$  is the union of three compact sub-surfaces with disjoint interior:  $\partial Y = X_+ \cup (\partial X_+ \times I) \cup X_-$  with  $X_\pm$  being connected and with  $\partial X_+ \neq \emptyset$ . We suppose that  $Y$  has corners exactly along  $\partial X_+ \amalg \partial X_-$ . Suppose that  $X_+$  is the quotient of a planar surface with non-empty boundary by a free action of a finite group. If the inclusion  $X_+ \subset H$  induces an isomorphism on fundamental groups then  $Y$  is diffeomorphic to  $(X_+ \times I, X_+ \times \{0\})$  by a diffeomorphism extending the given product structure on  $\partial X_+ \times I$ .

Applying this lemma to the submanifold  $\overline{X}_0$  in  $[Y_0/\text{Stab}(Y_0)] \times I$  with boundary  $Y_0/\text{Stab}(Y_0) \times \{0\}$  and  $\Sigma \cap (Y_0 \times I)/\text{Stab}(Y_0)$  we conclude that, provided that  $Y$  has non-empty boundary,  $\overline{X}_0$  has a product structure extending the given product structure on the boundary. We can choose these product structures on the various components so that they combine to produce an  $H$ -invariant product structure on  $[(Y_0 \times I) \setminus \nu \cap X]$  extending the given product structure on the boundary. Putting this together with the product structure on  $\nu \cap X$  gives the result in all cases where  $Y \neq S^2$ .

The final case we need to study is when  $Y = S^2$ . In this case  $H$  acts freely on  $S^2$ . This means that  $H$  is a cyclic group of order 1 or 2. If  $H$  is trivial, the result is immediate since any  $S^2 \subset S^2 \times I$  that separates the ends is isotopic to a level  $S^2$ . In the case when  $H$  is of order two, the quotient of  $S^2$  and  $\Sigma$  by  $H$  is  $\mathbb{R}P^2$ . Thus we need to show that  $\mathbb{R}P^2 = \Sigma/H$  embedded in  $\mathbb{R}P^2 \times I$  separates into two product regions. We take an embedded annulus connecting non-trivial embedded loops in the ends of  $\mathbb{R}P^2 \times I$ . We can take this annulus transverse to  $\Sigma/H$ . The intersection is a finite collection of circles, some trivial and one non-trivial. By a standard inner most disk argument, we remove all the trivial circles. This makes the intersection a circle separating the ends of the annulus and generating the fundamental group of  $\mathbb{R}P^2$ . We can deform  $\Sigma/H$  until this intersection circle is a level  $t$ . Cutting out a neighborhood of this annulus gives us a 3-ball  $B_0$  meeting each end,  $S^2 \times \partial_\pm I$  in a 2-disk,  $D_\pm$ . The intersection of  $\Sigma/H$  with  $\partial B_0$  is a disk  $D$  separating  $D_+$  and  $D_-$  and we have a product structure on  $\partial B_0 \setminus \text{int}(D_+ \cup D_-)$  and the boundary of  $D$  is a level circle in this product structure. It follows easily that there is a product structure  $B_0 = D_+ \times I$  extending the product structure already given on the boundary and with  $D$  being the disk at level  $t$ . This completes the proof in the case when the dimension of  $H$  is zero and hence completes the proof of the proposition.  $\square$

One consequence of this is that the restrictions of linear actions on balls to invariant sub-balls are also linear actions.

**Proposition 13.7.** *Let  $H$  be a compact group and let  $\mathcal{B} = H \times_K B$  be a family of balls with the natural left  $H$  action being linear. Suppose further that  $\mathcal{B}'$  is a family of compact 3-dimensional balls with  $\mathcal{B}' \subset \text{int } \mathcal{B}$  with the property that the inclusion  $\mathcal{B}' \subset \mathcal{B}$  induces a bijection on connected components. Suppose that  $\mathcal{B}'$  is  $H$ -invariant. Then there is a  $H$ -equivariant diffeomorphism  $\partial \mathcal{B} \times I \rightarrow \mathcal{B} \setminus \text{int } \mathcal{B}'$  where the domain is given the product of the given action on  $\partial \mathcal{B}$  and the trivial  $H$ -action on  $I$ . In particular, there is an  $H$ -equivariant isotopy from the inclusion of  $\mathcal{B}' \subset \mathcal{B}$  to an*



*H-equivariant diffeomorphism of  $\mathcal{B}' \rightarrow \mathcal{B}$ , and consequently the restriction of the  $H$ -action to  $\mathcal{B}'$  is linear.*

*Proof.* Clearly, it suffices to prove this in the case when  $\mathcal{B}$  is connected, and hence the action is a linear action of  $H$  on the unit 3-ball  $B$ . In this case  $\mathcal{B}'$  is a sub-ball  $B' \subset B$ . The sub-ball  $B'$  must contain a fixed point for the entire group action; in fact it must contain the origin except when  $H$  is a cyclic or dihedral group. The restriction of the action to a small ball  $B'' \subset B'$  centered at  $p$  is linear. The region  $B \setminus \text{int } B''$  is equivariantly isomorphic to  $S^2 \times I$  with the action being a linear action on  $S^2$  and trivial on  $I$ . Invoking the previous result we see that  $B' \setminus B''$  is itself equivariantly isomorphic to a product, which means that there is an equivariant isotopy from  $B'$  to the ball  $B''$ , establishing that the action on  $B'$  is equivariantly diffeomorphic to a linear action.  $\square$

### 13.1.1 Connected sum with linear actions on $S^3$ .

**Corollary 13.8.** *Let  $H \times S^3 \rightarrow S^3$  be a linear action and let  $B \subset S^3$  be an  $H$ -invariant ball. (We assume only that  $B$  is smooth, not that it has any special geometric properties with respect to the standard metric on  $S^3$ .) Then there is a diffeomorphism from  $\psi: S^3 \rightarrow S^3$  carrying the given linear action to a linear action with the property that  $\psi(B)$  is an invariant hemisphere in  $S^3$ . In particular, there is an orientation-reversing,  $H$ -equivariant diffeomorphism  $B \rightarrow S^3 \setminus \text{int } B$  that is the identity on  $\partial B$ .*

*Proof.* We use the usual round metric on  $S^3$ . It is  $H$ -invariant. Since the action of  $H$  on  $S^3$  is linear and leaves invariant a sub-ball, the action fixes a point of  $B$ , say  $b \in B$ . It follows that the  $H$ -action fixes the linear subspace spanned by  $\pm b$ . Acting by a rotation that sends  $b$  to the unit vector in the last coordinate direction conjugates  $H$  into  $O(3)$ . A small metric ball  $B_1$  centered at  $-b$  is  $H$ -invariant and disjoint from  $B$ , and hence  $B \subset S^3 \setminus \text{int } B_1$ . Stereographic projection from  $-b$  identifies the  $H$ -action on  $S^3 \setminus \text{int } B_1$  with a linear action of  $H$  on a ball in  $\mathbb{R}^3$ . Thus, it follows from Proposition 13.7 that  $B$  is equivariantly isotopic in  $S^3$  to  $S^3 \setminus \text{int } B_1$ . That is to say, there is a diffeomorphism  $S^3 \rightarrow S^3$  commuting with the  $H$ -action and sending  $B$  to  $S^3 \setminus \text{int } B_1$ . This allows us to assume that  $B$  is a metric ball centered at a fixed point of the action of  $H$ . Of course, there is an  $H$ -equivariant diffeomorphism of  $S^3$  to itself that contracts or expands  $B$  radially toward its center until it becomes a hemisphere. This allows us to make  $B$  a hemisphere fixed by the action of  $H \subset O(3)$ .  $\square$

**Proposition 13.9.** *Let  $H \times M \rightarrow M$  be an action of a compact group on a connected and simply connected 3-manifold. Suppose that this action is an  $H$ -equivariant connected sum of linear actions of  $H$  on families of 3-spheres. Then there is a diffeomorphism  $M$  to  $S^3$  transporting the given  $H$ -action to a linear action.*

*Proof.* Suppose that the  $H$ -equivariant connected sum in question comes from  $\mathcal{B} \subset M'$  and the involution  $\psi: \mathcal{B} \rightarrow \mathcal{B}$ . Form a finite graph whose vertices are the connected components of  $M'$  and whose edges are the orbits of the  $\psi$ -action on the

connected components of  $\mathcal{B}$ . The vertices of an edge are the connected components of  $M'$  that contain the  $\psi$ -orbit that goes with the edge. Since  $M$  is connected and simply connected, it follows that this graph is a tree. Hence, there is a vertex of order one; i.e., there is a connected component  $M'_0$  of  $M'$  that meets  $\mathcal{B}$  in a single ball,  $B_0$ . Let  $B_1 = \psi(B_0)$  and let  $M'_1$  be the component of  $M'$  containing  $B_1$ . One possibility is that  $B_1$  is in the same  $H$ -orbit as  $B_0$ , say  $B_1 = \tau \cdot B_0$  for some  $\tau \in H$ . Let  $M'_1 = \tau M'_0$ . Then  $M'_1 \cap \mathcal{B} = B_1$ . Thus,  $M$  is the connected sum of  $M'_0$  and  $M'_1$  along  $\psi: B_0 \rightarrow B_1$ . Let  $H'_0$  be the stabilizer of  $M'_0$  (which is also the stabilizer of  $M'_1$ ). Then  $H$  is generated by  $H'_0$  and  $\tau$ . Since the action of  $H'_0$  on each of  $M'_0$  and  $M'_1$  is linear, it follows from Corollary 13.8 that  $M'_i \setminus \text{int } B_i$  is  $H'_0$ -equivariantly diffeomorphic to the linear action on  $B_i$ . It is now easy to see that the action of  $H$  on the connected sum is equivariantly diffeomorphic to the linear action of  $H$  on  $S^3$  given by  $c \cdot L: H \rightarrow O(4)$  where  $L: H \rightarrow O(3)$  is the representation that gives the linear action of  $H$  on  $\partial B_0$  and  $c: H \rightarrow \mathbb{Z}/2\mathbb{Z}$  is the homomorphism with kernel  $H'_0$  and  $\mathbb{Z}/2\mathbb{Z}$  is embedded in  $O(4)$  centralizing  $L(H)$  and  $O(3)$ . This shows that the action is linear in this case.

The other possibility is that  $B_1$  is not contained in the  $H$ -orbit of  $B_0$ . In this case since  $M'_0 \setminus \text{int } B_0$  is  $H'_0$ -equivariantly diffeomorphic to  $B_0$  by a map extending the identity on the boundary, and since  $B_0$  and  $B_1$  are  $H'_0$ -equivariantly diffeomorphic, it follows that forming the connected sum along the  $H$ -orbit on  $B_0$  and  $B_1$  does not change the  $H$ -equivariant diffeomorphism type.

A standard induction on the number of 3-balls in the connected sum then gives the result.  $\square$

### 13.2 Actions on Canonical Neighborhoods

Regions of sufficiently large scalar curvature in a time-slice  $M_t$  of the Ricci flow have canonical neighborhoods. Our goal here is to show that there is an equivariant version of this result. One type of canonical neighborhood is an  $\epsilon$ -neck. Recall that an  $\epsilon$ -neck in a 3-manifold  $M$  is an embedding  $\psi: S^2 \times (-\epsilon^{-1}\epsilon^{-1}) \rightarrow M$  so that there is a positive constant  $R$  with the property that the pull back of  $R$  times the Riemannian metric on  $M$  is within  $\epsilon$  in the  $C^{[1/\epsilon]}$ -topology to the product of the round metric of curvature 1 on  $S^2$  and the standard metric on the interval  $(-\epsilon^{-1}, \epsilon^{-1})$ . The central 2-sphere of the neck is the image of  $S^2 \times \{0\}$ , and a point is said to be *at the center of the neck* if it lies in the central 2-sphere of the neck. Notice that on an  $\epsilon$ -neck  $N$  there is a 2-dimensional distribution of 2-planes of maximal sectional curvature. We denote by  $\mathcal{L}_N$  the line field on  $N$  orthogonal to this distribution. Suppose that we have a point  $x$  contained in the middle half of an  $\epsilon$ -neck. Suppose that  $\gamma$  and  $\gamma'$  are geodesics ending at  $x$  whose lengths, when measured with respect to the rescaled metric with  $R(x) = 1$ , are at least  $10^3$ . Then the angle at  $x$  between  $\gamma$  and  $\gamma'$  is within  $5 \cdot 10^{-3}$  either of 0, if  $\gamma$  and  $\gamma'$  lie on the same side of the 2-sphere factor of the  $\epsilon$ -neck containing  $x$ , or of  $\pi$ , if they lie on opposite sides.

### 13.2.1 Actions on $\kappa$ -solutions

One source of models for regions of large scalar curvature in a Ricci flow with surgery are  $\kappa$ -solutions. Let us recall the definitions and main results from Section 9 of [21]. A 3-dimensional  $\kappa$ -solution is a Ricci flow defined for  $-\infty < t \leq 0$  of complete, non-flat, orientable 3-manifolds of bounded, non-negative sectional curvature. Furthermore, it is required to be  $\kappa$ -non-collapsed in the sense that given any ball  $B(x, r, t)$  for which the sectional curvatures are bounded on the parabolic neighborhood  $P(x, r, t, -r^2)$  by  $r^{-2}$ , we have  $\text{Vol } B(x, t, r) \geq \kappa r^3$ . According to Proposition 9.83 of [21] any non-compact  $\kappa$ -solution either has strictly positive curvature, is isometric to  $S^2 \times \mathbb{R}$ , or is double covered by  $S^2 \times \mathbb{R}$  where the covering transformation is the product of the antipodal action on  $S^2$  and the involution on  $\mathbb{R}$  interchanging the ends. By Theorem 9.89 of [21] each time-slice of a compact  $\kappa$ -solution of positive curvature has a round metric.

Here is an initial classification of compact group actions on a  $\kappa$ -solution in the easy cases.

**Proposition 13.10.** *1. Any compact group of isometries of a compact  $\kappa$ -solution is finitely covered by a linear action on  $S^3$ .*

*2. Any isometric action of a compact group on  $S^2 \times \mathbb{R}$  is the product of a linear action on  $S^2$  and a linear action on  $\mathbb{R}$ .*

*3. Let  $\tau$  be the involution of  $S^2 \times \mathbb{R}$  that is the product of the antipodal map on  $S^2$  and the reflection in the origin on  $\mathbb{R}$ . Any isometric action on  $(S^2 \times \mathbb{R})/\tau$  is induced from a linear action on  $S^2$ .*

*Proof.* Since compact  $\kappa$ -solutions have round metrics, the first item is immediate.

An action on  $S^2 \times \mathbb{R}$  must preserve the family of 2-spheres and the perpendicular family of lines. Thus, projecting to the factors determines isometric actions on  $S^2$  and on  $\mathbb{R}$ . It is clear that the action is the product of these actions on the factors.

In the last case an action of  $H$  on the quotient lifts to an isometric action of  $\tilde{H}$  on  $S^2 \times \mathbb{R}$ , where  $\tilde{H}$  fits into the exact sequence:

$$0 \rightarrow \mathbb{Z}/2\mathbb{Z} \rightarrow \tilde{H} \rightarrow H \rightarrow \{1\}.$$

The kernel is central in fact is split by the orientation character on the line, so that  $\tilde{H} = H \times \mathbb{Z}/2\mathbb{Z}$ , where the second factor is generated by  $-\text{Id} \times R_0 \subset O(3) \times \text{Iso}(\mathbb{R})$ , and  $R_0$  is the antipodal map of  $S^2$ . The result follows immediately from the first case.  $\square$

Now let us turn to the most interesting case.

**Proposition 13.11.** *If  $(M, g(t))$ ,  $-\infty < t \leq 0$ , is a non-compact  $\kappa$ -solution of positive curvature and if  $(M, g(0))$  is invariant under a compact group  $H$ , then the  $H$ -action on  $M$  is equivariantly diffeomorphic to a linear action of  $H$  on  $\mathbb{R}^3$ .*

*Proof.* Since  $(M, g(0))$  has positive curvature, it has a soul which is a point. We claim that it has a soul invariant under the  $H$ -action. To see this, fix a compact  $H$ -invariant subset  $X \subset M$  and consider the set,  $A$ , of all minimal geodesic rays

starting at points of  $x$ . For each such ray  $\gamma$  let  $b_\gamma: M \rightarrow \mathbb{R}$  be the Busemann function for  $\gamma$ ; i.e.,  $b_\gamma(x) = \lim_{t \rightarrow \infty} d(x, \gamma(t)) - t$ . The super level sets of  $b_\gamma$ , denoted  $S_a(b_\gamma) = \{x \in M | b_\gamma(x) \geq a\}$ , are totally geodesically convex in the sense that if  $\gamma'$  is a geodesic arc with endpoints in  $S_a(b_\gamma)$ , then  $\gamma' \subset S_a(b_\gamma)$ . Now we consider  $f = \inf_{\gamma \in A} b_\gamma$ . This is an  $H$ -invariant function whose super level sets,  $S_a(f)$ , are totally geodesically convex. We claim that each super level set of  $f$  is compact. We can assume that  $a \leq -\text{diam}(X)$ . First notice that if  $x \in X$  then  $f(x) \geq -\text{diam}(X)$ , so that  $X$  is in the super level set  $S_a(f)$ . If  $S_a(f)$  is not compact, then there is a sequence,  $x_n \in S_a(f)$  tending to infinity. Fix a point  $x_0 \in X$  and take (minimal) geodesics  $\mu_n$  from  $x_0$  to  $x_n$ . Then  $\mu_n \subset S_a(f)$ . Passing to a subsequence we can arrange that the  $\mu_n$  converge to a geodesic ray  $\gamma$  from  $x_0$ . But clearly  $f(\gamma(t)) \leq b_\gamma(\gamma(t)) = -t$ , which contradicts the fact that  $\gamma \subset S_a(f)$  for some  $a > -\infty$ .

Let  $a_0$  be the maximum value of  $f$  and set  $C = f^{-1}(a_0)$ . This is a non-empty, compact, totally geodesic  $H$ -invariant subset of  $M$ . Lemma 62 in Section 11.4 and the argument immediately following it in [29] show that in this case of strictly positive sectional curvature  $S_{a_0}(f)$  is a point, and hence it is an  $H$ -invariant soul  $p_0$  for  $M$ .

Now consider  $r = d(p_0, \cdot)$ . This is an  $H$ -invariant function. For  $b > 0$  sufficiently small  $\nabla(r)$  is a smooth, non-vanishing  $H$ -invariant vector field on  $B(p_0, b) \setminus \{p_0\}$ . Furthermore, if  $b > 0$  sufficiently small the exponential map identifies the action of  $H$  on the closed ball  $\overline{B(p_0, b)}$  with a linear action on the closed ball of radius  $b$  in the tangent space  $T_{p_0}X$ . The general soul theory implies that there is a smooth vector field agreeing with  $\nabla r$  on  $\overline{B(p_0, b)}$  with the property that  $d(p_0, \cdot)$  is increasing along every flow line of this vector field (except at  $p_0$ ). Averaging the vector field over  $H$  allows us to assume that in addition it is  $H$ -invariant. As such it determines an  $H$ -invariant diffeomorphism:

$$M \setminus B(p_0, b) \cong \partial \overline{B(p_0, b)} \times [b, \infty),$$

and hence establishes an equivariant diffeomorphism from  $H \times M \rightarrow M$  to a linear action of  $H$  on  $\mathbb{R}^3$  given by the differential of the action at the  $H$ -invariant soul  $p_0$ .  $\square$

**Corollary 13.12.** *Let  $M$  be the final time-slice of a non-compact  $\kappa$ -solution of positive curvature. Suppose that  $H \times M \rightarrow M$  is an isometric action of a compact group, and let  $B \subset M$  be an invariant, closed 3-ball. Then the restriction of the action of  $H$  to  $B$  is equivariantly diffeomorphic to a linear action.*

*Proof.* Since by Proposition 13.11 the action on  $M$  is equivariantly diffeomorphic to a linear action on  $\mathbb{R}^3$ , there is a compact ball  $B_0$  containing  $B$  on which the action is equivariantly diffeomorphic to a linear action. The result is then immediate from Proposition 13.7.  $\square$

### 13.2.2 Actions on the standard solution

The other possible models for regions of large scalar curvature in a Ricci flow with surgery come from the standard solution. According to Section 12 of [21] this is

a Ricci flow  $(M, h_0(s))$ ,  $0 \leq s < 1$ . The Riemannian manifold  $(M, h_0(0))$  has an isometric action of  $O(3)$  which fixes a point  $q_0$ , called the *tip*, and which is equivariantly diffeomorphic to the natural linear action of  $O(3)$  on  $\mathbb{R}^3$ . This action<sup>10</sup> preserves the metric  $h_0(s)$  for every  $0 \leq s < 1$ . Any compact group action on any time-slice,  $(M, h_0(s))$ , is a subgroup of this  $O(3)$ -action and in particular fixes  $q_0$  and is equivariantly diffeomorphic to a linear action on  $\mathbb{R}^3$ .

Just as in the previous case, this leads to the following result for balls in a time-slice of the standard solution.

**Corollary 13.13.** *Let  $(M, h_0(s))$  be a time-slice of the standard solution. Suppose that  $H$  is a compact group acting isometrically on  $(M, h_0(s))$  and that  $B \subset M$  is an  $H$ -invariant 3-ball. Then the action of  $H$  on  $B$  is equivariantly diffeomorphic to a linear action.*

### 13.2.3 Canonical Neighborhoods in $\kappa$ -solutions and the standard solution

Here is a result that was established in Section 9 of [21] about canonical neighborhoods for non-compact  $\kappa$ -solutions.

**Proposition 13.14.** *For any  $\epsilon > 0$  there is  $C_0 = C_0(\epsilon) < \infty$ , with  $C_0 > 10\epsilon^{-1}$  such that the following holds. Suppose that  $(M, h(t))$ ,  $-\infty < t \leq 0$ , is a non-compact  $\kappa$ -solution of positive curvature. Then any point  $x \in (M, h(0))$  is either the center of an  $\epsilon/3$ -neck or there is a submanifold  $\mathcal{C} \subset (M, h(0))$  with the following properties:*

1.  $\mathcal{C}$  is diffeomorphic to an open 3-ball.
2. There an  $\epsilon/4$ -neck  $N(\mathcal{C})$  contained in  $\mathcal{C}$  and containing the end of  $\mathcal{C}$ .
3. The central 2-sphere  $\Sigma$  of  $N$  bounds a compact submanifold  $\mathcal{C}_0$  of  $\mathcal{C}$ , called the core of  $\mathcal{C}$ .
4. The metric has positive sectional curvature at every point of  $\mathcal{C}$ .
5. Given any points  $y, z \in \mathcal{C}$  and 2-planes  $P_y \subset T_y M$  and  $P_z \subset T_z M$  the ratio of the sectional curvature in the  $P_y$ -direction and that in  $P_z$  direction is between  $C_0$  and  $C_0^{-1}$ .
6. The diameter of  $\mathcal{C}$  is at most  $C_0 R^{-(1/2)}(x)$ .
7. The core  $\mathcal{C}_0$  contains  $x$ .

**Definition 13.15.** Given  $\epsilon > 0$  we fix  $C_0$  as in the proposition and we call a submanifold  $\mathcal{C}$  of the final time-slice of a non-compact  $\kappa$ -solution of positive curvature satisfying the conclusions 1 – 6 of the above proposition an  $\epsilon$ -cap. We define a *twisted  $\epsilon$ -cap* to be the quotient  $\mathcal{T}$  of  $S^2 \times (-C_0, C_0)$  by the involution  $\tau$  that is the product

<sup>10</sup>Actually, this result is only stated for  $SO(3)$  in [21] but by uniqueness result established in Section 12 of [21], given the initial metric, it holds for  $O(3)$

of the antipodal map on  $S^2$  with the map  $x \mapsto -x$  on the interval. Notice that there is an  $\epsilon/4$ -neck  $N \subset \mathcal{T}$  with compact complement and every point of  $N$  is the center of an  $\epsilon$ -neck in  $S^2 \times \mathbb{R}/\tau$ .

Clearly, any point of  $S^2 \times \mathbb{R}$  is the center of an  $\epsilon/3$ -neck and any point of  $S^q \times \mathbb{R}/\tau$  is either the center of an  $\epsilon/3$ -neck or is contained in a twisted  $\epsilon$ -cap.

Given  $\epsilon > 0$  for any  $s_0 < 1$ , there is  $C_1 = C_1(\epsilon, s_0)$  such that for any  $s \leq s_0$  the ball  $B(q_0, s, C_1)$  centered at the tip of the  $s$  time-slice of the standard solution has positive curvature and contains an  $\epsilon/4$ -neck whose complement is compact. Denoting by  $\Sigma$ , the central 2-sphere of of this neck, the compact submanifold of  $B(q_0, s, C_1)$  bounded by  $\Sigma$  is the *core* of this  $(C_1, \epsilon)$ -cap.

### 13.2.4 Standard Models for regions of large scalar curvature

In Section 16 of [21] the following is established (though not explicitly stated):

**Theorem 13.16.** (*Canonical neighborhood theorem*) *Given  $\epsilon > 0$  and  $\delta > 0$  sufficiently small there is an  $s_0 < 1$  and a function  $r(t)$  such that the following holds for  $C = C(\epsilon, s_0) = \max(C_0(\epsilon), C_1(s_0, \epsilon))$ . Let  $x \in M_t$  be a point of the  $t$  time-slice of a Ricci flow with surgery. If  $R(x) \geq r^{-2}(t)$  then, setting  $g'_t = R(x)g_t$ , one of the following holds:*

1. *There are a  $\kappa$ -solution  $(N, h(t))$ ,  $-\infty < t \leq 0$ , a point  $(p, 0)$  in its final time-slice with  $R(p, 0) = 1$ , and a smooth embedding  $\varphi: B_{h(0)}(p, C) \rightarrow M_t$  with  $\varphi(p) = x$  with  $\varphi^*g'_t$  within  $\delta$  in the  $C^{[1/\delta]}$ -topology of the restriction of  $h(0)$  to  $B_{h(0)}(p, C)$ .*
2. *There are  $s \leq s_0$  and a smooth embedding  $\varphi$  of the ball  $B_1(s) = B(q_0, s, C)$  centered at the tip of the  $s$  time-slice of the standard solution into  $M_t$  containing  $B_{g'_t}(x, 10\epsilon^{-1})$  with the property that  $\varphi^*g'_t$  is within  $\delta$  in the  $C^{[1/\delta]}$ -topology to the restriction of  $R(\varphi^{-1}(x))h_0(s)$  to  $B_1(s)$ .*

Now we can use this result to extract four kinds of models for regions of large scalar curvature.

**Corollary 13.17.** *Fix  $\epsilon > 0$  sufficiently small. Let  $s_0$  and  $C_0, C_1$  be fixed as in the previous theorem. Then there is a positive function  $r(t)$  such that the following holds. If  $x \in M_t$  has  $R(x) \geq r^{-2}(t)$ , then, setting  $g'_t = R(x)g_t$ , one of the following holds:*

1. *The connected component of  $M_t$  containing  $x$ , with its metric rescaled by  $R(x)$ , is within  $\epsilon$  in the  $C^{[1/\epsilon]}$ -topology to a round metric of constant curvature  $1/3$ .*
2.  *$x$  is the center of an  $\epsilon$ -neck in  $M_t$ .*
3. *There is a twisted  $\epsilon$ -cap whose core contains  $x$ .*

4. There is a non-compact  $\kappa$ -solution  $(N, h(t))$ ,  $-\infty < t \leq 0$ , a point  $(p, 0) \in (N, h(0))$  with  $R(p, 0) = 1$ , an  $\epsilon$ -cap  $\mathcal{C}(p)$  in  $(N, h(0))$  whose core contains  $p$ , and an embedding  $\varphi: \mathcal{C}(p) \rightarrow M_t$  sending  $p$  to  $x$ , such that  $\varphi^*g'_t$  is within  $\epsilon$  in the  $C^{[1/\epsilon]}$ -topology to the restriction of  $h(0)$  to  $\mathcal{C}(p)$ . Furthermore, the sectional curvature of  $M_t$  is positive on the image  $\varphi(\mathcal{C}(p))$ .
5. There are  $s \leq s_0$  and a smooth embedding  $\varphi$  of the ball  $B_1(s) = B(q_0, s, C_1)$  in the  $s$  time-slice of the standard solution into  $M_t$  containing  $B_{g'_t}(x, 10\epsilon^{-1})$  with the property that  $\varphi^*g'_t$  is within  $\epsilon$  in the  $C^{[1/\epsilon]}$ -topology to the restriction of  $R(\varphi^{-1}(x))h_0(s)$  to  $B_1(s)$  and the sectional curvatures on the image of  $\varphi$  are positive.

*Proof.* We take  $\delta \ll \epsilon$  and then fix  $r(t)$  depending on  $\epsilon, s_0, \delta$ . Three remarks are in order: (i) When the model is a ball  $B_{h(0)}(p, C)$  in a non-compact  $\kappa$ -solution and  $(p, 0)$  is the center of an  $\epsilon/3$ -neck in the  $\kappa$ -solution. In this case, since  $\delta \ll \epsilon$  it follows that  $x$  is the center of an  $\epsilon$ -neck in  $M_t$ ; (ii) since  $R(p, 0) = 1$  the sectional curvatures on  $\epsilon$ -caps  $\mathcal{C}(p)$  in non-compact  $\kappa$  solutions bounded below by a positive constant. Likewise in Case 5 the sectional curvatures on  $B(q_0, s, C_1)$  are bounded below by a positive constant independent of  $s \leq s_0$ . Taking  $\delta$  sufficiently small, it will then be true that in Cases 3 and 5 that the sectional curvatures on the image  $\varphi(\mathcal{C}(p))$  will be positive.  $\square$

**Definition 13.18.** Fixing  $\epsilon > 0$ . We say that  $\mathcal{C}(p)$  and  $B_{h_0(s)}(q_0, s, C_1)$  satisfying the conclusions of the previous theorem are *model strong  $\epsilon$ -caps*, and given  $x$ , the image of a map  $\varphi$  from one of these model strong  $\epsilon$ -caps as in the third or fifth item of the corollary is called a *strong  $\epsilon$ -cap neighborhood of  $x$* , or a *strong  $\epsilon$ -cap*. The neighborhood in the fourth item is called a *twisted  $\epsilon$ -cap*.

### 13.2.5 Limits of group actions

Recall the notion of a geometric limit of a sequence of based, Riemannian manifolds: A sequence of based, complete Riemannian manifolds  $\{(M_n, g_n, x_n)\}_{n \geq 1}$  is said to *converge geometrically* to a based complete Riemannian manifold  $(M_\infty, g_\infty, x_\infty)$  if there is an increasing sequence of open subsets  $U_n \subset M_\infty$ , each containing  $x_\infty$ , whose union is  $M_\infty$  and, for each  $n$  sufficiently large, a diffeomorphism  $\varphi_n: U_n \rightarrow \varphi_n(U_n) \subset M_n$  with  $\varphi_n(x_\infty) = x_n$  such that the  $\varphi_n^*g_n$  converge to  $g_\infty$ , uniformly in the  $C^\infty$ -topology on every compact subset of  $M_\infty$ .

Here is the main result we need about limits of group actions.

**Proposition 13.19.** *Let  $(M_n, g_n, x_n)$  be a sequence of based, complete Riemannian 3-manifolds and let  $H$  be a compact group. Suppose that for each  $n$ ,  $\psi_n: H \times M_n \rightarrow M_n$  is an effective, isometric group action. We equip  $H$  with an invariant metric, denoted  $d_H$ . We suppose that the following two conditions hold:*

1. *There is  $R < \infty$  such that for every  $n \geq 1$  and every element  $h \in H$ ,  $d(x_n, hx_n) < R$ .*

2. For any  $S > 0$  and  $\epsilon > 0$ , there is  $\delta = \delta(\epsilon, S) > 0$  such that for all  $n$  sufficiently large and for any  $x \in B(x_n, S)$  and any  $h, h' \in H$  with  $d_H(h, h') < \delta$  we have  $d(hx, h'x) < \epsilon$ .

If the based Riemannian manifolds converge geometrically to a limit  $(M_\infty, g_\infty, x_\infty)$ , then, after passing to a subsequence, there are:

1. an effective, isometric action  $\psi_\infty: H \times M_\infty \rightarrow M_\infty$ ,
2. an increasing sequence of  $H$ -invariant open sets  $V_n \subset M_\infty$ , each containing  $x_\infty$ , whose union is  $M_\infty$ , and
3. for each  $n$  an  $H$ -equivariant embedding  $\bar{\varphi}_n: V_n \rightarrow M_n$ , with  $d(\bar{\varphi}_n(x_\infty), x_n)$  tending to 0 as  $n \rightarrow \infty$ , such that the  $\bar{\varphi}_n^* g_n$  converge to  $g_\infty$ , uniformly in the  $C^\infty$ -topology on every compact subset of  $M_\infty$ .

*Proof.* One first goal is to pass to a subsequence and construct a limiting action of  $H$  on  $M_\infty$ . Let us consider a single element  $h \in H$ . The distance  $d(x_n, \psi_n(h)x_n)$  is bounded independent of  $n$ , and hence for all  $n$  sufficiently large  $\varphi_n^{-1}$  is defined on  $\psi_n(h)x_n$  and the distance in  $M_\infty$  between  $\varphi_n^{-1}(\psi_n(h)x_n)$  and  $x_\infty$  is bounded independent of  $n$ . Passing to a subsequence we arrange that the sequence  $\varphi_n^{-1}\psi_n(h)x_n$  converges to a point  $y(h) \in M_\infty$ . Passing to a further subsequence we can arrange that the differentials of  $\varphi_n^{-1}\psi_n(h)\varphi_n$  at  $x_\infty$  converge to an isometry from  $T_{x_\infty}M_\infty \rightarrow T_{y(h)}M_\infty$ . Since  $\psi_n(h)$  is an isometry and the  $\varphi_n$  are converging uniformly on compact sets to isometries, it follows that the  $\varphi_n^{-1}\psi_n(h)\varphi_n$  are converging uniformly on compact subsets of  $M_\infty$  to an isometry, which we call  $\psi_\infty(h)$ , of  $M_\infty$ , and this isometry is determined by  $y(h)$  and the limiting differential at  $x_\infty$ .

There is a finite set of elements  $D \subset H$  that generate a dense subgroup of  $H$ . Apply the result established in the previous paragraph to pass to a subsequence so that for every element of  $D$  the action of this element on  $M_n$  converges, uniformly on compact subsets, to an isometry of  $M_\infty$ . Then for every product  $p = d_1 \cdots d_k$  of elements of  $D$ , the actions of  $\psi_n(p)$  on  $M_n$ , converge uniformly on compact subsets, to the product of the limiting actions  $\psi_\infty(d_i)$  on  $M_\infty$ . That is to say, letting  $G(D) \subset H$  be the subgroup generated by  $D$ , there is an action  $\psi_\infty$  of  $G(D)$  on  $M_\infty$  and for each  $g \in G(D)$  the diffeomorphisms  $\varphi_n^{-1}\psi_n(g)\varphi_n$  converge uniformly on compact subsets to the isometry  $\psi_\infty(g)$ .

Now we extend  $\psi_\infty$  to action of all of  $H$  on  $M_\infty$ . Given  $h \in H$  there is a sequence  $g_i \in G(D)$  converging to  $h$ . The uniform continuity of the  $\psi_n$  on compact sets implies that the  $\psi_\infty(g_i)$  converge, uniformly on compact subsets of  $M_\infty$  to an isometry  $\psi_\infty(h)$ , which as the notation suggests, depends only on  $h$ . This defines an extension of  $\psi_\infty$  to an isometric action on  $M_\infty$ . Using the uniform continuity of the  $\psi_n$  on compact sets we see that for any  $h \in H$  the diffeomorphisms  $\varphi_n^{-1}\psi_n(h)\varphi_n$  converge uniformly on compact subsets to  $\psi_\infty(h)$ . The uniform continuity of the action in the sequence implies that the killing fields associated with unit vectors in the Lie algebra of  $H$  have uniformly bounded length under  $d\psi_n$ . Hence, passing to a subsequence we can arrange that the maps of the Lie algebra into the vector fields



on  $M_n$  converge uniformly on compact sets to a map of this Lie algebra into killing vector fields on  $M_\infty$ . Thus, the limiting action is a smooth action of  $H$  on  $M_\infty$ .

This completes the construction of the limiting effective, isometric action  $\psi_\infty$  of  $H$  on  $M_\infty$  with the property that given any compact set  $X \subset M_\infty$  and any  $\delta > 0$  for any  $h \in H$  for all  $n$  sufficiently large the restrictions of  $\varphi_n^{-1}\psi_n(h)\varphi_n$  and  $\psi_\infty(h)$  to  $X$  are within  $\delta$  in the  $C^\infty$ -topology of each other. It remains to replace the approximating diffeomorphisms  $\varphi_n: U_n \rightarrow M_n$  by equivariant diffeomorphism (shrinking the  $U_n$  but keeping their union equal to all of  $M_\infty$ ). For any compact subset  $X \subset M_\infty$ , the subset  $H \cdot X$  is compact and hence is contained in  $U_n$  for all  $n$  sufficiently large. Hence, for all  $n$  sufficiently large, for every  $h \in H$  the map  $\psi_n(h)^{-1}\varphi_n\psi_\infty(h)$  is defined on  $X$ . Furthermore, as  $h$  varies these maps are all close to each other in the  $C^\infty$ -topology on  $X$ , with the error going to zero (on  $X$ ) as  $n \mapsto \infty$ . Thus, passing to a subsequence we can suppose that eventually for each compact subset  $X$  of  $M_\infty$ , for all  $n$  sufficiently large, the restrictions of  $\psi_n(h)^{-1}\varphi_n\psi_\infty(h)$  to  $X$  are arbitrarily close together. Since the metrics on the union of the images of  $X$  under these maps are converging to the metric of  $g_\infty$  on this compact set, for all  $n$  sufficiently large we can take the center of mass of this set of points (parametrized by  $h \in H$  and integrated with respect to a Haar measure on  $H$  of total volume 1) as in [10]. This center of mass map determines a map  $\bar{\varphi}_n$  defined on  $X$ . By construction  $\bar{\varphi}_n$  is  $H$ -equivariant, and is  $C^\infty$ -close to  $\varphi_n$ , with the error estimates going to zero as  $n \rightarrow \infty$ . In particular,  $\bar{\varphi}_n$  is an embedding for all  $n$  sufficient large. As  $n \mapsto \infty$  these embeddings converge smoothly on  $X$  to an isometry in the sense that  $\bar{\varphi}_n^*g_n|_X$  converges in the  $C^\infty$ -topology to  $g_\infty|_X$ .

It is clear that  $d(\bar{\varphi}_n(x_\infty), x_n)$  tends to zero as  $n \rightarrow \infty$ .  $\square$

### 13.2.6 Group actions on strong $\epsilon$ -caps and twisted $\epsilon$ -caps

**Lemma 13.20.** *Let  $\mathcal{C}$  be a strong  $\epsilon$ -cap centered at  $x \in M_t$ . Suppose that  $H \times M_t \rightarrow M_t$  is an isometric action of a compact group. Let  $H_0$  be the set of elements in  $h \in H$  with the property that  $h\mathcal{C}_0 \cap \mathcal{C}_0 \neq \emptyset$ . Then  $H_0$  is a subgroup of finite index and there is an  $H_0$ -invariant 2-sphere in the  $\epsilon/4$ -neck  $N(\mathcal{C})$  that is isotopic in  $N(\mathcal{C})$  to the central 2-sphere of the neck.*

*Proof.* Let  $\Sigma$  be the central 2-sphere of  $N(\mathcal{C})$ . We claim that for any  $h \in H_0$  we have  $h \cdot \Sigma \cap \Sigma \neq \emptyset$ . For suppose  $h\Sigma$  is disjoint from  $\Sigma$ . Then either  $h\mathcal{C}_0 \subset \mathcal{C}_0$ ,  $\mathcal{C}_0 \subset h\mathcal{C}_0$  or  $\mathcal{C}_0 \cap h\mathcal{C}_0 = \emptyset$ . The last violates the assumption that  $h \in H_0$ . Neither of the first two is possible since the volume of  $h\mathcal{C}_0$  equals the volume of  $\mathcal{C}_0$ . This proves that  $h \cdot \Sigma \cap \Sigma \neq \emptyset$  for every  $h \in H_0$ .

Let  $d(\Sigma)$  be the diameter of  $\Sigma$ . It follows that if  $h_1, h_2 \in H_0$ , then  $h_1h_2\Sigma$  is contained in the  $2d(\Sigma)$  neighborhood of  $\Sigma$ , and in particular is contained in  $N(\mathcal{C})$ . It follows easily that  $h_1h_2\mathcal{C}_0 \cap \mathcal{C}_0 \neq \emptyset$ . This shows that  $H_0$  is closed under products; it is clearly closed under taking inverses, and hence is a subgroup of  $H$ , obviously of finite index.

Now we consider the unit vector field on  $N(\mathcal{C})$  that generates the line field  $\mathcal{L}_{N(\mathcal{C})}$  and points toward  $\mathcal{C}_0$ . This vector field is invariant under  $H_0$ . Using the flow

generated by this vector field we define a product structure on an open subset  $U$  of  $N(\mathcal{C})$  containing the middle half of  $N(\mathcal{C})$ :

$$\Sigma \times (-a, a) \rightarrow U.$$

Now for each  $h \in H_0$  the 2-sphere  $h \cdot \Sigma$  is the image under this product structure of the graph of a smooth function  $f_h: \Sigma \rightarrow (-a, a)$ . Let  $\bar{f}$  be the average of these functions over  $h \in H_0$ . We claim that the image of the graph of  $\bar{f}$  is an  $H_0$ -invariant 2-sphere as required.

The fact that action of  $H_0$  on  $\Sigma \times (-a, a)$  preserves the unit vector field in the  $t$  direction means that it is given by  $h \cdot (\sigma, t) = \bar{h}(\sigma), f_h(\bar{h}(\sigma) + t)$ , where  $h \rightarrow \bar{h}$  is an action of  $H$  on  $\Sigma$ . From the group law it follows that

$$f_{h_1 h_2}(\bar{h}_1 \bar{h}_2 \sigma) = f_{h_1}(\bar{h}_1 \bar{h}_2 \sigma) + f_{h_2}(\bar{h}_2 \sigma). \quad (13.1)$$

Applying this with  $h_2 = h_1^{-1}$ , and using the fact that  $f_e = 0$ , we have:

$$f_h(\sigma) = -f_{h^{-1}}(\bar{h}^{-1} \sigma). \quad (13.2)$$

Now

$$h(\sigma, \bar{f}(\sigma)) = (\bar{h}\sigma, f_h(\bar{h}\sigma) + \int_{g \in H_0} f_g(\sigma) dg),$$

whereas

$$(\bar{h}\sigma, f(\bar{h}\sigma)) = (\bar{h}\sigma, \int_{g \in H_0} f_g(\bar{h}\sigma)).$$

Thus, to show that the graph of  $\bar{f}$  is  $H$ -invariant we need to show:

$$\int_{g \in H_0} f_g(\bar{h}\sigma) dg = f_h(\bar{h}\sigma) + \int_{g \in H_0} f_g(\sigma) dg.$$

But, applying Equation 13.1 with  $h_1 = g$  and  $h_2 = g^{-1}h$ , and using the fact that the volume of  $H_0$  is one, we have

$$\begin{aligned} \int_{g \in H_0} f_g(\bar{h}\sigma) dg &= \int_{g \in H_0} f_h(\bar{h}\sigma) dg - \int_{g \in H_0} f_{g^{-1}h}(\bar{g}^{-1} \bar{h}\sigma) dg, \\ &= f_h(\bar{h}\sigma) - \int_{g \in H_0} f_{g^{-1}h}(\bar{g}^{-1} \bar{h}\sigma) dg. \end{aligned}$$

Now using Equation 13.2 we have

$$- \int_{g \in H_0} f_{g^{-1}h}(\bar{g}^{-1} \bar{h}\sigma) dg = \int_{g \in H_0} f_{h^{-1}g}(\sigma) dg,$$

which by the invariance of the measure on  $H_0$  under left multiplication is equal to

$$\int_{g \in H_0} f_g(\sigma) dg,$$

completing the proof.  $\square$

Now we are ready to show that the actions on the truncated versions of strong  $\epsilon$ -caps are linear.

**Proposition 13.21.** *Fix an integer  $N$ . The following holds for all  $\epsilon > 0$  sufficiently small, how small depending on  $N$ .*

1. *Suppose that  $\mathcal{C}$  is a strong  $\epsilon$ -cap in a closed Riemannian manifold  $M$ , that  $H \times M \rightarrow M$  is an isometric action of a compact group, with  $H$  having at most  $N$  connected components, and  $h \cdot \mathcal{C}_0 \cap \mathcal{C}_0 \neq \emptyset$  for all  $h \in H$ . Let  $X \subset \mathcal{C}$  be a compact  $H$ -invariant submanifold with boundary a 2-sphere in  $N(\mathcal{C})$  isotopic in  $N(\mathcal{C})$  to the central 2-sphere. Then the action  $H \times X \rightarrow X$  is equivariantly diffeomorphic to a linear action on the 3-ball.*

2. *Suppose that  $\mathcal{C}$  is a twisted  $\epsilon$ -cap and that  $X \subset \mathcal{C}$  is a compact  $H$ -invariant submanifold whose boundary is a 2-sphere in  $N(\mathcal{C})$  isotopic in  $N(\mathcal{C})$  to the central 2-sphere. Then  $H \times X \rightarrow X$  is double covered by an action  $\tilde{H} \times (S^2 \times I) \rightarrow S^2 \times I$  that is the product of a linear action on  $S^2$  and a linear action on the interval.*

*Proof.* Fix  $N < \infty$  and suppose that there is no  $\epsilon$  as required in 1. Then there is a sequence of  $\epsilon_n \rightarrow 0$  as  $n \rightarrow \infty$  and a sequence of counter examples  $H_n \times X_n \rightarrow X_n$  for contained in strong  $\epsilon_n$ -caps. Passing to a subsequence we can suppose that the model  $\epsilon$ -caps either are all contained in non-compact  $\kappa$ -solutions of positive curvature or are all contained in the standard solution. In the first case, let  $p_n$  be a soul for  $(N_n, h_n(0))$ . It is contained in the model for  $\mathcal{C}_n$  and we denote by  $x_n$  its image under the map from the model. We rescale the metrics so that  $R(p_n, 0) = 1$  and  $R(x_n) = 1$ . Then the  $(X_n, x_n, g'_n)$  converge as  $n \rightarrow \infty$  geometrically to the final time-slice of a non-compact  $\kappa$ -solution.

Each group  $H_n$  has at most  $N$  components and has dimension bounded above by 3, so passing to a subsequence we can assume that all the  $H_n$  are isomorphic. We identify all the  $H_n$  with  $H$ . We claim the two conditions in Proposition 13.19 are satisfied. Fix  $\epsilon_0 > 0$ . Then for all  $n$  sufficiently large  $\epsilon_n < \epsilon_0$ . For any  $\epsilon_0$ -neck in  $X$  with central 2-sphere  $\Sigma$  for any  $h \in H$  we have  $h \cdot \Sigma \cap \Sigma \neq \emptyset$ . Since there is an  $\epsilon_0/2$  neck at a uniformly bounded distance from  $p_n$  it follows that in  $X$  there is an  $\epsilon_0$ -neck at a uniformly bounded distance from  $x_n$ . Since the central 2-sphere of this neck is mapped so as to meet itself, it follows that  $h$  moves  $x_n$  a distance bounded independent of  $h \in H$  and of  $n$ . It is also true that any circle subgroup of  $H$  fixes two points on any central 2-sphere of an  $\epsilon_0$ -neck, and so all such circle groups have fixed points within a uniformly bounded distance of  $x_n$ . This, and the fact that the metrics are converging uniformly on compact sets as  $n \rightarrow \infty$ , imply that Condition 2 in the hypothesis of Proposition 13.19 holds. Hence, both conditions in the hypothesis of this proposition hold.

According to Proposition 13.19 there is a limiting action of  $H$  on the geometric limit. This limit is either a  $\kappa$ -solution or a time-slice of the standard solution. But these limit actions are automatically linear actions on  $\mathbb{R}^3$  so that by Proposition 13.19 for all  $n$  sufficiently large we have an equivariant diffeomorphism from  $X_n$  to an invariant ball in a linear action on  $\mathbb{R}^3$ . But as we have already seen, this implies that for all  $n$  sufficiently large, the action on  $X_n$  is linear. This completes the proof of the first case.

The case of twisted  $\epsilon$ -caps is analogous.  $\square$

### 13.3 Equivariant Ricci flow and linearity of the action on the surgery regions

#### 13.3.1 Step 1: An equivariant version of the Ricci flow with surgery.

Let  $M$  be a compact, orientable 3-manifold and let  $H \times M \rightarrow M$  be a smooth action of a compact group. We fix a Riemannian metric  $g$  on  $M$  that is  $H$ -invariant. Scaling it by a suitably large positive constant allows us to assume that  $g$  is also normalized. We also fix  $\epsilon > 0$  sufficiently small, how small depending on the number of connected components of  $H$  as in Proposition 13.21.

**Proposition 13.22.** *With proper choices one can construct a Ricci flow with surgery  $(\mathcal{M}, G)$  with initial conditions  $(M, g)$  satisfying the conclusions of Theorem 1.2 and an action  $H \times \mathcal{M} \rightarrow \mathcal{M}$  that preserves the levels  $M_t \subset \mathcal{M}$ , acts by isometry on each level, and preserves the flow lines on the smooth part of the Ricci flow with surgery.*

*Proof.* We begin with the  $H$ -equivariant compact Riemannian manifold  $(M, g)$ . Rescaling the metric  $g$  by constant if necessary, we can assume that the initial conditions are normalized. Since the solution to the Ricci flow equation is unique, it follows that the maximal Ricci flow  $(M, g(t))$ ,  $0 \leq t < t_0$ , with this initial data is  $H$ -invariant. At the limiting time, i.e., at the first surgery time, the open subset  $\Omega \subset M$  consisting of all points where the metrics  $g(t)$  converge smoothly to a limiting metric as  $t \rightarrow t_0^-$  is clearly  $H$ -invariant, as is the subset  $\Omega(\rho(t_0)) \subset \Omega$  where the scalar curvature of the limiting metric is at most  $\rho^{-2}(t_0)$ . (Here,  $\rho(t_0) = \bar{\delta}(t_0)r(t_0)$ , the functions on the right-hand side being the ones associated to  $\epsilon$  by Theorem 1.2.) Surgery is done on the ends of  $\Omega$ . We begin by recalling some of the central concepts in understanding these ends and then we show that these concepts have equivariant analogues, eventually leading to a proof that surgery can be done equivariantly.

Recall the notion of an  $\epsilon$ -horn in  $\Omega$ . We equip  $\Omega$  with the limiting metric as  $t \rightarrow t_0^-$ , denoted  $g(t_0)$ . Recall that an  $\epsilon$ -horn  $\mathcal{K}$  is the image of a proper embedding  $S^2 \times [0, \infty) \rightarrow \Omega$  with the following properties:

1. the restriction of the scalar curvature function of  $\Omega$  to goes to  $+\infty$  as we go to infinity in  $\mathcal{K}$ .
2. Each point of  $\mathcal{K}$  is the center of an  $\epsilon$ -neck in  $\Omega$  and the boundary of  $\mathcal{K}$  is a central 2-sphere in an  $\epsilon$ -neck.
3. The image of  $\partial\mathcal{K}$  is contained in  $\Omega(\rho(t_0))$ .

In Theorem 11.30 in [21] it was shown that for every  $\delta > 0$  there is  $R(\delta) < \infty$  such that for any any Ricci flow with singularity at time  $t_0$  and any  $\epsilon$ -horn  $\mathcal{K}$  in  $\Omega$  for this Ricci flow, all points of the  $\mathcal{K}$  with scalar curvature at least  $R(\delta)$  are centers of  $\delta$ -necks. Since  $\Omega$  is defined geometrically, it is an  $H$ -invariant subset of  $M$ , so that there is an induced isometric action  $H \times \Omega \rightarrow \Omega$ . For any  $\epsilon$ -horn  $\mathcal{K}$  let  $H_{\mathcal{K}}$  be the stabilizer in  $H$  of the end of  $\mathcal{K}$ .

**Claim 13.23.** *For any  $\delta > 0$  there is a sequence of  $H_{\mathcal{K}}$ -invariant  $\delta$ -necks  $N_n \subset \mathcal{K}$  tending to infinity in  $\mathcal{K}$ . The pull back from each  $N_n$  of action of  $H_{\mathcal{K}}$  is an action on  $S^2 \times (-\delta^{-1}, \delta^{-1})$  that is the product of a linear action on  $S^2$  with the trivial action on the interval.*

*Proof.* Let  $x_n \in \mathcal{K}$  be any sequence of points converging to the end of  $\mathcal{K}$ . Then, after passing to a subsequence, the based actions  $(\mathcal{K}, x_n, H_{\mathcal{K}})$ , with a sequence of rescaled metrics  $g_n$  with the property that in the metric  $g_n$  we have  $R(x_n) = 1$  converge geometrically to an action of  $H_{\mathcal{K}}$  on  $S^2 \times \mathbb{R}$  preserving the ends. (Notice that  $g_n = R(x_n)g$  and  $R(x_n) \rightarrow \infty$  as  $n \rightarrow \infty$ .) From [21] we know that for any such sequence there is a geometric limit that is  $S^2 \times \mathbb{R}$ , with the metric being the product of a round metric on  $S^2$  with the usual metric on  $\mathbb{R}$ . For each  $n$  let  $\Sigma_n$  be the central 2-sphere of an  $\delta$ -neck with  $x_n \in \Sigma_n$ . Since  $H_{\mathcal{K}}$  is compact and preserves the end of  $\mathcal{K}$ , for each element of this group we have  $h\Sigma_n \cap \Sigma_n \neq \emptyset$ , so that in the metric  $g_n$  every element of  $H_{\mathcal{K}}$  moves  $x_n$  a distance at most twice the diameter (in  $g_n$ ) of  $\Sigma_n$ , which is bounded by  $5\pi$ .

Let  $\mathcal{L}$  be the line field on  $\mathcal{K}$  orthogonal to the 2-plane field of maximal curvature directions. The group  $H_{\mathcal{K}}$  acts so as to preserve this line field and an orientation of it. This line field crosses  $\Sigma_n$  transversely with each line meeting  $\Sigma_n$  exactly once. This means that the leaf space of this line field is identified with  $\Sigma_n$ . The action of  $S^1$  on the line field then has two fixed points, meaning that there are two lines stabilized by the circle action, and hence point-wise fixed by the circle action. Thus, the circle has two fixed point on  $\Sigma_n$  both of which are within  $4\pi$  of  $x_n$  when distances are measured using  $g_n$ . From this it follows that the actions are uniformly continuous in the sense of Proposition 13.19. Thus, by that proposition after passing to a subsequence there is a limiting action of  $H_{\mathcal{K}}$  on the geometric limit of a subsequence, which as we have already said is  $S^2 \times \mathbb{R}$ . Consequently, the limiting action is the product of a linear action on  $S^2$  with the trivial action on  $\mathbb{R}$ . Proposition 13.19 also implies that given  $\epsilon > 0$  for all  $n$  sufficiently large, there is a point  $y_n$  whose distance from  $x_n$  (in  $g_n$ ) goes to zero as  $n \rightarrow \infty$  which is at the center of an  $H_{\mathcal{K}}$  invariant  $\epsilon$ -neck. Since the  $x_n$  converge to the end of  $\mathcal{K}$ , so do the  $y_n$ .  $\square$

At time  $t_0$  we can choose a  $H$ -invariant family of  $\epsilon$ -horns that make up all the ends of all connected components of  $\Omega$  that meet  $\Omega(\rho(t_0))$ . For each horn  $\mathcal{K}$  we construct an  $H_{\mathcal{K}}$ -invariant  $\bar{\delta}(t_0)$ -neck arbitrarily far out in the horn. Clearly, we can do this in such a way that the entire family of  $\delta(t_0)$ -necks is  $H$ -invariant. Once we have the  $H$ -invariant family of  $\bar{\delta}(t_0)$ -necks, we cut off each horn at the central 2-sphere of the neck in that horn. This allows us to cut off  $M$  at a  $H$ -invariant family of 2-spheres, where the stabilizer of each 2-sphere is isomorphic to a subgroup of  $O(3)$  and the action is  $\bar{\delta}(t_0)$ -close to the orthogonal action, meaning that there is an  $H$ -equivariant, almost isometric diffeomorphism from this collection of 2-spheres to a linear  $H$ -action on a family of 2-spheres.

The next step in the surgery process is to glue in the restriction to a 3-ball of the initial metric of the standard solution, where the gluing matches (up to an overall translation and reversal of sign) the distance function from the central point

of the initial metric of the standard solution with the interval coordinate in the  $H_K$ -invariant  $\bar{\delta}(t_0)$ -neck structure. The initial metric of the standard solution is  $O(3)$ -invariant. The gluing is done using a partition of unity which is chosen to be  $O(3)$ -invariant when written in the coordinates of the standard solution and hence depends only on the interval factor in the  $\bar{\delta}(t_0)$ -neck. This means that the  $H$ -action on the truncated version of  $M$  can be extended to an isometric  $H$ -action that is equivariantly diffeomorphic to a linear action on the family of balls we add in performing surgery. That is to say, the surgery procedure can be done in a  $H$ -equivariant fashion. Repeating this operation at each surgery time produces a  $H$ -equivariant Ricci flow with surgery defined for all time. This completes the proof of the proposition.  $\square$

### 13.3.2 Step 2: Examination of the components that disappear at finite time

In order to describe these components we introduce the following notion:

**Definition 13.24.** An  $\epsilon$ -tube  $\hat{T} \subset M$  is an open submanifold diffeomorphic to  $S^2 \times (0, 1)$  such that:

1.  $\hat{T}$  is a union of  $\epsilon$ -necks in  $M$ .
2. There is a disjoint union of two  $\epsilon$ -necks  $N_+ \amalg N_-$  contained in  $\hat{T}$  whose complement is compact. We denote by  $T \subset \hat{T}$  the open submanifold, also diffeomorphic to  $S^2 \times (0, 1)$ , between the central 2-spheres of  $N_+$  and  $N_-$ .

Given an  $\epsilon$ -tube  $\hat{T} \supset T$  we denote by  $U$  the union of  $T$  with the central thirds of  $N_+$  and  $N_-$ . Then  $T \subset U \subset \hat{T}$  and  $U$  is diffeomorphic to  $S^2 \times (0, 1)$ .

Suppose that  $t_0$  is a surgery time. Then for  $t < t_0$  but sufficiently close to it, the time-slices  $M_t$  form an ordinary Ricci flow, so that all these manifolds are identified and the flow is a flow of metrics  $g(t)$  on a fixed manifold, which we denote by  $M_{t_0}^-$ . As  $t \rightarrow t_0^-$  the metrics  $g(t)$  become singular at a compact subset  $X_{t_0} \subset M_{t_0}^-$ . Surgery at time  $t_0$  involves three operations. First, we cut  $M_{t_0}^-$  open along a finite  $H$ -invariant family of 2-spheres contained in  $M_{t_0}^- \setminus X_{t_0}$ . Denote by  $M'_{t_0}$  the result. It naturally contains  $X_{t_0}$ . Second, we remove an  $H$ -invariant family of components  $Y_{t_0}$  of  $M'_{t_0}$  with the property that  $X_{t_0} \subset Y_{t_0}$ . Third, we attach in an  $H$ -invariant way a family of 3-balls along the entire boundary of  $M_{t_0}^- \setminus Y_{t_0}$ .

We identify  $Y_{t_0}$  with a subset of  $M_{t_0}^-$  in the natural way. Then, with respect to any of the metrics  $g(t)$  for  $t < t_0$  sufficiently close to  $t_0$ , each point of  $Y_{t_0}$  has a canonical neighborhood. Thus, there are components of  $Y_{t_0}$  that are components of  $M_{t_0}^-$  and have positive curvature. All other components of  $Y_{t_0}$  are covered by  $\epsilon$ -necks,  $\epsilon$ -caps, and twisted  $\epsilon$ -caps. For components of  $Y_{t_0}$  that are components of  $M_{t_0}^-$  covered by these neighborhoods the possibilities are: (2a) those diffeomorphic to  $S^3$  and covered by two  $\epsilon$ -caps, possibly together with an  $\epsilon$ -tube, (2b) those diffeomorphic to  $\mathbb{R}P^3$  covered by an  $\epsilon$ -cap and a twisted  $\epsilon$ -cap possibly together with an  $\epsilon$ -tube, (2c) those diffeomorphic to an  $S^2$ -bundle over  $S^1$  covered by a union of  $\epsilon$ -necks, and (2d) those diffeomorphic to  $\mathbb{R}P^3 \# \mathbb{R}P^3$  covered by the union of two

twisted  $\epsilon$ -caps, possibly together with an  $\epsilon$ -tube. The possibilities for the topology of components of  $Y_{t_0}$  that are properly contained in components of  $M_{t_0}^-$  are the following: (3a) those diffeomorphic to  $S^2 \times I$  and contained in an  $\epsilon$ -tube, (3b) those diffeomorphic to  $B^3$  and contained in an  $\epsilon$ -cap possibly together with an  $\epsilon$ -tube, and (3c) those diffeomorphic to a twisted  $I$ -bundle over  $\mathbb{RP}^2$  and contained in twisted  $\epsilon$ -cap, possibly together with an  $\epsilon$ -tube.

**Proposition 13.25.** *Let  $t_0$  be a surgery time. and let  $C$  be a connected component of  $M_t$  for  $t < t_0$ , sufficiently close to  $t_0$ . Let  $H_C$  be the stabilizer in  $H$  of  $C$  and let  $\tilde{H}_C$  be the group of isometries of the universal covering  $\tilde{C}$  that cover elements of  $H_C$ . If  $C$  completely disappears at a surgery time  $t_0$  (i.e. if  $C$  is a component of  $Y_{t_0}$ ), then  $\tilde{C}$  has a homogeneous metric that is invariant under  $\tilde{H}_C$ .*

*Proof.* There are three possibilities.

**Case 1:**  $(C, g(t))$  has positive curvature for all  $t < t_0$  sufficiently close to  $t_0$ . Actually, there are two possibilities here: The first possibility is that  $(C, g(t))$  has positive sectional curvature and the diameter  $d(t)$  of this component converges to zero as  $t \rightarrow t_0^-$ . In this case rescaling the metrics  $(G, g(t))$  so that the diameter of the manifolds remains constant the metrics converge to a round metric. This limiting metric is invariant under the action of the stabilize  $H_C$  of  $C$  in  $H$  so that  $C$  is finitely covered by  $S^3$  with the round metric and  $H$  has a finite extension  $\tilde{H}$  by the fundamental group of  $C$  which acts on  $S^3$  via an embedding  $\tilde{H} \subset O(4)$ .

The second possibility is that  $C$  is a component of  $M_{t_0}^-$  and for all  $t < t_0$  sufficiently close to  $t_0$  the component  $C$  has positive sectional curvature in the metric  $g(t)$  but it is not converging to a point at time  $t_0$ . In this case, the Ricci flow applied to  $(C, g(t))$  exists for some finite time (longer than  $t_0 - t$ ) and at the limiting time the metric on  $C$  becomes round. Since the Ricci flow is equivariant under the stabilizer  $H_C$ , we see that  $C$  admits an  $H_C$ -invariant round metric. Thus, in Case 1 the component  $C$  has a round metric that is invariant under the stabilizer  $H_C$  of  $C$  in  $H$ .

**Case 2a:**  $C$  is the union of two  $\epsilon$ -caps and possibly an  $\epsilon$ -tube. Notice that  $H_C$  is a subgroup of finite index in  $H$  so that the number of its components is at most  $N$ . If  $C$  is a union of two  $\epsilon$ -caps, then  $C$  has positive curvature and is already covered by Case 1. Thus, we can assume that the cores of the two caps  $\mathcal{C}_0$  and  $\mathcal{C}'_0$  are disjoint. Thus,  $C$  is the union of the cores of these two caps and an  $\epsilon$ -tube  $T$  with the property that the boundaries of  $\mathcal{C}_0$  and  $\mathcal{C}'_0$  are cross sections for the line field  $\mathcal{L}_T$ . There is a point  $x$  in  $\mathcal{C}_0$  that is not contained in any  $\epsilon$ -neck. For any  $h \in H_C$  the image  $hx$  is contained in either  $\mathcal{C}_0$  or  $\mathcal{C}'_0$ . Thus, by Lemma 13.20 there is a subgroup  $H'_C$  of index at most two in  $H_C$  such that for every  $h \in H'_C$  the intersection of  $\mathcal{C}_0$  with its image under  $h$  is non-empty. In this case, there is an  $H'_C$ -invariant submanifold  $X_1 \subset C$  with boundary 2-sphere contained in  $T$  and a cross section for  $\mathcal{L}_T$ . According to Proposition 13.21 the action of  $H'_C$  on  $X_1$  is equivariantly diffeomorphic to a linear action on the 3-ball. We perform the analogous construction for  $X_2 \subset \mathcal{C}'_0$ , and do it  $H_C$ -equivariantly if  $H_{C'} \neq H_C$ . Then the  $H_C$  action on  $X_1 \amalg X_2$  is equivariantly diffeomorphic to a linear action. The region  $R$  between  $\partial X_1$  and  $\partial X_2$  is an  $H_C$ -invariant subset of  $T$  with boundary

transverse to the line field  $\mathcal{L}_T$ , which is  $H_C$ -invariant. Using this line field and the fact that the action on the boundary 2-spheres is linear, we see that there is an  $H_C$ -equivariant diffeomorphism from  $R$  to a linear action on  $S^2 \times I$ . It follows that there is an embedding  $H_C \subset O(3) \times O(1) \subset O(4)$  and an  $H_C$ -equivariant diffeomorphism from  $C$  to the induced linear action of  $H_C$  on  $S^3$ .

**Case 2b:  $C$  is the union of a twisted  $\epsilon$ -cap and an  $\epsilon$ -cap possibly together with an  $\epsilon$ -tube.** In this case there is a double cover  $\tilde{C}$  of  $C$  with an action of an extension  $\tilde{H}_C$  of  $H_C$  by a group of order 2 acting on  $\tilde{C}$  covering the given action of  $H_C$  on  $C$ . Thus, this case follows immediately from the previous.

**Case 2c:  $C$  is an  $S^2$ -bundle over  $S^1$  and, for every  $t < t_0$  sufficiently close to  $t_0$ , every point of  $C$  is the center of an  $\epsilon$ -neck.** Pass to the universal covering  $\tilde{C}$ , and let  $\tilde{H}$  be the group of isometries of  $\tilde{C}$  that normalize the group of covering transformations of  $\tilde{C} \rightarrow C$  and project to elements of  $H$ . Then there is an exact sequence:

$$\{1\} \rightarrow \mathbb{Z} \rightarrow \tilde{H} \rightarrow H \rightarrow \{1\}.$$

One possibility is that there is a circle subgroup of  $H$  whose orbits represent non-trivial elements in  $H_1(C)$ . Since the fundamental group of a compact, connected semi-simple group is finite, in this case it follows that the component of the identity  $H^0$  of  $H$  contains a central circle whose orbits represent non-trivial elements in  $H_1(C)$ . The quotient of the center of  $H^0$  by this group then acts effectively on the quotient  $C/S^1$ . This implies that the center of  $H^0$  has rank either one or two. The center of the identity component of the covering group  $\tilde{H}^0 \subset \tilde{H}$  is then either isomorphic to  $\mathbb{R}$  or  $S^1 \times \mathbb{R}$ , and the  $\mathbb{R}$  acts freely and properly discontinuously on  $\tilde{C}$  with quotient a 2-sphere.

**Claim 13.26.** *We can choose the  $\mathbb{R} \subset \tilde{H}^0$  to be a normal subgroup of  $\tilde{H}$ .*

*Proof.* If the center of  $\tilde{H}^0$  is  $\mathbb{R}$ , then this subgroup is a normal subgroup of  $\tilde{H}$ . If the center of  $\tilde{H}^0$  is isomorphic to  $S^1 \times \mathbb{R}$ , then  $\tilde{H}$  acts on this group through a finite image. It is easy to see that any finite subgroup of automorphisms of  $S^1 \times \mathbb{R}$  has an invariant  $\mathbb{R}$ -factor.  $\square$

Fix a normal subgroup  $\mathbb{R} \subset \tilde{H}$ . This group acts freely and properly discontinuously on  $\tilde{C}$  with quotient  $S^2$ . There is a cross section and hence there is a product structure  $\tilde{C} = S^2 \times \mathbb{R}$  so that  $\mathbb{R}$  actions by translation in the second factor. Since  $\mathbb{R}$  is a normal subgroup of  $\tilde{H}$ , the action of  $\tilde{H}$  preserves the foliation of  $\tilde{C}$  by the copies of  $\mathbb{R}$ . Let  $\bar{H} = \tilde{H}/\mathbb{R}$ . It is a compact group. Consequently,  $H^2(\bar{H}; \mathbb{R})$  is trivial and hence there is a splitting  $\tilde{H} = \mathbb{R} \rtimes \bar{H}$ . Let  $\Sigma = S^2 \times \{0\}$  and consider the intersection of  $\bar{H} \cdot \Sigma \rightarrow S^2 \times \mathbb{R}$  with  $\{x\} \times \mathbb{R}$ . This gives a function  $\psi_x: \bar{H} \rightarrow \mathbb{R}$  that varies smoothly with  $x \in S^2$ . We form the average of this function using Haar measure of volume 1 on  $\bar{H}$ . The result is a function  $\bar{\psi}: S^2 \rightarrow \mathbb{R}$  which is  $\bar{H}$  invariant in the sense that  $\bar{\psi}(hx) = \bar{\psi}(x)$  for every  $h \in \bar{H}$ . This means that the graph of  $\bar{\psi}$ , denoted  $\Sigma'$ , is a 2-sphere transverse to the  $\mathbb{R}$ -foliation that is invariant under  $\bar{H}$ . We define a product structure on  $\tilde{C}$  so that  $\Sigma' \times \mathbb{R}$  so that  $\Sigma'$  is the 2-sphere cross section at 0 and  $\mathbb{R}$  acts by translations. This product structure is invariant under  $\tilde{H}$ . Since the action of  $\bar{H}$  on  $\Sigma'$  is equivariantly diffeomorphic to a linear action, it follows there



is an  $\tilde{H}$ -equivariant diffeomorphism from  $\tilde{C}$  to an  $\tilde{H}$ -action on  $S^2 \times \mathbb{R}$  that is the product of a linear action on  $S^2$  and a linear action on  $\mathbb{R}$ .

Now we consider the case when there is no  $S^1 \subset H$  whose orbits represent non-trivial elements in  $H_1(C)$ . In this case we use the line field  $\mathcal{L}_C$  that is orthogonal to the 2-planes of maximal curvature. We suppose that we have chosen  $\epsilon > 0$  sufficiently small so that  $\epsilon^{-1}$  is much larger than the order  $N$  of the group of connected components of  $H$ . The line field  $\mathcal{L}$  integrates to give a foliation of the universal covering  $\tilde{C}$  by properly embedded lines with quotient space  $S^2$ . The group  $\tilde{H}$  acts on  $\tilde{C}$  preserving this foliation and hence there is an induced action of  $\tilde{H}$  on  $S^2$ . It follows that every element in the connected component of the identity of  $\tilde{H}$  acts on the quotient space with fixed points, that is to say, it stabilizes one of the flow lines in  $\tilde{C}$  of the line field  $\mathcal{L}$ . That element then fixes the flow line point-wise, and hence has fixed points on any 2-sphere cross section.

Fix an  $\epsilon$ -neck  $N_\epsilon$  in  $C$  with central 2-sphere  $S^2$ . Let  $H^0 \subset H$  be the subgroup of index at most 2 consisting of elements preserving the direction of the line field  $\mathcal{L}$ . We claim that every element  $h \in H^0$  either has the property that  $h \cdot S^2 \cap S^2 \neq \emptyset$  or  $hN_{\epsilon/3N} \cap N_{\epsilon/3N} = \emptyset$ . The reason is that if  $h \cdot S^2 \cap S^2 = \emptyset$  and yet  $hN_{\epsilon/3N} \cap N_{\epsilon/3N} \neq \emptyset$ , then for every  $1 \leq k \leq N$  the  $k^{\text{th}}$  power of  $h$  moves  $S^2$  so that it is contained in  $N_\epsilon$  but does not meet  $S^2$ . Since  $N$  is the order of the component group of  $H$ , it follows that for some  $1 \leq k \leq N$  the  $k^{\text{th}}$  power of  $h$  is in the component of the identity and hence, by the discussion above, fixes some point of  $S^2$ . This is a contradiction.

Once we have this dichotomy, it follows easily (provided that  $\epsilon^{-1}/N \geq 3$ ), that the subset of elements  $h \in H^0$  with the property that  $h \cdot S^2 \cap S^2 \neq \emptyset$  is a normal subgroup  $H'$  of  $H^0$  with finite cyclic quotient. Fix a product structure on an open subset of  $N$  that contains the middle half of  $N$  by integrating the line field  $\mathcal{L}_N$  from  $S^2$ . Then for each  $h \in H'$  the image  $h \cdot S^2$  is a cross section of the product structure and is the graph of a function from  $S^2 \rightarrow \mathbb{R}$ . Averaging these functions over  $H'$  gives a function whose graph is an  $H'$  invariant cross-section  $\Sigma$  contained in the middle third of  $N$ . The translates of  $\Sigma$  under  $H^0$  are a finite disjoint union of 2-spheres and the region between any successive ones is diffeomorphic to  $S^2 \times I$  by a diffeomorphism that sends  $\mathcal{L}_C$  to the tangent line field to the interval factors. The universal covering of  $C$  is obtained by gluing these product regions end-to-end infinitely in both directions. Let  $h_1: \Sigma \rightarrow \Sigma$  be the gluing map. To see that the  $H^0$ -action is equivalent to a linear action we need the following claim.

**Claim 13.27.** *Suppose that  $H \times S^2 \rightarrow S^2$  is a compact group action and that  $\psi: S^2 \rightarrow S^2$  is a diffeomorphism with the property that there is an automorphism  $\varphi: H \rightarrow H$  with  $\psi(hx) = \varphi(h)\psi(x)$  for all  $h \in H$  and all  $x \in S^2$ . Then there is a one-parameter family of diffeomorphisms  $\psi_t: S^2 \rightarrow S^2$  with  $\psi_0 = \psi$  and with  $\psi_t(hx) = \varphi(h)\psi_t(x)$  for all  $h \in H$ , all  $x \in S^2$ , and all  $0 \leq t \leq 1$ , such that there is a round metric on  $S^2$  invariant under  $H$  and  $\psi_1$ .*

*Proof.* Without loss of generality we can suppose that  $H$  acts effectively. Let's consider the case when  $H$  is a finite group. In this case  $Q = S^2/H$  is a 2-dimensional orbifold and  $\psi$  induces an orbifold isomorphism  $\bar{\psi}: Q \rightarrow Q$ . This orbifold isomorphism is isotopic through orbifold isomorphisms to one of finite order (the order of

the permutation of the exceptional points induced by  $\bar{\psi}$ ). This deformation lifts to a deformation of  $\psi$  as required with  $\psi_1$  of finite order modulo  $H$ . Thus, the group generated by  $H$  and  $\psi_1$  is finite, and the result follows.

Now suppose that  $H$  is of dimension one. Then the quotient space  $S^2/H$  is an interval and the result is elementary in this case.

Lastly, if  $H$  has dimension greater than 1, then it is  $SO(3)$  acting in the standard way on  $S^2$  and the map  $\psi$  is determined by  $\varphi$  and either is contained in  $SO(3)$  or together with  $H$  generates  $O(3)$ .  $\square$

We apply this to  $\tilde{C}$  which is a union  $\{S^2 \times I\}_{n=-\infty}^{\infty}$  with  $(S^2 \times \{1\})_i$  glued to  $(S^2 \times \{0\})_{i+1}$  by the map  $\psi_1$ . The subgroup  $H^0$  acts linearly on  $(S^2 \times I)_0$ . According to the previous claim we can deform the product structure on  $(S^2 \times I)_0$  in an  $H^0$ -invariant fashion and so that the group generated by  $H^0$  and  $\psi_1$  preserves a round metric on  $S^2$ . This shows that the action of  $H^0$  on  $S^2 \times \mathbb{R}^1$  is equivalent to a product of linear actions. This proves the result when  $H = H^0$ .

It remains to consider the case when  $H^0 \subset H$  is of index 2. In this case the subgroup  $H'$  of elements  $h \in \tilde{H}$  that preserve the direction of the line field and also fix a point of every  $S^2$  cross section form a normal subgroup with quotient an infinite dihedral group. Fix an element  $\tau \in \tilde{H}$  reversing direction of the line field  $\mathcal{L}_{\tilde{C}}$ . Then  $H'$  and  $\tau$  generate an extension of  $\mathbb{Z}/2\mathbb{Z}$  by  $H'$ . Averaging cross sections as before, we obtain a cross section  $\Sigma \subset \tilde{C}$  to the line field that is invariant under  $H'$  and  $\tau$ . The translates of  $\Sigma$  under  $\tilde{H}$  form a disjoint family of 2-spheres, glued end-to-end, by a diffeomorphism  $h_1$  to form  $\tilde{C}$ . Invoking the previous claim again we see that by deforming the product structure on the region bounded by  $\Sigma$  and one of its nearest translates, we can arrange that there is a round metric on  $\Sigma$  that is preserved by  $H', \tau$  and by the gluing diffeomorphism  $h_1$ . This produces a product structure  $\tilde{C} \cong S^2 \times \mathbb{R}$  with the property that  $\tilde{H}$  is the product of a linear action on  $S^2$  and a linear action on  $\mathbb{R}$ .

**Case 2d:  $C$  contains two disjoint quotients of an  $\epsilon$ -neck by an involution flipping their ends and these quotients are connected by an  $\epsilon$ -tube.** This case follows from the previous by passing to the two-sheeted covering.

This completes the proof of the proposition.  $\square$

Now let us examine the components of  $Y_{t_0}$  that are not components of  $M_{t_0}^-$ . We have an  $H$ -invariant family of  $\epsilon$ -necks  $N_j$ . The central 2-sphere of each neck separates  $\Omega(\rho(t_0))$  from the end on the  $\epsilon$ -horn containing it. We do surgery on the central 2-spheres on the neck (which form a disjoint union of  $H$ -invariant submanifolds on which the action is linear) and add 3-balls to these 2-spheres and extend the action to a disjoint union of linear actions of  $H$  on a disjoint union of 3-balls. If we consider  $t < t_0$  but  $t$  sufficiently close to  $t_0$ , there are is a disjoint union of  $\epsilon$ -tubes and  $\epsilon$ -tubes with either  $\epsilon$ -caps or twisted  $\epsilon$ -caps attached at one end that contains the disjoint union of the  $N_j$ . In fact, it is easy to arrange that the family of  $N_j$  has exactly two members in each of the  $\epsilon$ -tubes in this collection, one near each end, and each capped  $\epsilon$ -tube contains exactly one of the  $N_j$ , near its non-capped end. For any

such component  $T$ , we denote by  $H_T$  the stabilizer of the submanifold of  $T$  with boundary the central 2-spheres of all the  $N_j$  contained in  $T$ .

Let  $T$  be an  $\epsilon$ -tube in this collection with  $N_1$  and  $N_2$  being the  $\epsilon$ -necks near its ends. Either  $H_T$  is equal to the stabilizer of each  $N_1$  and  $N_2$ , or it contains these stabilizers as a subgroup of order two and  $H_T$  contains an element interchanging  $N_1$  and  $N_2$ . According to the surgery prescription, the action of the stabilizer of each  $N_j$  on that component equivariantly diffeomorphic to the product of a linear action on  $S^2$  and the trivial action on the interval. Using the flow lines of the line field on an  $\epsilon$ -tube, we see that the action on the region between the central 2-sphere near the ends of the tube is also equivariantly diffeomorphic to a product of linear actions on  $S^2$  and on the interval (possibly containing a flip interchanging the two ends).

In the case that  $T$  is an the  $\epsilon$ -tube capped with a twisted  $\epsilon$ -cap,  $H_T$  is equal to the stabilizer of the  $\epsilon$ -neck,  $N_j$ , that it contains. Passing to the double covering reduces this case to the previous one.

Lastly, if the component  $T$  is an  $\epsilon$ -tube capped by an  $\epsilon$ -cap, then  $H_T$  is equal to the stabilizer of the neck,  $N_j$ , that it contains. We must show that the action on the 3-ball cut off by the central 2-sphere of the  $N_j$  contained in  $T$  is linear. We know that near the boundary the action is a linear action on  $S^2$  times the trivial action on  $I$ . Since  $H_T$  is a subgroup of finite index in  $H$  according to Proposition 13.21 provided that we have chosen  $\epsilon > 0$  sufficiently small given the number of connected components of  $H$ , that the  $\epsilon$ -cap contains an  $H_T$ -invariant 3-ball  $\bar{C}$  on which the action is linear. Furthermore, the region between the central 2-sphere of  $N_j$  and  $\partial\bar{C}$  is a product region with the product structure being given by an  $H_T$ -invariant line field. Thus, the action on this region is a product of a linear action on  $S^2$  with the trivial action on the interval.

We have established the following:

**Proposition 13.28.** *Let  $t_0$  be a surgery time for a Ricci flow with surgery of compact 3-manifolds and let  $H$  be a compact group acting on this Ricci flow with surgery. Then each component of the  $H$ -invariant region  $Y_{t_0}$  of  $M_{t_0}^-$  that is removed by doing  $H$ -equivariant surgery at time  $t_0$  is one of the following types:*

1. *a component  $C$  of  $M_{t_0}^-$  with stabilizer  $H_C$  acting in such a way that it is covered by an isometric action on a manifold with a homogeneous metric modelled on  $Solv$ ,  $Nil$ ,  $\mathbb{R}^3$ ,  $S^3$ , or  $S^2 \times \mathbb{R}$ ,*
2. *diffeomorphic to  $S^2 \times I$  and the action of its stabilizer in  $H$  is equivariantly diffeomorphic to the product of a linear action on  $S^2$  and a linear action on  $I$ ,*
3. *diffeomorphic to a 3-ball and the action of its stabilizer is equivariantly diffeomorphic to a linear action, or*
4. *diffeomorphic to the complement of an open ball in  $\mathbb{R}P^3$  and the action of its stabilizer is equivariantly diffeomorphic to one one that lifts to an action on the double cover which is the product of a linear action on  $S^2$  and a linear action on  $I$ .*

Surgery of the third type produces a manifold after surgery equivariantly diffeomorphic to the manifold before surgery. Surgery of the second type is inverse to  $H$ -equivariant connected sum decomposition. Surgery of the fourth type does a connected sum decomposition and removes prime factors each diffeomorphic to  $\mathbb{R}P^3$  and each with the action of its stabilizer being covered by a linear action on  $S^3$ .

As a result, to prove the equivariant version of the Geometrization Conjecture for  $M = M_0$ , it suffices to prove it for  $M_t$  for any  $t < \infty$  sufficiently large.

### 13.3.3 Step 3: Proof of the Generalized Smith Conjecture

At this point we can give a proof of the Generalized Smith Conjecture, which says that any action of a compact group on  $S^3$  is equivariantly diffeomorphic to a linear action, using Ricci flow. (The usual Smith Conjecture is the case of orientation-preserving actions of prime order cyclic groups.)

Suppose that  $M$  is a simply connected 3-manifold and  $H \times M \rightarrow M$  is a compact group action. We choose an  $H$ -invariant metric on  $M$  and run the Ricci flow with surgery, producing a one-parameter family  $(M_t, g(t))$  of Riemannian manifolds with  $H$ -actions. Consider a surgery time  $t_0$  and the disjoint union of components of  $C$  of  $M_{t_0}^-$  that disappear at time  $t_0$ . This is an  $H$ -invariant subset of  $M_t$  for every  $t < t_0$ , sufficiently close to  $t_0$ . Since  $M$  is simply connected, each component of  $C$  is diffeomorphic to a  $S^3$ -sphere.

According to Proposition 13.28 the action of the stabilizer of  $C$ ,  $H_C \subset H$ , on  $C$  is equivariantly diffeomorphic to a linear action. It follows immediately that the action of  $H$  on  $C$  is equivariantly diffeomorphic to a disjoint union of linear actions of  $H$  on families of 3-spheres.

Surgery along the  $H$ -invariant family of  $\epsilon$ -tube components of  $C$  is done in an equivariant fashion and hence is an  $H$ -equivariant connected sum decomposition.

Surgery that removes an  $H$ -invariant family of  $\epsilon$ -tubes, each with a  $\epsilon$ -cap attached to the end removes a linear action on disjoint union of 3-balls and replaces it by another action with the same boundary, one that is also equivariantly diffeomorphic to a linear action. Hence, this operation does not change the equivariant diffeomorphism type. Thus, according to Proposition 13.28, up to equivariant diffeomorphism the effect of surgery in this Ricci flow with surgery is to remove a disjoint union of  $S^3$  with linear actions and to do equivariant connected sum decomposition.

Of course, according to one of the main results of [21], since  $M$  has trivial fundamental group Ricci flow with surgery applied to  $M$  completely disappears at some finite-time. At the final time  $T$ , we see that the manifold that is disappearing is a disjoint union of linear actions of  $H$  on a disjoint union of 3-spheres. As we move backwards in time across singularities we either (i) make no change up to equivariant diffeomorphism, (ii) add new disjoint unions of linear actions of  $H$  on families of 3-spheres, or (iii) do one or more  $H$ -equivariant connected sums. Hence, by Proposition 13.9 we show by induction moving backward in time across the finite number of singular times that at each time the  $H$ -action is equivariantly diffeomorphic to a linear action of  $H$  on a disjoint union of families of 3-spheres. In particular, this is true at the initial time, showing that the action of  $H$  on  $M$  is equivariantly diffeo-

morphic to a linear action of  $H$  on  $S^3$ . Notice that in this argument we needed no results outside of those proved directly from the existence of equivariant Ricci flow with surgery.

### 13.4 Proof of Theorem 13.4.

Now we consider the general case of a Ricci flow with surgery with an isometric action of a compact group  $H$ . By our analysis of the surgeries we see that for any  $t > 0$  the action  $H \times M_0 \rightarrow M_0$  is obtained from the action  $H \times M_t \rightarrow M_t$  by a sequence of operations of the following type: (i) addition of copies of  $S^3$ ,  $S^3/\Gamma$ ,  $S^2$ -bundles over  $S^1$ ,  $\mathbb{R}P^3$ , and  $\mathbb{R}P^3 \# \mathbb{R}P^3$  each with actions preserving locally homogeneous metrics; (ii) equivariant connected sum decomposition; and (iii) equivariant diffeomorphism. In particular, if the  $H$ -equivariant Ricci flow with surgery with  $(M, g(0))$  as initial conditions becomes extinct after finite time (which is automatic if  $\pi_1(M)$  is a free product of cyclic groups and finite groups), then  $M$  is an equivariantly diffeomorphic to an equivariant connected sum of isometric actions on round manifolds and manifolds with metrics locally modelled on  $S^2 \times \mathbb{R}$ . Furthermore, if we can show that for some  $t$  sufficiently large, the action of  $H$  on  $M_t$  satisfies the conclusions of Theorem 13.4 then the same is true for the action of  $H$  on  $M_0$ . For  $t$  sufficiently large the only components of the manifold  $M_t$  that are not aspherical are 3-spheres. Since we have already established the result for actions on disjoint union of 3-spheres, it suffices to prove the result for the disjoint union of the connected components of  $M_t$  that are not homeomorphic to 3-spheres. **This allows us to assume that every component of  $M_t$  is aspherical. We implicitly make this assumption from now on.**

Next, we study the decomposition as  $t \rightarrow \infty$  of the slices  $(M_t, g(t))$  of the  $H$ -invariant Ricci flow with surgery. There are finitely many hyperbolic manifolds  $H_1, \dots, H_k$  (with metrics of constant sectional curvature  $-1/2$ ) that appear as limits as  $t \rightarrow \infty$  of the locally non-collapsed regions of  $(M_t, (1/t)g(t))$ . Let  $\mathcal{H}$  be the disjoint union of the  $H_i$ . According to Proposition 13.19 there is an induced isometric action of  $H$  on  $\mathcal{H}$ . Let  $\overline{\mathcal{H}}$  the truncation along horospherical tori of area  $3w/4$  (where  $w$  is a suitably small, positive constant). Then the  $H$ -action on  $\mathcal{H}$  leaves  $\overline{\mathcal{H}}$  invariant, so that there is an induced  $H$ -action on  $\overline{\mathcal{H}}$ . Actually, we enlarge the truncation to  $\overline{\mathcal{H}}_1$  by adding a collar neighborhood of between horospherical tori to each boundary component. We choose this collar to be of a fixed length  $\epsilon^{-1}$  in the hyperbolic metric of sectional curvature  $-1/2$ . According to Proposition 13.19 for all  $t$  sufficiently large there are embeddings  $\Phi_t: \overline{\mathcal{H}}_1 \rightarrow (M_t, (1/t)g(t))$  are  $H$ -equivariant and converge as  $t \rightarrow \infty$  to an isometric embedding. The image of  $\overline{\mathcal{H}}$  under this embedding is a region, denoted  $M_t(w, +)$ , of  $(M_t, g(t))$  that contains all points  $x \in M_t$  that are not  $w$ -volume collapsed on the scale of the negative curvature. Also, according to Proposition 2.25, the boundary of  $M_t(w, +)$  consists of tori each of which is incompressible in  $M_t$ .

We turn now to the complement  $M_t(w, -) = M_t \setminus \text{int}(M_t(w, +))$  (for  $t$  sufficiently large). It is an  $H$ -invariant compact 3-manifold with geodesically convex boundary. The manifold  $M_t(w, -)$  is  $w$ -volume collapsed on the scale of the negative curvature

and the boundary of  $M_t(w, -)$  has a collar neighborhood, which, with respect to the metric  $(1/t)g(t)$ , is  $H$ -equivariantly almost isometric to an  $H$ -action preserving the hyperbolic metric on a disjoint union of regions in cups of a complete (possibly disconnected) hyperbolic manifolds bounded by parallel horospherical tori a distance  $\epsilon^{-1}$  apart, one such region in each cusp.

Our goal is to show the following:

**Proposition 13.29.** *For all  $t$  sufficiently large there is an  $H$ -invariant family of incompressible tori  $\mathcal{T}(t)$  in  $M_t(w, -)$  such that cutting  $M_t(w, -)$  open along these tori produces a compact 3-manifold  $\mathcal{Y}(t)$  with an  $H$ -action and each component of  $\mathcal{Y}(t)$  is of one of the following types:*

1. *a component diffeomorphic to  $T^2 \times I$  in such a way that the action of its stabilizer is equivariantly diffeomorphic to a product of a linear action on  $T^2$  and a linear action on the interval.*
2. *a component diffeomorphic to a twisted  $I$ -bundle over the Klein bottle and the action of its stabilizer is double covered by an action on  $T^2 \times I$  as in the first item.*
3. *a closed component with a flat metric that is invariant under the action of its stabilizer.*
4. *a closed component that is fibered over  $S^1$  with  $T^2$  fiber and a locally homogeneous metric (or Solv, Nil, or Flat type) invariant under the action of its stabilizer.*
5. *a Seifert fibration with incompressible boundary whose total space is diffeomorphic to neither  $T^2 \times I$  nor to a twisted  $I$ -bundle over the Klein bottle.*

First let us deal with flat components. Recall that for every  $x \in M_t(w, -)$  we take  $\rho(x)$  to be such that setting  $g'(x) = \rho^{-2}(x)g(t)$  the infimum of the sectional curvatures of  $B_{g'(x)}(x, 1)$  is  $-1$ . Since the volume of this ball is at most  $w$ , this implies that provided  $w > 0$  is chosen sufficiently small, given  $\epsilon$ , for each  $x$  there is an Alexandrov space  $B(\bar{x}, 1)$  with curvature  $\geq -1$  and of dimension 0, 1 or 2 such that  $B_{g'(x)}(x, 1)$  is within  $\epsilon$  in the Gromov-Hausdorff distance from  $B(\bar{x}, 1)$ . Suppose that for some  $x$ , the ball has dimension zero. Then rescale the metric on the connected component  $C_t$  of  $M_t(w, -)$  containing  $x$  so that the diameter is 1. The rescaled metric is close in the Gromov-Hausdorff distance to an Alexandrov ball of curvature  $\geq -1$  and dimension either 1, 2 or 3. If there is a sequence of  $x_{t_n} \in M_{t_n}(w, -)$  for  $t_n \rightarrow \infty$  with the rescaled metrics on the connected component  $C_{t_n}$  of  $M_{t_n}(w, -)$  of diameter one being uniformly volume non-collapsed then these rescaled Riemannian manifolds converge to a compact flat manifold. According to Proposition 13.19, passing to a subsequence so that the stabilizers of the components in question are all isomorphic, this common group is also represented as a group of isometries of the limiting flat manifold. Furthermore, for all  $n$  sufficiently large, there are equivariant diffeomorphisms from the limit action to the action on  $C_{t_n}$ , showing that  $C_{t_n}$  has a flat metric invariant under its stabilizer. This allows us to

assume that the nearby Alexandrov balls are all of dimension 1 or 2. This implies that there is a further decomposition of  $M_t(w, -)$  into two types of pieces: those close to interior points of open intervals, and those (possibly after rescaling further) that are close to 2-dimensional Alexandrov space of curvature  $\geq -1$ .

Now let us fix  $t$  sufficiently large and let us consider the open set  $U_1(t)$  of  $M_t(w, -)$  as given in Section 12. Since  $X_1(t)$  and  $U_1(t)$  are defined geometrically, they are  $H$ -invariant subsets. Furthermore, as we have seen there is a line field on  $U_1(t)$  that makes a small angle with any geodesic ending at a point  $y \in U_1(t)$  provided that the geodesic has length at least  $10^{-3}$  in the metric  $g'(y)$ . It follows that we can average the line field over  $H$  and produce an  $H$ -invariant line field with the same property. Each component of  $U_1(t)$  fibers over a 1-manifold and hence is of one of the following types;

1. a component of  $M_t$  that fibers over the circle with fiber  $T^2$  or  $S^2$ ,
2. an open subset diffeomorphic to  $T^2 \times (0, 1)$  and the action of its stabilizer is equivalent to the product of a linear action on  $T^2$  and a linear action on  $(0, 1)$ ,
3. an open subset diffeomorphic to  $S^2 \times (0, 1)$  and the action of its stabilizer is equivalent to the product of a linear action on  $S^2$  with a linear action on the interval.

In fact, the hypothesis that every component of  $M_t$  is aspherical implies that there are no components that fiber over the circle with  $S^2$  as fiber.

The next step in Section 12 was to take a slightly smaller subset  $U'_1(t) \subset U_1(t)$  whose ends are of the form  $U(x_{\mathcal{E}})$  and whose external boundary consists of locally flat surfaces. Consider a component of a level surface for the distance function from  $x_{\mathcal{E}}$  approximately halfway from  $x_{\mathcal{E}}$  to the external frontier of  $U(x_{\mathcal{E}})$ . This is a cross section for the line field on  $U'_1(t)$ . We can approximate it arbitrarily closely by a smooth cross section  $\Sigma(x_{\mathcal{E}})$ . Since the diameter of each component  $U(x_{\mathcal{E}})$  is much larger than the distance from its external boundary to the frontier of  $U_1(t)$ , it follows that the stabilizer of the frontier of  $U_1(t)$  closest to  $U(x_{\mathcal{E}})$  maps  $\Sigma(x_{\mathcal{E}})$  to another cross section in  $U(x_{\mathcal{E}})$ . Thus, we can average these cross section over the stabilizer of this component of the frontier of  $U_1(t)$ , to produce a smooth cross section invariant under this stabilizer. This allows us to choose smooth cross sections  $\Sigma(x_{\mathcal{E}})$ , exactly one cross section in each neighborhood  $U(x_{\mathcal{E}})$  of an end of  $U'_1(t)$ . We do this in such a way that the entire collection is  $H$ -invariant. We denote this  $H$ -invariant collection of tori and 2-spheres by  $\mathcal{T}_0(t)$ .

Next, as in Section 12.1.1 we expand  $U'_1(t)$  to a larger open submanifold  $U''_1(t)$  by adding to  $U'_1(t)$  all of its complementary components  $V_0(x_i)$  as in Item 1 of Corollary 12.11. There is a subset of the ends of  $U'_1(t)$  that are ends of  $U''_1(t)$  and correspondingly a subset  $\mathcal{T}_1(t)$  of the surface components of  $\mathcal{T}_0(t)$  near to and parallel to these ends. Since  $U''_1(t)$  is geometrically defined, it is  $H$ -invariant and hence so is the collection  $\mathcal{T}_1(t)$  of 2-spheres and tori.

The subset of  $U''_1(t)$  external to the surfaces  $\mathcal{T}_1(t)$  is diffeomorphic to a product of  $\mathcal{T}_1(t)$  with an open interval. We denote by  $W'_1(t)$  the complementary compact submanifold of  $U''_1(t)$ . The boundary of  $W'_1(t)$  is  $\mathcal{T}_1(t)$ . According to Proposition 12.12

and the fact that  $M_t(w, -)$  is aspherical each connected component of  $W'_1(t)$  is of one of the following types:

1. a  $T^2$ -bundle over the circle or the union of two twisted  $I$ -bundles over the Klein bottle glued together along their boundary,
2.  $T^2 \times I$  or  $S^2 \times I$ , or
3. a twisted  $I$ -bundle over the Klein bottle, a solid torus, or the 3-ball.

Because of the Smith conjecture and the fact that any action on  $T^2 \times I$  is equivalent to the product on a linear action on  $T^2$  and a linear action on the interval we see that the actions of the stabilizers of components of the above type are equivalent to

1. in the second case the product of a linear action on the surface with a linear action on  $I$ ,
2. in the third case a linear action on the 3-ball, the product of a linear action on  $D^2$  with a linear action on  $S^1$  or an action double covered by the product of a linear action on  $T^2$  and a linear action on  $I$ .

The surfaces  $\mathcal{T}_1(t)$  do not agree with the surfaces  $\Sigma(\mathcal{E})$  which form the boundary of  $W_1$ . Nevertheless, each neighborhood of the ends,  $U(\mathcal{E})$ , of  $U'_1(t)$  contain one component of  $\mathcal{T}_1(t)$  which is a cross-section of the line field and hence isotopic in  $U(\mathcal{E})$  to the surface  $\Sigma(\mathcal{E})$ . Thus, while the geometry near the boundaries is different, the components of  $M_t(w, -) \setminus \text{int}W'_1(t)$  are diffeomorphic to the corresponding components of  $W_2$ . In particular, any component of the result of cutting  $M_t(w, -)$  open along  $\mathcal{T}_1(t)$  is either a component of  $W'_1(t)$  or is a union of the total space of a Seifert fibration possibly with solid tori and/or solid cylinders attached so as to kill the homotopy class of the generic fiber. The ends of the solid cylinders are contained in 2-sphere components of  $\mathcal{T}_1(t)$ , and each 2-sphere component of  $\mathcal{T}_1(t)$  contains two ends of solid cylinders. Consider a component  $X$  of  $W'_1(t)$  that is diffeomorphic to  $S^2 \times I$ . Since each component of  $M_t$  is aspherical, exactly one of the boundary components of  $X$  bounds a 3-ball  $B(X)$  that does not contain  $X$ . We remove from  $\mathcal{T}_1(t)$  both boundary components of  $X$  together with all the components of  $\mathcal{T}_1(t)$  contained in  $B(X)$ . We do this for all such components  $X$  diffeomorphic to  $S^2 \times I$ . The resulting collection  $\mathcal{T}_2(t)$  is an  $H$ -invariant sub-collection of  $\mathcal{T}_1(t)$  and consists only of tori. Each component of cutting  $M_t(w, -)$  open along this new collection is obtained from a component of cutting  $M_t(w, -)$  open along  $\mathcal{T}_1(t)$  by attaching 3-balls along all the boundary 2-spheres. Thus, the components of the result of cutting  $M_t(w, -)$  open along  $\mathcal{T}_2(t)$  are total spaces of Seifert fibrations possibly with one or more solid tori added so as to kill the homotopy class of the generic fiber, as well the components of  $W'_1(t)$  of the following types:

1. a  $T^2$ -bundle over the circle or the union of two twisted  $I$ -bundles over the Klein bottle glued together along their boundary,



2.  $T^2 \times I$ , or
3. a twisted  $I$ -bundle over the Klein bottle or a solid torus.

**Lemma 13.30.** *Without loss of generality we can assume that, in addition to the above description of the components of the result of cutting  $M_t(w, -)$  open along the  $H$ -invariant family of tori  $\mathcal{T}_2(t)$ , the following hold. Every component of  $\mathcal{T}_2(t)$  is a 2-torus and each component of  $\mathcal{T}_2(t)$  is either incompressible in  $M_t(w, -)$  or is compressible on exactly one side and bounds a solid torus in  $M_t(w, -)$ , a solid torus that contains no component of  $\mathcal{T}_2(t)$  in its interior.*

*Proof.* Consider all components of  $\mathcal{T}_2(t)$  that are compressible tori. For any such torus either bounds a solid torus in  $M_t(w, -)$  or bounds a non-trivial knot complement in  $M_t(w, -)$  and is compressible on the other side. Suppose that there are 2-torus components of  $\mathcal{T}_2(t)$  that are compressible in  $M_t(w, -)$  yet do not bound solid tori. We take the collection of these components whose knot complements are minimal, in the sense that they are not properly contain other knot complements bounded by a component  $\mathcal{T}_2(t)$ . The submanifolds that these tori bound form an  $H$ -invariant family of disjoint knot complements in  $M_t(w, -)$ . We replace each of these minimal knot complements by a solid torus in such a way that kernel on first homology of the inclusion of the 2-torus boundary into the solid torus is the same as the kernel of the inclusion of the 2-torus into the knot complement. This this collection of subgroups of first homology is stabilized by the  $H$ -action, and consequently the  $H$ -action on this collection of tori extends to an  $H$ -action on the solid tori.

**Claim 13.31.** *Replacing the knot complements with solid tori in this manner and extending the actions over the solid tori does not change the ambient manifold up to  $H$ -equivariant diffeomorphism.*

*Proof.* Let us consider the operation restricted to one such torus component  $T$  of  $\mathcal{T}_2(t)$  and restrict to the stabilizer  $H_T$  of that component. Let  $K$  be the knot complement bounded by  $T$  and let  $S$  be the solid torus that we add to  $T$ . Since there is a product neighborhood of  $T$  on which the action of  $H_T$  is the product of a linear action on  $T$  with the trivial action on  $I$ , we can deform the metric in an  $H_T$ -equivariant fashion until  $T$  is totally geodesic. There is a compressing disk  $D \subset X = M_t(w, -) \setminus \text{int}K$  for  $T$ . A regular neighborhood  $P$  of  $T \cup D$  in  $X$  is the complement of a 3-ball in a solid torus, and the union  $K \cup_T P$  is a manifold with 2-sphere boundary and cyclic fundamental group. Since  $M_t$  is aspherical, it must be the case that this 3-manifold is the 3-ball. In particular, the kernels of  $H_1(T) \rightarrow H_1(K)$  and  $H_1(T) \rightarrow H_1(P)$  together generate  $H_1(T; \mathbb{Z})$ . Now consider the union of  $S \cup_T P$ . It is a union of a solid torus and a punctured solid torus and it has trivial first homology. Thus, it is diffeomorphic to a 3-ball. Hence  $K \cup_T P$  and  $S \cup_T P$  are diffeomorphic, and as a result  $K \cup_T X$  and  $S \cup_T X$  are also diffeomorphic. Now perform this operation simultaneously on all components of  $\mathcal{T}_2(t)$  that are compressible but do not bound solid tori. This argument shows that the result of replacing the knot complements by solid tori yields a manifold diffeomorphic to  $M_t(w, -)$ .

Now let us consider the action of the stabilizer  $H_T$  of  $T$ . It stabilizes the kernel of  $\pi_1(T) \rightarrow \pi_1(X)$  and hence, by the equivariant version of Dehn's Lemma and the Loop Theorem ([18]) there is an  $H_T$ -invariant family of disjointly embedded 2-disks  $\mathcal{D}$  in  $X$  with the property that the boundary of each generates the kernel of  $\pi_1(T) \rightarrow \pi_1(X)$ . Let  $k$  be the number of disks in the family  $\mathcal{D}$ . In this way we create an action of  $H$  on a manifold diffeomorphic to  $M_t(w, -)$ . We still need to establish is that the diffeomorphism between the manifolds can be chosen to be  $H$ -equivariant.

Let  $\tilde{P}$  be a regular neighborhood of  $T \cup \mathcal{D}$ . This is the complement of a disjoint collection of  $k$  three-balls in a solid torus. Thus,  $S \cup \tilde{P}$  is the complement of  $k$  three-balls in  $S^3$ . Since  $M_t(w, -)$  is acyclic, each of the 2-sphere boundary components of  $\tilde{P}$  separates  $M_t(w, -)$ . The group  $H_K$  permutes the complementary components of  $M_t(w, -) \setminus (S \cup \tilde{P})$ . If  $k > 1$ , this contradicts the fact that  $M_t(w, -)$  is acyclic. This shows that there is a single compressing disk  $(D, \partial D) \subset (X, T)$  that is  $H_T$ -invariant. Doing this for each such component of  $\mathcal{T}_2(t)$  that is compressible and bounds a knot complement, we find an  $H$ -invariant family of  $\mathcal{P} = \{P_1, \dots, P_k\}$  with each  $P_i$  being diffeomorphic to a complement of a 3-ball in a solid torus. The union of  $\mathcal{P}$  and the collection of solid tori  $\{S_1, \dots, S_k\}$  added to these components is then an  $H$ -equivariant family  $\mathcal{B}$  of 3-balls in the newly constructed manifold. Let  $Y$  be its complement. Then  $Y$  is identified with  $M_t(w, -) \setminus \cup_{i=1}^k (K_i \cup P_i)$  and the  $H$ -actions match under these identifications. Since we have already established the Generalized Smith Conjecture, it follows that there is an  $H$ -equivariant diffeomorphism from  $\mathcal{B}$  to  $\cup_{i=1}^k K_i \cup P_i$  extending the given identifications on the boundary. This completes the proof of the claim.  $\square$

This allows us to assume that if a component of  $\mathcal{T}_2(t)$  is a compressible 2-torus then it bounds a solid torus in  $M_t(w, -)$ . Since each component of  $M_t$  is aspherical, such a torus bounds a solid torus on only one side. We take a maximal collection of such solid tori, maximal in the sense that every solid torus bounded by an element of  $\mathcal{T}_2(t)$  is contained in one of these, and none of these is properly contained in a larger one. This is an  $H$ -invariant subset. We remove from  $\mathcal{T}_2(t)$  all components contained in the interiors of this collection of solid tori. This produces a new  $H$ -invariant family of tori, which we call  $\mathcal{T}'(t)$ , with the property that if  $T^2$  is a component of  $\mathcal{T}'(t)$  that is compressible in  $M_t(w, -)$ , then it bounds a solid torus which is one of the components of cutting  $M_t(w, -)$  open along  $\mathcal{T}'(t)$ .  $\square$

Each component of the result of cutting  $M_t(w, -)$  open along  $\mathcal{T}'(t)$  is of one of the following types:

1. a  $T^2$ -bundle over the circle or the union of two twisted  $I$ -bundles over the Klein bottle glued together along their boundary,
2. a solid torus,
3. a Seifert fibration not diffeomorphic to a solid torus,
4. the union of a Seifert fibration and one or more solid tori glued along boundary components of the Seifert fibration in such a way as to kill the generic fiber.

**Claim 13.32.** *No component of the result of cutting  $M_t(w, -)$  open along  $\mathcal{T}'(t)$  is the union of a Seifert fibration with one or more solid tori attached so as to kill the homotopy class of the generic fiber of the Seifert fibration structure.*

*Proof.* Suppose that there is such a component  $Y$  which is the union of a Seifert fibration with total space  $Z$  and a collection of one or more solid tori added along the boundary components of  $Z$ . The boundary components of  $Y$  are boundary components of  $Z$ . Since there is at least one solid torus in  $Y \setminus Z$  that kills the generic fiber of the Seifert fibration structure on  $Z$ , each of the boundary components of  $Y$  is compressible in  $Y$ . Since the elements of  $\mathcal{T}'(t)$  are compressible on exactly one side, that side being a solid torus component of the result of cutting  $M_t(w, -)$  open along  $\mathcal{T}'(t)$  it follows that  $Y$  is closed. Thus,  $\pi_1(Y)$  is identified with the quotient of  $\pi_1(Z)$  by the normal subgroup generated by the generic fiber of the Seifert fibration. This means that  $\pi_1(Y)$  is isomorphic to the orbifold fundamental group of a 2-dimensional orbifold, which contradicts the fact that  $Y$  is aspherical.  $\square$

A similar argument shows the following:

**Claim 13.33.** *If  $Y$  is a component of the result of cutting  $M_t(w, -)$  open along  $\mathcal{T}'(t)$  which is the total space of a Seifert fibration and which is not diffeomorphic to a solid torus, and if  $T$  is a component of  $\partial Y$  that is compressible in  $M_t(w, -)$ , then the Seifert fibration structure on  $Y$  extends over solid torus bounded by  $T^2$ .*

*Proof.* The Seifert fibration structure of  $Y$  will extend over the solid torus unless the homotopy class in  $Y$  of the boundary of the non-trivial disk in the solid torus  $\tau$  that  $T$  bounds is the same as that of the generic fiber. Suppose that this were the case. If  $Y$  has a boundary component  $T'$  distinct from  $T$ , then  $T'$  is compressible on the side containing  $Y$ , which is a contradiction as before. If  $\partial Y = T$ , then the fundamental group of the closed 3-manifold  $Y \cup \tau$  is the same as that of the quotient 2-dimensional orbifold for the Seifert fibration on  $Y$ . This means that  $Y$  is not aspherical.  $\square$

Now we remove from the collection  $\mathcal{T}'(t)$  all tori that bound solid tori and we call the resulting  $H$ -invariant family  $\mathcal{T}(t)$ . We let  $\mathcal{Y}(t)$  be the result of cutting  $M_t(w, -)$  open along  $\mathcal{T}(t)$ . By what we just established, each component  $Y$  of  $\mathcal{Y}(t)$  is a Seifert fibration not diffeomorphic to solid tori or is diffeomorphic to  $T^2$ -bundle over the interval or the circle, a twisted  $I$ -bundle over the Klein bottle, or the union of two twisted  $I$ -bundles over the Klein bottle. It follows from the fact that no component  $Y$  is a solid torus that the boundary components of  $Y$  are incompressible in  $Y$ . It follows by Van Kampen's theorem that each 2-torus in  $\mathcal{T}(t)$  is incompressible in  $M_t(w, -)$ . This completes the proof of Proposition 13.29.

To complete the proof of Theorem 13.4 it suffices to find a collection  $\widehat{\mathcal{T}}(t)$  of incompressible tori and Klein bottles in  $M_t$  with the properties stated in that theorem for the collection  $\widehat{\mathcal{T}}(P)$ . Since the boundary components of  $M_t(w, -)$  are incompressible tori in  $M_t$ , it follows that  $\widehat{\mathcal{T}}_1(t) = \mathcal{T}(t) \amalg \partial M_t(w, -)$  is an  $H$ -invariant family of incompressible tori in  $M_t$ . The components of the result,  $\widehat{\mathcal{Y}}_1(t)$ , of cutting

$M_t$  open along  $\widehat{\mathcal{T}}(t)$  are of the types listed in Proposition 13.29 and components diffeomorphic to truncations along horospherical tori of complete hyperbolic manifolds of finite volume. The base orbifold of a compact Seifert fibration with incompressible boundary either has an interior that admits a locally homogeneous metric of finite area or is isomorphic to the annulus, the Möbius band, or the disk with two exceptional points of order 2. If it is orientable, the total space of a Seifert fibration with one of these three exceptional bases is diffeomorphic to  $T^2 \times I$  or the twisted  $I$ -bundle over the Klein bottle. Thus, each component of  $\mathcal{Y}_1(t)$  either has an interior that admits a locally homogeneous Riemannian metric of finite volume or is diffeomorphic to either  $T^2 \times I$  or to the twisted  $I$ -bundle over the Klein bottle. We shall modify the tori  $\widehat{\mathcal{T}}_1(t) \subset M_t$  so as to remove the components of  $\widehat{\mathcal{Y}}(t)$  that are diffeomorphic to  $T^2 \times I$  or to twisted  $I$ -bundles over the Klein bottle. First we remove from  $\widehat{\mathcal{T}}_1(t)$  all components that bound either  $T^2 \times I$  or a twisted  $I$ -bundle over the Klein bottle on each side. This replaces  $\widehat{\mathcal{T}}_1(t)$  by a smaller  $H$ -invariant collection of incompressible tori denoted  $\widehat{\mathcal{T}}_2(t)$ . We denote by  $\mathcal{Y}_2(t)$  the result of cutting  $M_t$  open along  $\widehat{\mathcal{T}}_2(t)$ . The components of  $\mathcal{Y}_2(t)$  are those listed in Proposition 13.29 and truncated hyperbolic manifolds of finite volume. Furthermore, any component of  $\widehat{\mathcal{Y}}_2(t)$  that is diffeomorphic to  $T^2 \times I$  is bordered on each side by a component that is not diffeomorphic to either  $T^2 \times I$  nor to the twisted  $I$ -bundle over the Klein bottle. For each component of  $\mathcal{Y}_2(t)$  diffeomorphic to  $T^2 \times I$  we replace the two boundary tori of that component by the middle torus in that component. By ‘middle torus’ we mean a torus in the interior of the component, parallel to each boundary component, that is invariant under the stabilizer of this component. We can do this in every product component of  $\mathcal{Y}_2(t)$  so as to produce a new  $H$ -invariant family of incompressible tori  $\widehat{\mathcal{T}}_3(t)$  with the property that the components of the result,  $\widehat{\mathcal{Y}}_3(t)$ , of cutting  $M_t$  open along this new collection are of the types listed in Proposition 13.29 and truncated hyperbolic manifolds of finite volume, and furthermore, no component of  $\widehat{\mathcal{Y}}_3(t)$  is diffeomorphic to  $T^2 \times I$ . Next, we consider the components of  $\widehat{\mathcal{Y}}_3(t)$  that are diffeomorphic to twisted  $I$ -bundles over the Klein bottle. Each such component contains a Klein bottle isotopic to the zero section that is invariant under the stabilizer of that component. We replace the boundary torus of such components by these invariant Klein bottles. Again we can do this in an  $H$ -invariant way, resulting in an  $H$ -invariant family of incompressible tori and Klein bottles  $\widehat{\mathcal{T}}(t)$  with the property that the components,  $\mathcal{Y}(t)$ , of the result of cutting  $M_t$  open along these surfaces are all of the types listed in Proposition 13.29 and truncated hyperbolic manifolds of finite volume, and furthermore, so that no component of  $\mathcal{Y}(t)$  is diffeomorphic to  $T^2 \times I$  or to a twisted  $I$ -bundle over the Klein bottle. This shows:

**Proposition 13.34.** *For every  $t$  sufficiently large, there is an  $H$ -invariant family of incompressible tori and Klein bottles,  $\widehat{\mathcal{T}}(t)$ , in  $M_t$  such that the components of the result of cutting  $M_t$  open along this family are of the types listed in Proposition 13.29 or are truncations of complete hyperbolic 3-manifolds of finite volume along horospherical tori. Furthermore, no component of the result of cutting  $M_t$  open along  $\widehat{\mathcal{T}}(t)$  is diffeomorphic to either  $T^2 \times I$  or a twisted  $I$ -bundle over the Klein bottle.*

By Lemma 5.1 it follows that we have an  $H$ -invariant family  $\widehat{\mathcal{T}}(t)$  of incompress-

ible tori and Klein bottles with the property that every component of  $M_t \setminus \widehat{\mathcal{T}}(t)$  has a complete homogeneous metric of finite volume of Solv, Nil, Flat,  $\mathbb{H}^2 \times \mathbb{R}$ ,  $\widehat{PSL}_2(\mathbb{R})$  or hyperbolic type. It remains to show that these locally homogeneous metrics of finite volume can be chosen to be  $H$ -invariant, or equivalently that each component has a locally homogeneous metric of finite volume invariant under the stabilizer of that component. We have already established by limiting arguments that disjoint union of the the hyperbolic metrics on the truncated hyperbolic components are invariant under  $H$  and that the flat metrics on the flat components are  $H$ -invariant. All the other cases follow from [17]. (There only the case of finite groups is considered there but the same arguments work for compact groups.)

## References

- [1] B. N. Apanasov, ‘Conformal Geometry of Discrete Groups and Manifolds,’ in Expositions in Mathematics, **vol. 32**, Walter de Gruyter, 2000, Berlin-New York, 523 pages.
- [2] L. Bessières, G. Besson, M. Boileau, S. Maillot, J. Porti ‘Weak collapsing and geometrisation of aspherical 3-manifolds,’ *preprint* arXiv:math/0706.2065, 2007.
- [3] Y. Burago, M. Gromov and G. Perelman, ‘A. D. Aleksandrov spaces with curvatures bounded below,’ (Russian. Russian summary) Uspekhi Mat. Nauk 47 (1992), no. 2(284), 3–51, 222; translation in Russian Math. Surveys 47 (1992), no. 2, 1–58.
- [4] H-D. Cao and X-P. Zhu, ‘A complete proof of the Poincaré and geometrization conjectures—application of the Hamilton-Perelman theory of the Ricci flow,’ Asian J. Math. 10 (2006), no. 2, 165–492.
- [5] T. Colding and W. Minicozzi, ‘Estimates for the extinction time for the Ricci flow on certain 3-manifolds and a question of Perelman,’ J. Amer. Math. Soc. 18 (2005), no. 3, 561–569 arXiv:math/0308090.
- [6] J. Dinkelbach and B. Leeb, ‘Equivariant Ricci flow with surgery and applications to finite group actions on geometric 3-manifolds,’ *preprint* arXiv:math/0801.0803, 2008.
- [7] K. Fukaya and T. Yamaguchi, ‘The fundamental groups of almosts nonnegatively curved manifolds,’ Ann. of Math. (2) **136** (1992), 253 – 333.
- [8] D. Gabai, R. Meyerhopf, and P. Milley, ‘Minimum volume cusped hyperbolic 3-manifolds,’ *preprint* arXiv:0705.4325
- [9] Gromov, M. “Metric structures for Riemannian and non-Riemannian spaces,” Birkhäuser, Boston, 1985.
- [10] K. Grove and Karsher, How to conjugate  $C^1$ -close group actions. Math. Z. **132** (1973), 11–20.

- [11] R. Hamilton, ‘Three manifolds of positive Ricci curvature,’ *Jour. Diff. Geom.* **17** (1982), 255–306.
- [12] R. Hamilton, ‘Non-singular solutions of the Ricci flow on three-manifolds,’ *Comm. Anal. Geom.* **7** (1999), 695–729.
- [13] R. Hamilton, ‘Four-manifolds with positive isotropic curvature,’ *Comm. Anal. Geom.*, **5** (1997), 1–92.
- [14] V. Kapovitch, ‘Perelman’s Stability Theorem,’ *preprint* arXiv:math/0703002, 2007.
- [15] B. Kleiner and J. Lott, ‘Notes on Perelman’s paper,’ URL: <http://www.math.lsa.umich.edu/re-search/ricciflow/perelman.html>.
- [16] B. Kleiner and J. Lott, ‘Geometrization of Three-Dimensional Orbifolds via Ricci Flow,’ *preprint* arXiv:math.DG/1101.3733v2, 2011.
- [17] W. Meeks and P. Scott, ‘Finite group actions on 3-manifolds,’ *Invent. Math.* **86** (1986), 287–346.
- [18] W. Meeks and S-T. Yau, ‘Topology of three-dimensional manifolds and the embedding problems in minimal surface theory,’ *Ann. of Math* **112** (1980), 441 – 485.
- [19] Milka, A.D., *Ukrain. Geometr. Sb. Vyp* 4(1967)
- [20] J. Milnor, ‘A unique decomposition theorem for 3-manifolds,’ *Amer. J. Math.* Vol. **84** (1962), 1–7.
- [21] J. Morgan and G. Tian, ‘Ricci Flow and the Poincaré Conjecture,’ *Clay Mathematics Monographs*, vol. **3**, American Math. Society, Providence 521 pages, 2007.
- [22] G. D. Mostow, ‘Quasi-conformal mappings in n-space and the rigidity of the hyperbolic space forms,’ *Publ. Math. IHES* 34 (1968) 53104.
- [23] P. Orlik, ‘Seifert manifolds,’ *LMN Vol. 291*, Springer-Verlag, Berlin-New york, 1972, 155pp.
- [24] J-P. Otal, ‘The hyperbolization theorem for fibered 3-manifolds,’ *SMF/AMS Texts and Monographs* **7**, translated from the 1996 French original by Leslie D. Kay, American Mathematical Society, Providence, 2001, xiv+126 pages.
- [25] J-P. Otal, ‘Thurston’s hyperbolization of Haken manifolds,’ *Surveys in differential geometry*, Vol. III Cambridge, MA, 77–194, Int. Press, Boston, MA, 1998.
- [26] G. Perelman, ‘The entropy formula for Ricci flow and its geometric applications,’ *preprint* arXiv:math.DG/0211159, 2002.

- [27] G. Perelman, ‘Ricci flow with surgery on three-manifolds,’ preprint arXiv.math.DG/0303109, 2003.
- [28] G. Perelman, ‘Finite extinction time for the solutions to the Ricci flow on certain three-manifolds,’ *preprint* arXiv:math/0307245, 2003.
- [29] P. Petersen, ‘Riemannian Geometry, 2nd edition,’ Graduate Texts in Mathematics 171, Springer-Verlag, Berlin-New York, 2006, 401 pp.
- [30] P. Scott, ‘The geometries of 3-manifolds,’ Bull. London Math. Soc. **15** (5), 401 – 487, 1983.
- [31] H. Seifert, ‘Topologie Dreidimensionaler Gefaseter Räume,’ Acta Mat. **60** (1933), 147 – 238.
- [32] T. Shioya and T. Yamaguchi, ‘Collapsing three-manifolds under a lower curvature bound,’ J. Differential Geom. **56** (2000), 1 – 66.
- [33] T. Shioya and T. Yamaguchi, ‘Volume collapsed three-manifolds with a lower curvature bound,’ Math. Ann. **333** (2005), no. 1, 131–155.
- [34] V. A. Toponogov, ‘The metric structure of Riemannian spaces with nonnegative curvature which contains straight lines,’ in Translations of the AMS Series 2 Vol. 70 1968, 225-239.
- [35] F. Waldhausen, ‘Eine Klasse von 3-dimensionalen Mannigfaltigkeiten: I, II’ (German) Invent. Math. **3** (1967), 308–333; *ibid.* **4** (1967) 87–117.